









## Original Research Report

## DIFFERENCES IN AEROBIC CAPACITY AND RUNNING SPEED ACROSS VARIOUS SOMATOTYPE STRUCTURES AND BODY FAT COMPOSITIONS AMONG PROFESSIONAL FOOTBALL ATHLETES IN INDONESIA

Dhoni Akbar Ghozali<sup>1,2\*</sup> , Muhammad Syauqi Ridhallah<sup>2</sup>, Syania Shabrina<sup>3</sup>, Ahmad Isnaini Shidqi Nurhani<sup>3</sup>, Yunia Hastami<sup>2</sup> , Dwi Rahayu<sup>3</sup>, Lukman Aryoseto<sup>3</sup>, Selfi Handayani<sup>2</sup> , Siti Munawaroh<sup>2</sup> , Nanang Wiyono<sup>2</sup> , Agus Sugeng Riyanto<sup>4</sup>, Enrico Ananda Budiono<sup>3</sup> , Annisa Aghnia Rahma<sup>3</sup> , Muhana Fawwazy Ilyas<sup>2</sup> 

<sup>1</sup>Department of Sports Medicine, Faculty of Medicine, Universitas Sebelas Maret, Surakarta, Indonesia

<sup>2</sup>Department of Anatomy and Embryology, Faculty of Medicine, Universitas Sebelas Maret, Surakarta, Indonesia

<sup>3</sup>Faculty of Medicine, Universitas Sebelas Maret, Surakarta, Indonesia

<sup>4</sup>Bhayangkara Football Club, Bekasi, Indonesia

## ABSTRACT

This study investigates the effects of somatotype structure and body fat composition on aerobic capacity and running speed. This study used an analytical observational design with a cross-sectional approach. The subjects were 27 professional football athletes from Bhayangkara Football Club in Bekasi, Indonesia. This study measured several variables, namely age, playing position, somatotype structure assessed using the Somatotype Rating Form and Heath-Carter Somatochart, body fat composition measured using the Brozek and Siri formulas, aerobic capacities determined by the maximum rate of oxygen consumption (VO<sub>2</sub> max) through the Yo-Yo Intermittent Recovery Test Level 2, and running speed over a 30 m distance. The data were presented as mean ± standard deviation (SD), frequency (n), and percentage. The Shapiro-Wilk test was used to determine the normality of the data distribution. The statistical analyses included one-way analysis of variance (ANOVA) followed by the post-hoc least significant difference (LSD) test, the Kruskal-Wallis test followed by the post-hoc Mann-Whitney test, as well as the independent t-test, the Mann-Whitney test, and Pearson's or Spearman's correlation tests. A p-value of less than 0.05 was considered statistically significant. The athletes predominantly exhibited a mesomorph-endomorph somatotype (88.9%). Significant correlations were found between mesomorph rating and running speed ( $r = -0.548$ ;  $p = 0.003$ ), body fat composition and aerobic capacity ( $r = -0.448$ ;  $p = 0.019$ ), as well as age and aerobic capacity ( $r = -0.515$ ;  $p = 0.006$ ). Significant differences in aerobic capacity were observed among various age groups ( $p = 0.031$ ). There were also differences in body fat composition ( $p = 0.003$ ) and running speed ( $p = 0.036$ ) between the two somatotypes. These findings underscore the importance of individualized training and conditioning programs that take into account the unique body compositions and ages of athletes.

**Keywords:** Aerobic capacity; athletes; body fat composition; football; healthy lifestyle

\***Correspondence:** Dhoni Akbar Ghozali, Department of Anatomy and Embryology, Faculty of Medicine, Universitas Sebelas Maret, Surakarta, Indonesia. E-mail: [dhoniakbar@staff.uns.ac.id](mailto:dhoniakbar@staff.uns.ac.id)

## Article history

• Submitted 05/03/2024 • Revised 18/04/2024 • Accepted 09/05/2024 • Published 10/06/2024

**How to cite:** Ghozali DA, Ridhallah MS, Shabrina S, et al (2024). Differences in Aerobic Capacity and Running Speed Across Various Somatotype Structures and Body Fat Compositions among Professional Football Athletes in Indonesia. *Folia Medica Indonesiana* 60 (2), 103-110. doi: <https://doi.org/10.20473/fmi.v60i2.55757>



Copyright: © 2024 Folia Medica Indonesiana.

This is an open-access article distributed under the terms of the Creative Commons Attribution

License as stated in <https://creativecommons.org/licenses/by-nc-sa/4.0/deed.id>.

pISSN:2355-8393, eISSN: 2599-056x

## Highlights:

1. This study emphasizes the significance of understanding professional football players' various somatotype structures and body fat compositions as an important contribution to the area of sports science.
2. This study lays the groundwork for future research to better understand the intricate interactions between somatotype structures, body composition, and athletic performance.
3. Since a one-size-fits-all approach may not be successful, this study advocates for tailored evaluations of training regimens to cater to athletes' unique requirements and skills, which may lead to higher overall performance.

## INTRODUCTION

Football involves intricate movements such as sprinting, jumping, ball control, and rapid acceleration and deceleration. Athletes have to maintain peak physical condition to optimize their performance. The physical attributes of football players encompass strength, endurance, speed, flexibility, muscle mobility, fundamental motor skills, coordination, agility, and responsiveness (Akenhead 2014, Scheunemann 2014). Sprinting is recognized as an essential skill for scoring goals and determining the outcome of a game. Previous studies have underscored the significant influence of an athlete's somatotype on running speed and endurance (Marangoz 2018, Ryan-Stewart et al. 2018). Somatotype refers to the classification of human body types or compositions based on anthropometric characteristics, which are further categorized into mesomorphy, ectomorphy, and endomorphy types. It has been demonstrated that ectomorph and mesomorph types of amateur soccer players exhibit faster running speeds compared to those of endomorph-type players (Primasoni 2012, Anggitasari et al. 2019).

Football players must possess high levels of endurance, a paramount attribute given the prolonged and strenuous nature of football matches. Aerobic endurance is the ability of the heart, lungs, and blood vessels to use oxygen as energy to carry out activities for a long period of time. VO<sub>2</sub> max, which refers to the maximum rate of oxygen consumption, is a common way to gauge aerobic endurance. An average football player has a VO<sub>2</sub> max of 55–65 mL/kg/min (Akenhead 2014, Modric et al. 2020). Young (2020) demonstrated that genetics, cardiovascular system, training habits, lifestyles, and body compositions can impact a player's VO<sub>2</sub> max. Body composition is an important factor that must be considered when determining VO<sub>2</sub> max. Body composition includes the amount and distribution of fat, muscle, and bone tissue. An excess of adipose tissue can impede the adequate circulation of blood throughout the body, diminishing the delivery of oxygen to muscle cells. Additionally, extra body fat can lessen pulmonary strain, disturbing the exchange of oxygen and carbon dioxide. It has been found that futsal players with a lower body fat percentage exhibit higher VO<sub>2</sub> max or aerobic endurance (Jayanti et al. 2019, Damayanti & Adriani 2021). Excess fat can hasten tiredness, which affects an athlete's ability to maintain heart and lung endurance.

Numerous researchers, such as Cinarli et al. (2022) and Marangoz (2018), have examined the relationship between athletes' somatotype and running speed. However, there is a scarcity of research concerning professional football players.

To our knowledge, the existing research on the somatotypes of professional football players still primarily focuses on the variations among them according to the playing positions, whether it is a striker, midfielder, defender, or goalkeeper. Furthermore, a noticeable gap exists in the literature, as no prior studies have endeavored to elucidate the impact of body fat percentage on maximal oxygen uptake (VO<sub>2</sub> max). This study aimed to fill this void by scrutinizing the influence of somatotype structures and body fat composition on professional football athletes' aerobic capacities and running speed.

## MATERIALS AND METHODS

This study employed an analytical observational design and specifically utilized a cross-sectional approach (Probandari et al. 2020). The study was conducted in November 2021 at the home base of Bhayangkara Football Club (FC) in Bekasi and the Universitas Sebelas Maret Stadium in Surakarta, Indonesia. This study obtained ethical clearance from the Health Research Ethics Committee of Dr. Moewardi Regional General Hospital, Surakarta, Indonesia, under reference No. 895/IX/HREC/2021 on 21/9/2021. This study included professional football athletes from Bhayangkara FC who played in the Indonesian League 1 during the 2021/2022 season. The criteria for inclusion in this study were limited to male players between the ages of 18 and 35 who followed a routine training program provided by the coaching team. Meanwhile, the exclusion criteria were players who were sick or injured and those training for the Indonesian national team at the time, so they could not participate in this research until its completion. Additionally, athletes who refused to participate in this study were also excluded. Potential bias in this study was minimized through total sampling, standardized measurements using calibrated instruments, and ethical clearance. This research used a total sampling technique (Sumardiyono et al. 2020), which resulted in the selection of a sample consisting of 27 athletes who agreed to participate and provided informed consent.

This study measured several variables of the players, including age, playing position (i.e., goalkeeper, back, midfielder, and forward), somatotype structures, body fat composition, aerobic capacity, and running speed. The somatotype of all subjects was assessed by measuring their body height, body weight, skinfold (i.e., triceps, subscapular, suprascapular, and calf), bone diameter (i.e., width of elbow and knee joints), and muscle size (i.e., biceps and calf circumference). Each measurement was performed three times prior to calculating the average values. The calculation results were then

entered into the Somatotype Rating Form, and the final results were calculated on the Heath-Carter Somatochart according to the dominant somatoplots (Carter & Heath 1971). The somatoplot results were divided into four categories depending on the chief points: central, endomorph, mesomorph, and ectomorph.

The body fat percentage was measured according to the skinfold thickness at seven locations: the triceps, subscapular, mid-axilla, pectoral, abdomen, supra-iliac, and thigh. The average was calculated after conducting the measurements three times using a slim guide caliper. Data from the skinfold thickness measurements were entered into the seven-location Jackson-Pollock formula to determine body density values (Jackson & Pollock 1978). The body density values were then entered into the Brozek and Siri formulas to obtain body fat percentages (Brožek et al. 1963, Siri 1993). Following that, the aerobic capacities were assessed by measuring the VO<sub>2</sub> max using the Yo-Yo Intermittent Recovery Test Level 2 (Bangsbo et al. 2008). The measurements were carried out in the field using a 25-meter track for each test participant. Lastly, the running speed was determined by observing the time a participant took to run quickly over a distance of 30 m. The running speed was calculated by dividing the running distance by the time taken to finish the run.

The data obtained were analyzed using IBM SPSS Statistics for Windows, version 27.0 (IBM Corp., Armonk, N.Y., USA). The data were reported in the form of mean  $\pm$  standard deviation (SD), frequency (n), and percentage. The normal distribution of the data was determined using the Shapiro-Wilk test. In the statistical analyses, the one-way analysis of variance (ANOVA) followed by the post-hoc least significant difference (LSD) test were employed. Furthermore, we employed the Kruskal-Wallis test followed by the post-hoc Mann-Whitney test, as well as the independent t-test, the Mann-Whitney test, and Pearson's or Spearman's correlation tests. Finally, we conducted a path analysis by utilizing multiple linear regression. The statistical significance was indicated by a value of  $p < 0.050$  (Sumardiyo et al. 2020).

## RESULTS

A total of 27 professional football athletes were involved in this study. The study subjects had a mean age of  $26.519 \pm 4.611$  years and a mean body mass index (BMI) of  $23.258 \pm 1.555$  kg/m<sup>2</sup>. The majority of the subjects exhibited a somatotype structure that was predominantly mesomorph-endomorph (88.9%). Furthermore, the mean body fat composition was  $9.461 \pm 2.381\%$  when measured using the Brozek formula and  $7.949 \pm 2.622\%$

according to the Siri formula. The subjects had an average aerobic capacity (VO<sub>2</sub> max) of  $56.467 \pm 3.616$  and a running speed of  $26.300 \pm 1.213$  km/hour. Table 1 presents the overall characteristics of the sample.

Table 1. Characteristics of the entire study participants (n=27).

Characteristic	Mean $\pm$ SD	n (%)
Age [years]	26.519 $\pm$ 4.611	
18–20		4 (14.8%)
21–25		7 (25.9%)
26–30		9 (33.3%)
31–35		7 (25.9%)
Body mass index [kg/m <sup>2</sup> ]	23.258 $\pm$ 1.555	
Position		
Goal Keeper		3 (11.1%)
Back		10 (37.0%)
Midfielder		11 (40.7%)
Forward		3 (11.1%)
Somatotype structures		
Mesomorph-endomorph		24 (88.9%)
Mesomorph-ectomorph		3 (11.1%)
Somatotype rating		
Endomorphy rating	3.389 $\pm$ 0.725	
Mesomorphy rating	4.796 $\pm$ 0.880	
Ectomorphy rating	2.278 $\pm$ 0.641	
Body fat compositions (Brozek formula)	9.461 $\pm$ 2.381	
Body fat compositions (Siri formula)	7.949 $\pm$ 2.622	
Aerobic capacity (VO <sub>2</sub> max)	56.467 $\pm$ 3.616	
Running speed [km/h]	26.300 $\pm$ 1.213	

Legend: Nominal data are presented as frequencies and percentages. Numerical data are presented as means  $\pm$  standard deviations (SD).

There were differences in aerobic capacity (VO<sub>2</sub> max) among various age groups ( $p=0.031$ ). Significant differences were observed between the 18–20 and 31–35 age groups (mean difference= $5.897$ ;  $p=0.017$ ) and between the 26–30 and 31–35 age groups (mean difference= $4.204$ ;  $p=0.041$ ). However, the 18-20 and 26–30 age groups had a higher average than the 31–35 age groups. Apart from that, there were also significant differences in body fat composition measured using the Brozek formula (mean difference= $4.113$ ;  $p=.003$ ) as well as running speed (mean difference= $1.536$ ;  $p=0.036$ ) among the study subjects with different somatotype structures. Specifically, the mesomorph-endomorph group exhibited higher body fat composition and running speed than the mesomorph-ectomorph group. All results of differences in body fat composition, aerobic capacity, and running speed across different age groups, playing positions, and somatotype structures are presented in Table 2.

Table 2. Differences in body fat composition, aerobic capacity, and running speed between various age, position, and somatotype structure groups.

	Body fat compositions (Brozek formula)	Body fat compositions (Siri formula)	Aerobic capacity (VO2 max)	Running speed [km/h]
Age (years)	p=0.070 <sup>a</sup>	p=0.376 <sup>a</sup>	p=0.031 <sup>*a</sup>	p=0.252 <sup>a</sup>
18–20 years	7.710±3.349	6.416±2.342	59.200±1.289	26.099±1.643
21–25 years	8.704±1.961	7.474±2.341	56.734±3.496	25.577±1.051
26–30 years	9.451±1.930	8.232±2.099	57.507±3.924	26.750±1.171
31–35 years	11.232±1.945	8.935±3.550	53.303±2.168	26.561±1.038
Positions	p=0.895 <sup>a</sup>	p=0.595 <sup>b</sup>	p=0.318 <sup>a</sup>	p=0.658 <sup>a</sup>
Goal Keeper	10.302±2.129	9.800±2.306	53.123±0.822	25.753±1.228
Back	9.634±2.799	8.148±3.170	56.342±4.128	26.073±1.199
Midfielder	9.128±2.303	7.491±2.242	57.572±2.897	26.559±1.101
Forward	9.268±2.355	7.109±2.473	56.180±5.191	26.655±1.961
Somatotype structures	p=0.003 <sup>*c</sup>	p=0.084 <sup>d</sup>	p=0.747 <sup>c</sup>	p=0.036 <sup>*d</sup>
Mesomorph- endomorph	9.918±2.044	8.226±2.630	56.386±3.751	26.471±1.162
Mesomorph- ectomorph	5.805±1.772	5.732±1.219	57.120±2.721	24.935±0.675

Legends: <sup>a</sup>One-way ANOVA, <sup>b</sup>Kruskall-Wallis test, <sup>c</sup>independent t-test, and <sup>d</sup>Mann-Whitney test were utilized. An asterisk (\*) signifies any statistical significance (p<0.050).

There was a significant correlation between mesomorphy rating and running speed ( $r=-0.548$ ;  $p=0.003$ ). Additionally, negative correlations were observed between body fat composition (measured using the Brozek formula) and aerobic capacity ( $r=-$

$0.448$ ;  $p=0.019$ ) as well as between age and aerobic capacity ( $r=-0.515$ ;  $p=0.006$ ). Meanwhile, there was no relationship found between other variables. All correlation results between variables are presented in [Table 3](#).

Table 3. Correlation between somatotype structure, body fat compositions, aerobic capacity, and running speed.

		Correlation coefficient	p
Endomorphy rating	Aerobic capacity (VO2 max)	-0.243	0.222 <sup>a</sup>
	Running speed	0.239	0.230 <sup>a</sup>
Mesomorphy rating	Aerobic capacity (VO2 max)	0.141	0.484 <sup>b</sup>
	Running speed	-0.548	0.003 <sup>*b</sup>
Ectomorphy rating	Aerobic capacity (VO2 max)	-0.033	0.872 <sup>b</sup>
	Running speed	-0.074	0.713 <sup>b</sup>
Body fat compositions (Brozek formula)	Aerobic capacity (VO2 max)	-0.448	0.019 <sup>*a</sup>
	Running speed	0.230	0.249 <sup>a</sup>
Body fat compositions (Siri formula)	Aerobic capacity (VO2 max)	-0.347	0.076 <sup>b</sup>
	Running speed	0.525	0.525 <sup>b</sup>
Age	Aerobic capacity (VO2 max)	-0.515	0.006 <sup>*a</sup>
	Running speed	0.346	0.077 <sup>a</sup>
Body mass index	Aerobic capacity (VO2 max)	-0.299	0.130 <sup>a</sup>
	Running speed	-0.010	0.960 <sup>a</sup>

Legend: <sup>a</sup>Pearson correlation test and <sup>b</sup>Spearman correlation test were utilized. An asterisk (\*) signifies any statistical significance (p<0.050).

## DISCUSSION

The highest rate at which oxygen can be carried up and used by the body during intense activities is referred to as VO2 max. Aerobic capacity, defined as the highest level of oxygen consumption during intense activities that use large muscle groups, is a measure of aerobic fitness. A spike in VO2 max implies that aerobic capacity has increased. A prior study has established an association between maximum aerobic capacity and improved match

performance and team achievement ([Modric et al., 2020](#)). Enhanced aerobic capacity would enable a player to reach and sustain a higher work rate, thus exerting a more significant influence during a game. Various factors, such as genetics, the cardiovascular system, training habits, lifestyle, and body composition, can impact a player's VO2 max ([Akenhead 2014, Young 2020](#)).

Our research indicated significant differences in aerobic capacity (VO2 max) between various age



groups. Individuals in the age groups of 18–20 and 26–30 years had a higher average aerobic capacity than those in the 31–35 age group. According to prior research, the consequences of aging are particularly identified in the context of fitness (Wickramarachchi et al. 2023). The consistent decrease in VO<sub>2</sub> max, notably appearing at the ages of 20–30 and diminishing by around 10% every decade, demonstrates a significant decline in the capacity to exercise. On top of that, the pace of this reduction intensifies with age. It indicates that the present research's findings are in line with the previous study.

A previous cross-sectional study conducted by King & Lowery (2024) revealed that VO<sub>2</sub> max diminishes nearly 50% on average throughout the ages of 20 and 80. This decline must be attributed to either or both cardiac output and arterial-venous oxygen concentration. Cardiac output and arterial-venous oxygen concentration have been shown to fall by 25% with age. Because stroke volume has been predicted to be preserved throughout life, the vast majority of cardiac output reduction tends to be attributed to the decrease in maximum heart rate. The decline in VO<sub>2</sub> max results from an alteration in the arterial-venous oxygen concentration, which illustrates changes in oxygen efficiency and balance. This implies the critical role of mitochondrial and muscle mass performance. On the other hand, the decrease in cardiac function is accounted for by a variety of mechanisms, including increased circulation afterload, arterial-ventricular load mismatching, diminished intrinsic contractility of the myocardium, disrupted autonomic control, and physical deconditioning (King & Lowery 2024).

This current study discovered significant differences in body fat composition, as measured using the Brozek formula, among football players with varied somatotype structures. Individuals in the mesomorph-endomorph group exhibited higher body fat compositions compared to individuals classified in the other somatotype category. Football players are constantly exposed to extensive physical exercises and matches. Hence, their body composition becomes a significant factor affecting their performances. Body fat composition has been measured using numerous techniques, one of which is the Brozek formula. The formula estimates body density through observations of skinfold thickness. The Brozek formula might represent a more reliable alternative than the Siri formula for measuring body density as a percentage of body fat composition. Additionally, it is widely recognized that somatotype variations obviously impact athletic performance and are considered one of the most significant considerations in athlete recruitment. Somatotype determines a person's physical form on the basis of three elements: endomorphy (related to

adiposity), mesomorphy (related to musculoskeletal development), and ectomorphy (related to linearity or slenderness) (Almeida et al. 2021, Cinarli et al. 2022).

The endomorphy component of somatotype, which is associated with adiposity, exhibited a positive correlation with body fat percentage as determined by the Brozek formula. Individuals with a higher endomorphy component might be more likely to have a greater body fat percentage, reflecting a predisposition towards increased adipose tissue (Galić et al. 2016). On the contrary, the mesomorphy component, which is linked to musculoskeletal development, showed an inverse correlation with body fat percentage. Higher mesomorphy scores might be associated with increased lean muscle mass and decreased body fat, indicating a potential protective effect against adiposity. In addition, higher mesomorphy scores indicate strength and power-to-weight ratio, potentially leading to reduced body fat percentage (Sukanta 2014, Bertuccioli et al. 2022). Individuals with the ectomorphy component, which is related to linearity, might demonstrate lower body fat due to a slender body build. However, this could vary depending on factors such as muscle mass and overall fitness.

Running speed is a fundamental attribute in football since it influences an athlete's ability to execute offensive and defensive maneuvers. Athletes with different somatotype structures may demonstrate varying running speeds due to variations in their body compositions. This research revealed a significant difference in running speed between various somatotype structures, with the mesomorph-endomorph group exhibiting a higher running speed than the mesomorph-ectomorph group. A higher level of mesomorphic features contributes to a larger amount of muscle mass. It may benefit athletes who face significant physical challenges throughout competition and training. In contrast, the endomorphy component showed that having greater body fat mass might be advantageous for absorbing and releasing power, resulting in faster running speed compared to individuals in other somatotype variations (Bujak et al. 2016).

The mesomorphy component is more likely to exhibit a positive correlation with running speed. This is because athletes with higher mesomorphy scores may benefit from increased muscle mass and strength, contributing to improved sprinting abilities. However, this research found a negative correlation between mesomorphy rating and running speed. The negative correlation is consistent with several possible explanations. Individuals with a mesomorphic feature are associated with a higher proportion of muscle mass. While muscle mass

contributes to strength and power, it also adds to body mass (Gutnik et al. 2015). In terms of running, particularly when endurance is necessary, individuals with a larger body mass may require more energy to move. Therefore, they may face challenges in achieving maximum running speeds compared to individuals with a lower body mass.

Individuals exhibiting mesomorphy often have a higher percentage of fast-twitch muscle fibers, which are beneficial for short bursts of power. However, they may also experience fatigue more quickly during prolonged endurance activities such as running (de la Iglesia et al. 2020). According to Sanders et al. (2017), running speed is strongly influenced by aerobic capacity, which is the ability of the cardiovascular system to supply oxygen to muscles during sustained physical activity. While individuals with mesomorphy may prioritize strength training over cardiovascular training, it potentially leads to lower endurance and running speed. However, it is essential to note that other factors, such as training regimens, cardiovascular fitness, or individual variations, could also contribute to running speed.

This study found a correlation between the body fat composition of professional football athletes, as measured using the Brozek formula, and their aerobic capacity (VO<sub>2</sub> max). The negative correlation between body fat composition and aerobic capacity ( $r=-0.448$ ;  $p=0.019$ ) indicated that higher body fat levels are associated with lower aerobic capacity. Aerobic capacity, as determined by VO<sub>2</sub> max, is influenced by various factors, including body composition, cardiovascular system efficiency, and muscle oxygen utilization during exercise. On the other hand, body fat composition is a crucial determinant of overall health, with implications for metabolic function and cardiovascular well-being (Frank et al. 2019, Young 2020). Prior studies conducted by Green et al. (2018) and Littleton & Tulaimat (2017) have identified several mechanisms that contribute to the observed correlation. One of which is the physiological impact of body fat on the cardiovascular and respiratory systems. Excessive body fat can lead to decreased lung function, resulting in reduced lung compliance and impaired efficiency in oxygen uptake and utilization during aerobic activities. Moreover, excessive body fat slows the dynamic responses of the VO<sub>2</sub> max during cycling and delays the phase of vasodilation in muscle contraction. Adipose tissue secretes bioactive substances known as adipokines, which may negatively affect the cardiovascular system, further compromising aerobic capacity as measured by VO<sub>2</sub> max (Clemente-Suárez et al. 2023).

### Strength and limitations

This study highlights the importance of recognizing athletes' diverse somatotype structures and body fat compositions as a valuable contribution to the field of sports science. It provides a foundation for further research to enhance our understanding of the complex interplay between somatotype structures, body composition, and athletic performance. We also consider incorporating individualized assessments to optimize training programs for professional football athletes since a one-size-fits-all approach may not be effective. By doing so, they can better address each athlete's specific needs and capabilities, potentially leading to improved overall performance.

While this study contributes valuable insights into the relationships between somatotype structures, body fat compositions, and physical performance indicators among professional football athletes in Indonesia, several limitations should be acknowledged. Firstly, the cross-sectional design limited the establishment of causality, and longitudinal studies would provide a more comprehensive understanding of the dynamic interactions observed. Furthermore, the study did not consider potential psychological or nutritional aspects that could influence the observed relationships.

### CONCLUSION

Relationships between somatotype structure, body fat composition, and physical performance parameters were found among professional football players in Indonesia. The athletes predominantly had a mesomorph-endomorph somatotype. Significant relationships were found between mesomorph rating and running speed, body fat composition (determined by the Brozek formula) and aerobic capacity (VO<sub>2</sub> max), and between age and aerobic capacity. The aerobic capacities of the athletes differed significantly across various age groups. Their body fat compositions and running speeds also varied depending on their somatotype structures. Therefore, individualized training and conditioning programs are crucial considering the different body compositions and ages of the athletes.

### Acknowledgment

We would like to thank the athletes of the Bhayangkara Football Club, Bekasi, Indonesia, for their hospitality and assistance during the process of gathering data and specimens.

### Conflict of interest

None.

### Ethical consideration

This study received approval from the Health Research Ethics Committee of Dr. Moewardi Regional General Hospital, Surakarta, Indonesia, under reference No. 895/IX/HREC/2021 on 21/9/2021.

### Funding disclosure

None.

### Author contribution

DAG, YH, DR, LA, SH, SM, NW, and ASR contributed to the conception and design of this study. MSR, SS, and AISN collected, assembled, analyzed, and interpreted the data. MFI, EAB, and AAR drafted the article. DAG, YH, DR, LA, SH, SM, NW, and ASR contributed to the critical revision of the article for important intellectual content. All the authors gave their approval to the final article.

### REFERENCES

- Akenhead RM (2014). Examining the physical and physiological demands of elite football (thesis). Northumbria University. Available at: [https://www.researchgate.net/publication/265049750\\_Examining\\_the\\_Physical\\_and\\_Physiological\\_Demands\\_of\\_Elite\\_Football](https://www.researchgate.net/publication/265049750_Examining_the_Physical_and_Physiological_Demands_of_Elite_Football).
- Almeida LG, Numata-Filho ES, Dos-Santos GA, et al (2021). Anthropometric profile and functional performance of capoeira competitors in the world games. *International Journal of Morphology* 39, 969–976. doi: 10.4067/S0717-95022021000400969.
- Anggitasari ED, Dieny FF, Candra A (2019). Correlation of somatotype with physical fitness of football athletes. *Jurnal Keolahragaan* 7, 11–22. doi: 10.21831/jk.v7i1.21188.
- Bangsbo J, Iaia FM, Krstrup P (2008). The yo-yo intermittent recovery test. *Sports Medicine* 38, 37–51. doi: 10.2165/00007256-200838010-00004.
- Bertuccioli A, Sisti D, Amatori S, et al (2022). A new strategy for somatotype assessment using bioimpedance analysis: Stratification according to sex. *Journal of Functional Morphology and Kinesiology* 7, 86. doi: 10.3390/jfmk7040086.
- Brožek J, Grande F, Anderson JT, et al (1963). Densitometric analysis of body composition: Revision of some quantitative assumptions\*. *Annals of the New York Academy of Sciences* 110, 113–40. doi: 10.1111/j.1749-6632.1963.tb17079.x.
- Bujak Z, Gierczuk D, Hubner-Wozniak E, et al (2016). Anthropometric profile and anaerobic capacity of martial arts and combat sports athletes. *Ido Movement for Culture. Journal of Martial Arts Anthropology*. doi: 10.14589/ido.16.2.7.
- Carter JEL, Heath BH (1971). Somatotype methodology and kinesiology research. American Association for Health, Physical Education, and Recreation, Washington, DC, USA. Available at: <https://files.eric.ed.gov/fulltext/ED084219.pdf>.
- Cinarli FS, Buyukcelebi H, Esen O, et al (2022). Does dominant somatotype differentiate performance of jumping and sprinting variables in young healthy adults? *International Journal of Environmental Research and Public Health* 19, 1–11. doi: 10.3390/ijerph191911873.
- Clemente-Suárez VJ, Redondo-Flórez L, Beltrán-Velasco AI, et al (2023). The role of adipokines in health and disease. *Biomedicines* 11, 1–41. doi: 10.3390/biomedicines11051290.
- Damayanti C, Adriani M (2021). Correlation between percentage of body fat with speed and cardiorespiratory endurance among futsal athletes in Surabaya. *Media Gizi Indonesia* 16, 53. doi: 10.20473/mgi.v16i1.53-61.
- Frank AP, de Souza Santos R, Palmer BF, et al (2019). Determinants of body fat distribution in humans may provide insight about obesity-related health risks. *Journal of Lipid Research* 60, 1710–1719. doi: 10.1194/jlr.R086975.
- Galić BS, Pavlica T, Udicki M, et al (2016). Somatotype characteristics of normal-weight and obese women among different metabolic subtypes. *Archives of Endocrinology and Metabolism* 60, 60–65. doi: 10.1590/2359-3997000000159.
- Green S, O'Connor E, Kiely C, et al (2018). Effect of obesity on oxygen uptake and cardiovascular dynamics during whole-body and leg exercise in adult males and females. *Physiological Reports* 6, 1–14. doi: 10.14814/phy2.13705.
- Gutnik B, Zuoza A, Zuozienė I, et al (2015). Body physique and dominant somatotype in elite and low-profile athletes with different specializations. *Medicina (B Aires)* 51, 247–252. doi: 10.1016/j.medic.2015.07.003.
- IBM Corp. 2020. IBM SPSS Statistics for Windows, Version 27.0. Armonk, NY: IBM Corp. Available at: <https://www.ibm.com/support/pages/download-ibm-spss-statistics-27>.
- Jackson AS, Pollock ML (1978). Generalized equations for predicting body density of men. *British Journal of Nutrition* 40, 497–504. doi: 10.1079/BJN19780152.
- Jayanti R, Huldani H, Asnawati A (2019). The relationship of body fat percent and maximum oxygen capacity in prospective hajj pilgrimages. *Homeostasis: Jurnal Mahasiswa Pendidikan Dokter* 2, 87–92. doi: 10.20527/ht.v2i1.432.
- King J, Lowery DR (2024). Physiology, cardiac

- output. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/30648673>.
- de la Iglesia R, Espinosa-Salinas I, Lopez-Silvarrey FJ, et al (2020). A potential endurance algorithm prediction in the field of sports performance. *Frontiers in Genetics*. doi: [10.3389/fgene.2020.00711](https://doi.org/10.3389/fgene.2020.00711).
- Littleton SW, Tulaimat A (2017). The effects of obesity on lung volumes and oxygenation. *Respiratory Medicine* 124, 15–20. doi: [10.1016/j.rmed.2017.01.004](https://doi.org/10.1016/j.rmed.2017.01.004).
- Marangoz I (2018). The determination of the relationship between somatotypes and speed of sub-elite athletes. *Asian Journal of Education and Training* 4, 220–223. doi: [10.20448/journal.522.2018.43.220.223](https://doi.org/10.20448/journal.522.2018.43.220.223).
- Modric T, Versic S, Sekulic D (2020). Aerobic fitness and game performance indicators in professional football players; playing position specifics and associations. *Heliyon* 6, 1–6. doi: [10.1016/j.heliyon.2020.e05427](https://doi.org/10.1016/j.heliyon.2020.e05427).
- Primasoni N (2012). Benefits of protein to support sports activities, growth and development of early childhood (thesis). Universitas Negeri Yogyakarta. Available at: <https://staffnew.uny.ac.id/upload/198405212008121001/penelitian/manfaat-protein-untuk-perkembangan.pdf>.
- Probandari AN, Pamungkasari EP, Febrinasari RP, et al (2020). Quantitative research methods: Strategies for writing health research proposals. Surakarta. UNS Press. Available at: <https://opac.perpusnas.go.id/DetailOpac.aspx?id=1350296#>.
- Ryan-Stewart H, Faulkner J, Jobson S (2018). The influence of somatotype on anaerobic performance. *Barbosa TM. PLoS One* 13, 1–11. doi: [10.1371/journal.pone.0197761](https://doi.org/10.1371/journal.pone.0197761).
- Sanders GJ, Turner Z, Boos B, et al (2017). Aerobic capacity is related to repeated sprint ability with sprint distances less than 40 meters. *International Journal of Exercise Science* 10, 197–204. Available at: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=PMC5360375>.
- Scheunemann T (2014). Come on, Indonesia: Curriculum & basic guidelines for Indonesian football for early, youth & senior ages. Gramedia Pustaka Utama. Available at: <https://gpu.id/book/87866/ayo-indonesia>.
- Siri WE (1993). Body composition from fluid spaces and density: Analysis of methods. 1961. *Nutrition* 9, 480–491. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8286893>.
- Sukanta S (2014). Somatotype, body composition and explosive power of athlete and non-athlete. *LASE Journal of Sport Science* 5, 26–34. doi: [10.1515/ljss-2016-0023](https://doi.org/10.1515/ljss-2016-0023).
- Sumardiyono, Probandari AN, Widyaningsih V (2020). Basic statistics for health and medicine; Analysis using SPSS version 23.0. Surakarta. UNS Press. Available at: <https://opac.perpusnas.go.id/DetailOpac.aspx?id=1297802>.
- Wickramarachchi B, Torabi MR, Perera B (2023). Effects of physical activity on physical fitness and functional ability in older adults. *Gerontology and Geriatric Medicine*. doi: [10.1177/23337214231158476](https://doi.org/10.1177/23337214231158476).
- Young S (2020). OCR A-level physical Education student guide 1: Physiological factors affecting performance. Hodder Education. Available at: <https://www.abebooks.com/9781510472082/OCR-A-level-Physical-Education-Student-1510472088/plp>.

