# Correlation of Working Style to Stress, Coping Mechanism and Metabolic Syndrome Risk in Sedentary Workers

# Dominikus Raditya Atmaka<sup>1</sup>, Shintia Yunita Arini<sup>2</sup>, Bian Shabri Putri Irwanto<sup>3</sup>, Asri Meidyah Agustin<sup>4</sup>, Aliffah Nurria Nastiti<sup>5</sup>, Muh. Agus Prianto<sup>6</sup>, Andi Suci Rahman<sup>7</sup>

<sup>1,4,5</sup>Department of Nutrition, Faculty of Public Health, Universitas Airlangga, Indonesia
 <sup>2,3</sup>Department of Occupational Safety and Health, Faculty of Public Health, Universitas Airlangga, Indonesia
 <sup>1,2,3,4,5</sup>Universitas Airlangga, Campus C Mulyorejo, Surabaya, East Java, 60115 Indonesia
 <sup>6,7</sup>HCBP and Culture Department, Head Holding Kalla Group, Makassar, Indonesia

#### ABSTRACT

**Introduction:** Emotional eating, as a method of stress coping mechanism in workers, makes a person tend to consume more energy daily. Uncontrolled daily intake will lead to a high probability of a person developing metabolic syndrome in the future. The aim of this study is to analyze the relationship between working style to stress, coping mechanisms, and metabolic syndrome risk in sedentary workers. **Method:** This research is a cross-sectional study conducted at Kalla Group. Measurement of blood biochemistry (total cholesterol, blood glucose and uric acid) used FORA 6 Plus, blood pressure was measured using Omron digital sphygmomanometer 7130, waist circumference using Seca 201 measuring tape, and height using GEA microtoise. Measurements of body composition used Omron Karada Scan Hbf 375 Body Fat Composition Monitor Bioimpedance Analyzer. Stress was measured using Kessler Psychological Distress Scale K10 instrument. Coping mechanism was measured using Carver's COPE Inventory. **Result:** Significant correlations were found for cholesterol levels (p = 0.044), waist circumference (p = 0.036), blood pressure (p = 0.042), and muscular percentage (p = 0.744), blood glucose (p = 0.128), uric acid (p = 0.380), visceral fat percentage (p = 0.128), and fat mass (p = 0.547). **Conclusion:** Significant correlations were found between working style for cholesterol levels, waist circumference, blood pressure, and muscular percentage, suggesting that different working styles may influence these health aspects.

Keywords: degenerative disease, metabolic syndrome, sedentary workers, stress coping mechanism, working style

#### **Corresponding Author:**

Dominikus Raditya Atmaka E-mail: dominikus.raditya@fkm.unair.ac.id Telephone: +6287780621888

# **INTRODUCTION**

Work style is how an employee combines and expresses the professional, organizational, political, and moral value during daily activities in the workplace (Moslehpour *et al*, 2019). Every person has their own work style and each work style determined job satisfaction and personal stress. For example, a study in Romania showed work style has impact on positive interpersonal relationship, motivation, social support, and more effective and efficient management (Niculiță, 2015). For this reason, it is important to know how work style may impact further life, stress level, and potential health issues in the future.

The adult age group is an age group that is vulnerable to degenerative diseases due to nutritional problems (Gennuso *et al*, 2013; Shuremu, Belachew and Hassen, 2023). This is shown by the increasingly high incidence of obesity which continues to increase every year, as can be seen in the Indonesian Basic Health Research data where, in 2013, the obesity rate was only 14.8%, but increased rapidly in 2018 to reach 21.8% (Ministry of Health Republik Indonesia, 2018). The high potential for overnutrition in adults can trigger various other health problems related to metabolism (Ryan and Heaner, 2014; Christoffersen *et al*, 2023).

Metabolic syndrome is a collection of symptoms that indicate a decrease in the body's ability to carry

©2025 IJOSH All right reserved. Open access under CC BY NC–SA license doi:10.20473/ijosh.v14i1.2025.15-25. Received June 26, 2024; 1<sup>st</sup> revision April 09, 2024; 2<sup>nd</sup> revision August 22, 2024; Accepted December 03, 2024; Published: April 2025. Published by Universitas Airlangga.

*Cite this as:* Atmaka, D. R. et al. (2025) 'Correlation of Working Style to Stress, Coping Mechanism and Metabolic Syndrome Risk in Sedentary Workers', The Indonesian Journal of Occupational Safety and Health, 14(1), pp. 15-25.

out metabolic functions such as cholesterolemia, hypertension, central obesity, and insulin resistance (Grundy, 2016; Hoyas and Leon-Sanz, 2019). Based on data from the Indonesia Family Life Survey (IFLS), the prevalence of metabolic syndrome in Indonesia is 21.6%, which is dominated by low levels of HDL cholesterol (66.41%), hypertension (64.45%), and central obesity (43.21%) (Zahtamal, Prabandari and Setyawati, 2014; Sigit et al, 2020). Metabolic syndrome leads to various degenerative diseases in the future such as stroke, diabetes mellitus, coronary heart disease, and others. The incidence of metabolic syndrome in adults is often caused by wrong food consumption patterns, low physical activity, and stress that is not managed properly (Listyandini et al, 2021; Christoffersen et al, 2023).

Mental emotional disorder is a health problem that often occurs in the adult age group and its prevalence continues to increase due to stress that is not managed properly (Pertiwi, Denny and Widjasena, 2017; Rose et al, 2017). Based on Indonesian Basic Health Research data, it is known that there has been an increase in the number of people with mental emotional disorders from 6% in 2013 to 9.8% in 2018 (Health, 2015; Ministry of Health Republik Indonesia, 2018). Uncontrolled stress is usually caused by personal inability to find appropriate stress coping mechanisms as an effort to regulate stress that occurs within themselves (Graves et al, 2021). High stress levels in a person will produce more of the hormone cortisol which creates hunger and satiety signals (van der Meij et al, 2018; Stammers et al, 2020). This will cause a person to have the desire to eat as an effort to relieve stress from within. This is called the emotional eating phenomenon. The high incidence of emotional eating in adults makes a person tend to consume more energy daily (Tan and Chow, 2014; Serin and Şanlıer, 2018; Rachmawati, Anantanyu and Kusnandar, 2019). Uncontrolled daily intake will lead to a high probability of a person developing metabolic syndrome in the future (Samuel and Cohen, 2018).

Work that does not involve a lot of physical activity, or called sedentary work, is a type of work that involves minimal movement for long periods of time (Park *et al*, 2018). Sedentary workers sit more at work and do a lot of activities in front of the screen. Monotonous work conditions and minimal movement cause workers to experience higher levels of work stress and the potential for emotional eating (Tan and Chow, 2014; Samuel and Cohen, 2018; Serin and Şanlıer, 2018). The high level of emotional eating in the workplace makes workers snack more often and have a poor diet (high in sugar, salt and fat). This makes the incidence of metabolic syndrome and degenerative diseases more common in sedentary workers (Upadhyay et al, 2018). Until now there has been no research related to stress coping mechanisms for dealing with stress in sedentary workers in Eastern Indonesia. Compared to conditions in the western part of Indonesia, the eating habits of people in the eastern part of Indonesia are very different, for example, by the high consumption of high fat food like organs, high fish consumption, and high coconut milk. For this reason, this research was conducted with the aim of looking at the relationship between working style, stress, and coping mechanisms to the metabolic syndrome risk in sedentary workers.

#### METHOD

This research is an observational study with the research design used is cross-sectional. The research was conducted at the Kalla Group in Makassar, South Sulawesi, Indonesia. Kalla Group was chosen because it is one of the largest companies in Eastern Indonesia which controls various sectors ranging from infrastructure, transportation, to expeditions. The research was conducted between June and August 2023. The population selected was Kalla Group workers in the headquarters and branch offices in South Sulawesi Province. The subjects selected were permanent employees of the Kalla Group who had worked for at least six months, were workers with a sedentary activity level, and were willing to take part in a series of research as evidenced by signing an informed consent. A total of 133 workers from total of 1600 workers as population were selected using a purposive sampling technique according to the proportion of workers based on work subsector.

The independent variable measured in this study is working style. Meanwhile the dependent variables measured include stress level, coping mechanism, metabolic syndrome (cholesterol, blood glucose, uric acid, waist circumference, and blood pressure), and body composition (muscular percentage, visceral fat percentage, and fat mass percentage). Working style is measured using Working Style Assessment Questionnaire made by East London NHS Foundation Trust. The questionnaire consisted of five multiple choice questions, with which each option of choices having a different referral to a specific working style. The highest ranks for all five questions are then concluded to identify the working style of each worker. The results are then divided each respondent into four types of working style namely analytical, driver, amiable, and expressive.

Metabolic syndrome measurements are carried out using several parameters, namely total cholesterol levels, blood glucose levels, uric acid levels, waist circumference, and blood pressure. Measurement of blood biochemistry (total cholesterol, blood glucose and uric acid) using the FORA 6 Plus blood multifunctional monitoring system. Blood pressure was measured using an Omron digital sphygmomanometer 7130. Blood pressure measurements were carried out when the subject was at rest, had not been doing work in the last one hour, and had not been doing physical activity in the last one hour.

Waist circumference was measured using a Seca 201 measuring tape. Height was measured using the GEA microtoise. Measurements of body weight, muscular percentage, visceral fat and fat mass were carried out using the Omron Karada Scan Hbf 375 Body Fat Composition Monitor series Bioimpedance Analyzer. Body composition measurements were taken by wearing the lightest clothing possible, not wearing metal jewelry that is conductive, not carrying any items, and not wearing footwear. Subjects were asked to stand on the tool and hold the tool for approximately three seconds.

Stress measurement was carried out using the Kessler Psychological Distress Scale K10 instrument. Coping mechanism measurements were carried out using the COPE Inventory instrument. The K10 questionnaire used consists of 10 questions and the coping mechanism questionnaire consists of 24 questions. The questionnaire is given online using Google Form and is a self-assessment based on the results of performance assessments over the last three months.

Data processing was carried out using STATA 12 software. Normality testing was carried out using the Kolmogorov Smirnov test, followed by hypothesis relationship testing carried out using the Chi Square test for categorical data (nominal and ordinal) with the Pearson method for normally distributed data, and with the Fisher's exact method. for data that are not normally distributed. Linear regression for ratio data was carried out using the One Way ANOVA test. The research was approved by the Health Research Ethics Committee, Faculty of Public Health, Universitas Airlangga with number 96/EA/KEPK/2023.

# RESULTS

Table 1. presents a comprehensive overview of the respondents' characteristics, revealing out of the total respondents, 74 were male, accounting for 55.64% of the sample, and slightly older demographic with 69 respondents were older than 30 years, representing 51.88% of the sample. Stress levels are predominantly severe with 39 respondents (29.32%) falling into the moderate stress category and 70 respondents (52.63%) were classified as having severe stress. A significant portion utilizing problem-focused coping mechanisms were found with 74 respondents (55.64%) who employed problem-focused coping strategies. Analytical working styles are prevalent with 62 respondents (46.62%) were analytical. Metabolic health indicators such as cholesterol, blood glucose, and uric acid levels show a varied distribution, with a notable portion of the population having high measurements namely 64 respondents (48.12%) had high cholesterol levels, 26 respondents (19.55%) had high blood glucose levels, and 39 respondents (29.32%) had high uric acid levels.

Waist circumference and blood pressure data indicate a substantial proportion of respondents at risk where 68 respondents (51.13%) were classified as high risk based on their waist circumference, 57 respondents (42.86%) were in the prehypertension category, 10 respondents (7.52%) had Stage 1 hypertension, and 2 respondents (1.50%) had Stage 2 hypertension. Muscle mass (84 respondents (63.16%) had a low muscular percentage), visceral fat (29 respondents (21.80%) had high visceral fat and 17 respondents (12.78%) had very high visceral fat), and overall fat mass percentages (43 respondents (32.33%) had a high fat mass percentage and 72 respondents (54.14%) had a very high fat mass percentage) highlight areas of concern regarding body composition and potential health risks.

Table 2 illustrates various correlations between working style and health indicators. Significant correlations were found for cholesterol levels (p = 0.044), waist circumference (p = 0.036), blood pressure (p = 0.042), and muscular percentage (p = 0.032), suggesting that different working styles may influence these health aspects. No significant correlations were found between working style for

Characteristics	Frequency (n)	Percentage (%)
Sex		
Male	74	55.64
Female	59	44.36
Age		
≤30 years old	64	48.12
> 30 years old	69	51.88
Stress Category		
Well	16	12.03
Mild disorder	8	6.02
Moderate disorder	39	29.32
Severe disorder	70	52.63
Coping Mechanism		
Emotional focused	59	11 36
coping	57	.50
Problem focused coping	74	55.64
Working Style		
Amiable	15	11.28
Analytical	62	46.62
Driver	35	26.31
Expressive	21	15.79
Cholesterol		
Normal	69	51.88
High	64	48.12
Blood Glucose		
Normal	107	80.45
High	26	19.55
Uric Acid		
Normal	94	70.68
High	39	29.32
Waist Circumference		
Normal	65	48.87
High Risk	68	51.13
Blood Pressure		
Normal	64	48.12
Pre Hypertension	57	42.86
Stage 1 Hypertension	10	7.52
Stage 2 Hypertension	2	1.50
Metabolic Syndrome Status		
Normal	96	72.18
Metabolic Syndrome	37	28.82
Muscular Percentage		
Low	84	63.16
Normal	34	25.56
High	10	7.52
Very High	5	3.76

 Table 1. Respondent Characteristics

Advanced Table 1. Respondent Characteristics

Characteristics	Frequency (n)	Percentage (%)			
Visceral Fat Percentage					
Normal	87	65.42			
High	29	21.80			
Very High	17	12.78			
Fat Mass Percentage					
Normal	18	13.53			
High	43	32.33			
Very High	72	54.14			

stress level (p = 0.683), coping mechanisms (p = 0.744), blood glucose (p = 0.128), uric acid (p = 0.380), visceral fat percentage (p = 0.128), and fat mass (p = 0.547), indicating that working style might not have a strong impact on these factors.

#### DISCUSSION

#### **Respondent Characteristics**

The data provided in Table 1 highlight key aspects of respondents' health, including stress levels, coping mechanisms, and metabolic health indicators, all of which are relevant for understanding health risks in sedentary workers. Research suggests that chronic stress, especially when it reaches severe levels as shown by 52.63% of respondents in data, significantly increases the risk of cardiovascular diseases and other stress-related conditions (Proia *et al*, 2016; Anestal, 2022; Hoveling *et al*, 2022; Shchaslyvyi, Antonenko and Telegeev, 2024).

Problem-focused coping mechanisms, which were predominant in 55.64% of respondents, are known to be effective in mitigating stress's negative health impacts, as they focus on addressing the cause of stress directly (Anestal, 2022; Theodoratou and Argyrides, 2024). Moreover, the high percentage of respondents with metabolic health issues, including elevated cholesterol (48.12%), blood glucose (19.55%), and uric acid (29.32%), signals a significant risk for cardiovascular diseases, diabetes, and other metabolic conditions. Elevated waist circumference and high blood pressure among respondents further support these health risks, indicating potential for increased cardiovascular disease prevalence(Salk, Hyde and Abramson, 2017; Bonnet et al, 2020; Aly et al, 2022).

The implications of these findings emphasize the need for targeted health interventions, particularly

	Working Style									
-	Am	iable	Analytical		Di	Driver		Expressive		p*
	n	%	n	%	n	%	n	%		
Stress Level										
Mild Disorder	1	0.75	7	5.26	5	3.75	3	2.65		
Moderate Disorder	1	0.75	6	4.50	1	0.75	0	0.00	6.558	0.683
Severe Disorder	3	2.65	18	13.53	13	9.77	5	3.75		
Well	10	7.50	21	15.79	16	12.03	13	9.77		
		C	oping M	echanism						
Emotional Focused	7	5.26	29	21.80	16	12.03	7	5.26	8.084	0.044
Problem Focused	8	6.00	33	24.81	19	14.29	14	10.52		
			Chole	sterol						
High	7	5.26	27	20.30	14	10.52	16	12.03	9.917	0.128
Normal	8	6.00	35	26.32	21	15.79	5	3.75		
			Blood (	Glucose						
High	6	4.50	18	13.53	7	5.26	8	6.00	3.075	0.380
Normal	9	6.77	44	33.08	28	21.05	13	9.77		
		W	aist Circ	umference						
High Risk	13	9.77	29	21.80	16	12.03	10	7.50	8.566	0.036
Normal	2	1.50	33	39.85	19	14.29	11	8.27		
			Blood P	ressure						
Prehypertension	4	3.00	24	18.00	17	12.78	12	9.00		
Stage 2 Hypertension	0	0.00	3	2.65	6	4.50	4	3.00	13.441	0.042
Stage 1 Hypertension	1	0.75	0	0.00	1	0.75	24	18.00		
Normal	10	7.50	35	26.32	11	8.27	17	12.78		
		Μ	uscular 1	Percentage						
Low	11	8.27	30	26.50	28	21.05	15	13.25		
High	3	2.65	6	4.50	1	0.75	0	0.00	18.288	0.032
Very High	0	0.00	3	2.65	2	1.50	0	0.00		
Normal	1	0.75	22	16.54	5	3.75	6	0.75		
		Vis	ceral Fat	Percentage	•					
High	3	2.65	13	9.77	10	7.50	3	2.65	9.911	0.128
Very High	5	3.75	3	2.65	5	3.75	5	3.75		
Normal	8	6.00	45	33.83	20	15.00	13	9.77		
			Fat I	Mass						
High	3	2.65	21	15.79	11	8.27	8	6.00	4.974	0.547
Very High	11	8.27	29	21.80	21	15.79	10	7.50		
Normal	1	0.75	11	8.27	4	3.00	3	2.65		

 Table 2. Correlation of Working Style to Stress Level, Coping Mechanism, Metabolic Syndrome Indicator, and Body Composition

those addressing stress management, metabolic health monitoring, and lifestyle changes. Behavioral stress reduction programs (BSRPs), which have shown promise in reducing stress and improving health outcomes, could be particularly useful for populations experiencing both psychological and metabolic stress (Kivimäki, Bartolomucci and

Kawachi, 2023; Hill, Mostafa and Obeng-Gyasi, 2024). By integrating stress management with health monitoring, interventions could mitigate the elevated risk factors seen in this population. Future health policies and programs may benefit from incorporating problem-focused coping strategies and stress-reduction techniques as part of a comprehensive approach to managing stress and metabolic health (James *et al*, 2023; Ebrahim *et al*, 2024).

#### **Correlation Test**

The relationship between working style and various health indicators, such as stress level, coping mechanisms, metabolic syndrome indicators, and body composition, is complex and multifaceted. As stated by Moslehpour (2019), working style is how an employee combines and expressed the professional, organizational, political, and moral value during daily activities in workplace. Every person has their own work style and each work style determines job satisfaction and personal stress. Working style significantly impacts an individual's stress level. High job demands, low control, and imbalance between effort and reward are wellestablished predictors of work-related stress. The job demand-control (JDC) model posits that high job demands and low job control lead to increased stress levels (International Labour Organization, 2016). Similarly, the effort-reward imbalance (ERI) model suggests that a disproportionate balance between the efforts put into work and the rewards received exacerbates stress (Goetzel et al, 2014; Balogh et al, 2019; Kinman, 2019).

# **Stress Level**

There was no significant correlation found between working style and stress level (p = 0.683). However, the data suggest that individuals with an analytical working style experienced higher levels of severe stress (13.53%) compared to other working styles. In contrast, amiable individuals showed the highest percentage of being well (7.50%). The lack of significant correlation suggests that while certain working styles might exhibit higher stress levels, other factors likely contribute to stress management and perception. This aligns with existing literature indicating that individual stress responses are multifactorial, influenced by personality, work environment, and coping mechanisms (Arnett, Žukauskiene and Sugimura, 2014; Nordander et al, 2016; Anestal, 2022; Khairy, 2022).

# **Coping Mechanism**

Coping mechanisms are strategies employed by individuals to manage stress. Working style influences the adoption of specific coping mechanisms. For example, individuals in high-stress jobs may rely more on problem-focused coping, such as time management and seeking support, while those in lower-stress environments might engage in emotion-focused coping, such as relaxation and avoidance (Anestal, 2022; Khairy, 2022; Aras, Tandiayuk and Fatihah, 2023).

The study found no significant correlation between working style and coping mechanisms (p = 0.744). Problem-focused coping was more prevalent across all working styles, with analytical individuals showing the highest percentage (24.81%). This suggests that regardless of working style, individuals may prefer problem-focused coping strategies, which involve tackling the problem directly to alleviate stress. This finding is consistent with the coping theory which posits that problem-focused coping is often more effective in controllable stress situations (Graves *et al*, 2021; Lestari, 2021; Khairy, 2022).

#### **Metabolic Syndrome**

Metabolic syndrome is a cluster of conditions that increase the risk of heart disease, stroke, and diabetes. Indicators include abdominal obesity, high blood pressure, elevated fasting glucose, high triglycerides, and low HDL cholesterol levels. Sedentary working styles, characterized by prolonged sitting and lack of physical activity, are strongly associated with metabolic syndrome. Studies have shown that sedentary behavior increases the risk of developing metabolic syndrome by up to 50% (Grundy, 2016; Hoyas and Leon-Sanz, 2019; Listyandini *et al*, 2021).

There was a significant correlation between working style and cholesterol levels (p = 0.044). Analytical individuals had the highest percentage of high cholesterol (20.30%). A significant correlation was found at waist circumference (p = 0.036), with analytical individuals again showing the highest risk (21.80%). Significant correlation was observed at blood pressure (p = 0.042). Analytical individuals had a higher incidence of prehypertension (18.00%). The significant correlation between working style and certain metabolic syndrome indicators suggests that individuals with an analytical working style may be at higher risk for cardiovascular and metabolic diseases. This could be due to the nature of analytical work, which may involve prolonged periods of sedentary behavior and higher stress levels, known risk factors for these conditions (Rahma Listyandini et al, 2021; Khairy, 2022; Aras, Tandiayuk and Fatihah, 2023).

# **Body Composition**

Body composition, which includes the proportion of fat and non-fat mass in the body, is influenced by working style. Physically demanding jobs often result in lower body fat percentages and higher muscle mass, whereas sedentary jobs can lead to increased body fat and decreased muscle mass. Research indicates a significant correlation between sedentary working styles and higher body mass index (BMI) (Barnes *et al*, 2015; Park, 2017; Srikanthan *et al*, 2021).

Significant correlation on muscular percentage was found (p = 0.032), with analytical individuals showing higher percentages of low muscular percentage (26.50%). Meanwhile no significant correlation was found for these metrics for other metrics (visceral fat, fat mass). The significant correlation between working style and muscular percentage highlights the potential impact of work environment and style on physical health. Analytical work, often associated with sedentary behavior, may lead to lower muscle mass, increasing the risk for various health issues (Santos *et al*, 2014; Tseng *et al*, 2014; Park, 2017; Wang and Torriani, 2020).

#### Recommendation

Integrating these findings, it is clear that working style is a crucial determinant of overall health. High-stress working styles exacerbate stress levels and may lead to maladaptive coping mechanisms, while sedentary working styles contribute to adverse metabolic outcomes and poor body composition. Consequently, interventions aimed at modifying working styles, such as increasing job control, ensuring adequate rewards, promoting physical activity, and implementing stress management programs, can mitigate these negative health effects. It is important to assess each workers early to their working style from their recruitment process in order to place workers in the right position to prevent them from unsuitable job desk and interest and may decrease the work stress and increase the job satisfaction, for example, by using several assessment tools like Myers-Briggs Type Indicator (MBTI), DISC assessment, or the Big Five Personality Traits. Each assessment can identify personal preferences in working style like introversion or extraversion which then significantly affects their cooperation in work. This reason may enhance self-awareness among employees and improve team collaboration by highlighting diverse

strength and work preferences. This may allow the head of work team to tailor the leadership style and communication strategies to personalize among different working styles.

Several technical recommendations act as efforts which need to be done as well to prevent workers from future metabolic syndrome development by reduction of stress possibilities happening during work. The leaders need to be able to customize several activities to reduce stress based on diverse working style environment, like conducting team building activities, implementing cross-functional training, creating diversity-inclusive policies, utilizing diverse communication and collaboration working tools, giving one-on-one regular feedback, using open communication channels through different communication styles, and aligning team goals based on individual strengths. A supportive and collaborative work environment enhances social support, which is critical for stress reduction and improved mental health. Research shows that workers with strong social support tend to have lower stress levels and better coping mechanisms. Encouraging teamwork and open communication enables workers to problem-solve together, fostering an analytical working style that can alleviate stress. In hope these effort can accommodate the needs of the team to work in better environment following the diverse working styles (Bush et al, 2014; Goetzel et al, 2014; Samudera, Efendi and Indarwati, 2021).

Regular movement breaks, such as 5-minute standing or stretching breaks every hour, reduce physical strain and prevent mental fatigue. Frequent breaks to walk or stretch have been linked to improvements in cholesterol, blood sugar, and waist circumference. Consider using standing desks or walking meetings to integrate more movement into the workday, as this can reduce the harmful effects of prolonged sitting. Offering employees access to mindfulness training, yoga, and cognitive-behavioral therapy workshops can help improve emotional regulation and promote adaptive coping strategies, reducing reliance on maladaptive mechanisms like avoidance. Such programs help individuals manage stress by reframing negative thoughts and promoting relaxation techniques that can reduce mental and physical stress indicators (Goetzel et al, 2014; Hosker, Elkins and Potter, 2019; Shchaslyvyi, Antonenko and Telegeev, 2024; Theodoratou and Argyrides, 2024).

Periodic health assessments that monitor cholesterol, blood glucose, and body composition

can help workers stay aware of their health risks. Providing incentives like gym memberships or health bonuses for maintaining good metabolic health can motivate workers to engage in healthier lifestyles. Providing access to healthy meals in the workplace or encouraging participation in group fitness activities can reduce the risks associated with metabolic syndrome, including high cholesterol and blood glucose levels. These measures also have positive effects on mental well-being, creating a more productive work environment(Zahtamal, Prabandari and Setyawati, 2014; Hoyas and Leon-Sanz, 2019; Rahma Listyandini *et al*, 2021; Anestal, 2022; Kivimäki, Bartolomucci and Kawachi, 2023)

# CONCLUSION

Significant correlations were found between working style for cholesterol levels, waist circumference, blood pressure, and muscular percentage, suggesting that different working styles may influence these health aspects. No significant correlations were found between working style for stress level, coping mechanisms, blood glucose, uric acid, visceral fat percentage, and fat mass, indicating that working style might not have a strong impact on these factors. This discussion highlights the interconnectedness of working style with stress levels, coping mechanisms, metabolic syndrome indicators, and body composition. Understanding these relationships is essential for developing effective workplace interventions to promote employee health and well-being.

#### **CONFLICT OF INTEREST**

The author declare there is no conflict of interest during the process of the research.

#### AUTHOR CONTRIBUTION

DRA and SYA: designing the study method, funding acquisition, analyzing the data, writing the manuscript. BSPI: managing resources, data collection, data curation, editing manuscript. AMA and ANN: managing resources, data collection, and data management. MAP and ASR: managing the research process, managing the respondents, managing the resources, communicating with stakeholders.

#### ACKNOWLEDGMENTS

We thank Kalla Group for the coordination of data collection during the study, and to the Faculty of Public Health for the funding through the Penelitian Dosen Pemula scheme.

#### REFERENCES

- Aly, N.M. et al (2022) 'Association between COVID-19 Stress, Coping Mechanisms and Stress-related Oral Conditions among Egyptian Adults: A Cross-Sectional Study', *Scientific Reports*, 12(1). https:// doi.org/10.1038/s41598-022-22961-z
- Anestal, C. (2022) Stress, Coping, and Disease Awareness with Metabolic Disease Risk: A Longitudinal Cohort Study. Available at: http:// library.ucf.edu
- Aras, R.A., Tandiayuk, S. and Fatihah, R.N. (2023)
  'The Relationship Between Stress Coping Strategies and Work-Life Balance Among Bank Employees', Jurnal Office: Jurnal Pemikiran Ilmiah dan Pendidikan Administrasi Perkantoran, 9(1), pp. 101–108. Available at: http://ojs.unm. ac.id/jo.
- Arnett, J.J., Žukauskiene, R. and Sugimura, K. (2014) 'The New Life Stage of Emerging Adulthood at Ages 18-29 Years: Implications for Mental Health', *The Lancet Psychiatry*. Elsevier Ltd, pp. 569–576. https://doi.org/10.1016/S2215-0366(14)00080-7
- Balogh, I. et al (2019) 'Work-related neck and upper limb disorders - Quantitative exposureresponse relationships adjusted for personal characteristics and psychosocial conditions', BMC Musculoskeletal Disorders, 20(1). https:// doi.org/10.1186/s12891-019-2491-6
- Barnes, T.L. et al (2015) 'Snacking Behaviors, Diet Quality, and Body Mass Index in a Community Sample of Working Adults', Journal of the Academy of Nutrition and Dietetics, 115(7), pp. 1117–1123. https://doi.org/10.1016/j. jand.2015.01.009
- Bonnet, J.P. et al (2020) 'Breakfast Skipping, Body Composition, and Cardiometabolic Risk: A Systematic Review and Meta-Analysis of Randomized Trials', *Obesity*, 28(6), pp. 1098– 1109. https://doi.org/10.1002/oby.22791
- Bush, D.E. et al (2014) 'Evaluation of a pilot promotora program for Latino forest workers in Southern Oregon', American Journal of

*Industrial Medicine*, 57(7), pp. 788–799. https:// doi.org/10.1002/ajim.22347

- Christoffersen, T. et al (2023) 'Development of an Approach for Identifying Overnutrition among Older Adults in Community Health Care Settings: - an Opinion Paper', Clinical Nutrition Open Science, 47, pp. 102–120. https://doi. org/10.1016/j.nutos.2022.12.008
- Ebrahim, O.S. *et al* (2024) 'Perceived Stress and Anxiety among Medical Students at Helwan University: A Cross-Sectional Study', *Journal of Public Health Research*, 13(1). https://doi. org/10.1177/22799036241227891.
- Gennuso, K. et al (2013) 'Sedentary Behavior, Physical Activity, and Markers of Health in Older Adults', Medicine and Science in Sports Exercise, 45, pp. 1493–500. https://doi.org/10.1249/ MSS.0b013e318288a1e5.Sedentary.
- Goetzel, R.Z. et al (2014) 'Do Workplace Health Promotion (Wellness) Programs Work?', Journal of Occupational and Environmental Medicine, 56(9), pp. 927–934. https://doi.org/10.1097/ JOM.000000000000276.
- Graves, B.S. *et al* (2021) 'Gender Differences in Perceived Stress and Coping among College Students', *PLoS ONE*, 16(8), pp. 1-12. https:// doi.org/10.1371/journal.pone.0255634.
- Grundy, S.M. (2016) 'Metabolic Syndrome Update', *Trends in Cardiovascular Medicine*, 26(4), pp. 364–373. https://doi.org/10.1016/j. tcm.2015.10.004.
- Health, I.M. of (2015) 'Infodation : Situasi Kesehatan Kerja', Pusat Data dan Informasi Kementerian Kesehatan RI [Preprint]. Available at: https://www. kemkes.go.id/download.php?file=download/ pusdatin/infodatin/infodatin-kesja.pdf.
- Hill, M., Mostafa, S. and Obeng-Gyasi, E. (2024) 'Chronic Stress Indicator: A Novel Tool for Comprehensive Stress Analysis', International Journal of Environmental Research and Public Health, 21(3), pp. 1-27. https://doi.org/10.3390/ ijerph21030302.
- Hosker, D.K., Elkins, R.M. and Potter, M.P. (2019) 'Promoting Mental Health and Wellness in Youth Through Physical Activity, Nutrition, and Sleep', *Child and Adolescent Psychiatric Clinics of North America*, 28(2), pp. 171–193. https://doi. org/10.1016/j.chc.2018.11.010.
- Hoveling, L.A. *et al* (2022) 'Socioeconomic Differences in Metabolic Syndrome Development: Examining the Mediating Role of Chronic Stress using the Lifelines Cohort Study', *BMC Public*

*Health*, 22(1), pp. 1-11. https://doi.org/10.1186/ s12889-022-12684-1.

- Hoyas, I. and Leon-Sanz, M. (2019) 'Nutritional Challenges in Metabolic Syndrome', *Journal* of Clinical Medicine, 8(9), p. 1301. https://doi. org/10.3390/jcm8091301.
- International Labour Organization (2016) Workplace Stress: A Collective Challenge, International Labour Organization.
- James, K.A. et al (2023) 'Understanding the Relationships between Physiological and Psychosocial Stress, Cortisol and Cognition', Frontiers in Endocrinology, 14, pp. 1-20. https:// doi.org/10.3389/fendo.2023.1085950
- Khairy, H.A. (2022) 'Coping Strategies' Effect on Occupational Stress and its Reflection on Employees' Turnover Intention in Hotels: The Moderating Role of Social Exchange', *Journal* of the Faculty of Tourism and Hotels-University of Sadat City, 6(2), pp. 90-110. https://doi. org/10.21608/mfth.2022.274406.
- Kinman, G. (2019) 'Effort-reward Imbalance in Academic Employees: Examining Different Reward Systems', *International Journal of Stress Management*, 26(2), pp. 184–192. https://doi. org/10.1037/str0000128.
- Kivimäki, M., Bartolomucci, A. and Kawachi, I. (2023) 'The Multiple Roles of Life Stress in Metabolic Disorders', *Nature Reviews Endocrinology*, 19(1), pp. 10–27. https://doi. org/10.1038/s41574-022-00746-8
- Lestari, R.F. (2021) 'Strategi Coping pada Mahasiswa Ditinjau dari Big Five Personality', *Acta Psychologia*, 3(1), pp. 29–37. Available at: http:// journal.uny.ac.id/index.php/acta-psychologia.
- van der Meij, L. *et al* (2018) 'Hair Cortisol and Work Stress: Importance of Workload and Stress Model (JDCS or ERI)', *Psychoneuroendocrinology*, 89(July), pp. 78–85. https://doi.org/10.1016/j. psyneuen.2017.12.020
- Ministry of Health Republik Indonesia (2018) 'Riset kesehatan dasar (Riskesdas) 2018', Jakarta: Badan Penelitian dan Pengembangan Kesehatan Kementrian Kesehatan Republik Indonesia [Preprint].
- Moslehpour, M. *et al* (2019) 'Organizational Climate and Work Style: The Missing Links for Sustainability of Leadership and Satisfied Employees', *Sustainability (Switzerland)*, 11(1), pp. 1-17. https://doi.org/10.3390/su11010125.
- Niculiță, Z. (2015) 'The Relationship Between Work Style and Organizational Climate for Romanian

Employees', *Procedia Economics and Finance*, 32, pp. 1042–1049. https://doi.org/10.1016/ s2212-5671(15)01566-x

- Nordander, C. *et al* (2016) 'Exposure-response Relationships for Work-related Neck and Shoulder Musculoskeletal Disorders - Analyses of Pooled Uniform Data Sets', *Applied Ergonomics*, 55, pp. 70–84. https://doi.org/10.1016/j. apergo.2016.01.010.
- Park, J.H. et al (2018) 'Association between Sedentary Time and Cardiovascular Risk Factors in Korean Adults', Korean Journal of Family Medicine, 39(1), pp. 29–36. https://doi. org/10.4082/kjfm.2018.39.1.29.
- Park, S. (2017) 'Comparison of Body Composition between Fashion Models and Women in General', *Journal of Exercise Nutrition & Biochemistry*, 21(4), pp. 22–26. https://doi.org/10.20463/ jenb.2017.22.
- Pertiwi, E., Denny, H. and Widjasena, B. (2017) 'Hubungan Antara Beban Kerja Mental Dengan Stres Kerja Dosen Di Suatu Fakultas', Jurnal Kesehatan Masyarakat (e-Journal), 5(3), pp. 260-268.
- Proia, P. et al (2016) 'Lactate as a metabolite and a regulator in the central nervous system', *International Journal of Molecular Sciences*, 17(9), pp.1-20. https://doi.org/10.3390/ ijms17091450.
- Rachmawati, Y., Anantanyu, S. and Kusnandar (2019) 'Emotional Eating, Snacking Behavior and Nutritional Status among Adolescents', *International Journal of Public Health Science*, 8(4), pp. 413–418. https://doi.org/10.11591/ijphs. v8i4.20398.
- Listyandini, R. et al (2021) 'The Dominant Factor of Metabolic Syndrome among Office Workers', Journal of Health Science and Prevention, 5(1), pp. 40–48. https://doi.org/10.29080/jhsp. v5i1.421.
- Rose, D.M. et al (2017) 'Associations of Fatigue to Work-related Stress, Mental and Physical Health in an Employed Community Sample', BMC Psychiatry, 17(1), pp. 1–8. https://doi. org/10.1186/s12888-017-1237-y.
- Ryan, D. and Heaner, M. (2014) 'Guidelines (2013) for Managing Overweight and Obesity in Adults', *Obesity*, 22(S2), pp. i–xvi. Available at: https:// doi.org/10.1002/oby.20818.
- Salk, R.H., Hyde, J.S. and Abramson, L.Y. (2017) 'Gender Differences in Depression in Representative National Samples: Meta-analyses

of Diagnoses and Symptoms', *Psychological Bulletin*, 143(8), pp. 783–822. https://doi. org/10.1037/bul0000102.

- Samudera, W.S., Efendi, F. and Indarwati, R. (2021) 'Effect of Community and Peer Support based Healthy Lifestyle Program (CP-HELP) on self-care behavior and fasting blood glucose in patient with type 2 Diabetes Mellitus', *Journal* of Diabetes and Metabolic Disorders, 20(1), pp. 193–199. https://doi.org/10.1007/s40200-021-00729-y.
- Samuel, L. and Cohen, M. (2018) 'Expressive Suppression and Emotional Eating in Older and Younger Adults: An Exploratory Study', *Archives* of Gerontology and Geriatrics, 78, pp. 127–131. https://doi.org/10.1016/j.archger.2018.06.012
- Santos, D.A. *et al* (2014) 'Reference Values for Body Composition and Anthropometric Measurements in Athletes', *PLoS ONE*, 9(5). https://doi. org/10.1371/journal.pone.0097846.
- Serin, Y. and Şanlıer, N. (2018) 'Emotional Eating, the Factors which Affect Food Intake and Basic Approaches of Nursing Care', *Journal of Psychiatric Nursing*, 9(2), pp.135-146. https:// doi.org/10.14744/phd.2018.23600.
- Shchaslyvyi, A.Y., Antonenko, S. V. and Telegeev, G.D. (2024) 'Comprehensive Review of Chronic Stress Pathways and the Efficacy of Behavioral Stress Reduction Programs (BSRPs) in Managing Diseases', *International Journal of Environmental Research and Public Health. Multidisciplinary Digital Publishing Institute (MDPI)*, 21(8), pp. 1-55. https://doi.org/10.3390/ijerph21081077.
- Shuremu, M., Belachew, T. and Hassen, K. (2023) 'Nutritional Status and its Associated Factors among Elderly People in Ilu Aba Bor Zone, Southwest Ethiopia: A Community-based Crosssectional Study', *BMJ Open*, 13(1). https://doi. org/10.1136/bmjopen-2022-067787.
- Sigit, F.S. *et al* (2020) 'The Prevalence of Metabolic Syndrome and its Association with Body Fat Distribution in Middle-aged Individuals from Indonesia and the Netherlands: A Cross-sectional Analysis of Two Population-based Studies', *Diabetology and Metabolic Syndrome*, 12(1). https://doi.org/10.1186/s13098-019-0503-1.
- Srikanthan, P. et al (2021) 'Sex Differences in the Association of Body Composition and Cardiovascular Mortality', Journal of the American Heart Association, 10(5). https://doi. org/10.1161/JAHA.120.017511.

- Stammers, L. et al (2020) 'Identifying stress-related eating in behavioural research: A review', *Hormones and Behavior*, 124, pp. 1-13. https:// doi.org/10.1016/j.yhbeh.2020.104752.
- Tan, C.C. and Chow, C.M. (2014) 'Stress and Emotional Eating: The Mediating Role of Eating Dysregulation', *Personality and Individual Differences*, 66, pp. 1–4. https://doi.org/10.1016/j. paid.2014.02.033.
- Theodoratou, M. and Argyrides, M. (2024) 'Neuropsychological Insights into Coping Strategies: Integrating Theory and Practice in Clinical and Therapeutic Contexts', *Psychiatry International. Multidisciplinary Digital Publishing Institute (MDPI)*, pp. 53–73. https:// doi.org/10.3390/psychiatryint5010005.
- Tseng, L.A. *et al* (2014) 'Body Composition Explains Sex differential in Physical Performance among

Older Adults', *Journals of Gerontology - Series A Biological Sciences and Medical Sciences*, 69(1), pp. 93–100. https://doi.org/10.1093/ gerona/glt027.

- Upadhyay, J. *et al* (2018) 'Obesity as a Disease', *Medical Clinics of North America*, 102(1), pp. 13– 33. https://doi.org/10.1016/j.mcna.2017.08.004.
- Wang, B. and Torriani, M. (2020) 'Artificial Intelligence in the Evaluation of Body Composition', *Seminars in Musculoskeletal Radiology*, 24(1), pp. 30–37. https://doi.org/10.1055/s-0039-3400267.
- Zahtamal, Z., Prabandari, Y.S. and Setyawati, L. (2014) 'Prevalensi sindrom metabolik pada pekerja perusahaan', Kesmas: Jurnal Kesehatan Masyarakat Nasional (National Public Health Journal), 9(2), pp. 113–120.