

# Indonesian Journal of Tropical and Infectious Disease

Vol. 8 No. 2 May–August 2020

## Research Report

### Identification of SCC MEC Methicillin-Resistant Staphylococcus Aureus (MRSA) From Hospitals' Clinical Samples in Jambi using Polymerase Chain Reaction (PCR)

Humaryanto<sup>1\*</sup>, Hanina<sup>1</sup>, Lipinwati<sup>1</sup>, Charles Apul Simanjuntak<sup>1</sup>

<sup>1</sup>Faculty of Medicine and Health Science, University of Jambi, Jambi Indonesia

Received: 8<sup>th</sup> April 2019; Revised: 29<sup>th</sup> January 2020; Accepted: 23<sup>rd</sup> April 2020

#### ABSTRACT

*Staphylococcal cassette chromosome mec (SCCmec) is one of the mobile genetic elements of Methicillin-Resistant Staphylococcus aureus (MRSA) that carries many resistance genes and allows SCCmec to move from one bacterium to another. Twelve types of SCCmec have been identified throughout the world. Identification of SCCmec type is needed to determine the pattern of MRSA resistance in a particular region. This study aimed to identify the type of SCCmec MRSA from clinical samples. Specifically, this study was conducted at the Biomolecular Laboratory of the Faculty of Medicine and Health Sciences of Jambi University in June 2018-February 2019. Culture was carried out on 100 clinical specimens of festering wound swabs from inpatients at hospitals in Jambi City. A total of 32 samples of Staphytest plus test positive were tested using Cefoxitin disc diffusion method and MecA Polymerase Chain Reaction (PCR). There were 14 samples identified as MRSA isolates, namely twelve samples (85.72%) of SCCmec type III, one sample (7.14%) of SCCmec type II, and one sample (7.14%) of SCCmec type IVb. The results were different from previous studies where all MRSA isolates (100%) in Indonesia were SCCmec type III, although most SCCmec types were still dominated by SCCmec type III. This study concludes that there has been a shift in the content of SCCmec in MRSA isolate originating from hospitals in Jambi city.*

**Keywords:** MRSA, MecA, SCCMec, genetic, resistance

#### ABSTRAK

*Staphylococcal cassette chromosome mec (SCCmec) merupakan salah satu elemen genetik yang mobile pada Methicillin Resistant Staphylococcus aureus (MRSA) yang membawa beberapa gen resistensi dan memungkinkan SCCmec berpindah dari satu bakteri ke bakteri lainnya. Terdapat dua belas tipe SCCmec yang telah teridentifikasi kasi di seluruh dunia. Identifi kasi tipe SCCmec sangat diperlukan untuk mengetahui pola resistensi MRSA di suatu wilayah tertentu. Penelitian ini bertujuan untuk mengidentifikasi kasi tipe SCCmec MRSA dari sampel klinik. Penelitian ini dilakukan di Laboratorium Biomolekuler Fakultas Kedokteran dan Ilmu Kesehatan Universitas Jambi pada bulan Juni 2018-Februari 2019. Kultur dilakukan terhadap 100 spesimen klinik berupa swab luka yang bernanah pada pasien yang dirawat inap di Rumah Sakit di Kota Jambi. Sebanyak 32 sampel yang positif pada Uji Staphytest plus diuji dengan Cefoxitin Disk Difusion Metode dan Polymerase Chain Reaction (PCR) MecA. Terdapat 14 sampel yang teridentifikasi kasi sebagai isolat MRSA. Sebanyak 12 sampel (85,72%) merupakan SCCmec tipe III, satu sampel (7,14%) SCCmec tipe II dan satu sampel (7,14%) SCCmec tipe IVb. Hasil penelitian ini berbeda dengan penelitian sebelumnya dimana seluruh (100%) isolat MRSA di Indonesia merupakan SCCmec tipe III, meskipun tipe SCCmec terbanyak masih didominasi oleh SCCmec tipe III. Kesimpulan dari penelitian ini adalah mulai ditemukannya perubahan kandungan SCCmec pada isolat MRSA yang berasal dari rumah sakit di Kota Jambi.*

**Kata kunci:** MRSA, MecA, SCCMec, genetic, resistensi

---

\* Corresponding Author:

humaryanto\_fkik@unja.ac.id

**How to Cite:** Humaryanto., Hanina., Lipinwati., Chaeles Apul Simanjuntak. Identification of SCC MEC Methicillin-Resistant *Staphylococcus Aureus* (MRSA) From Hospitals' Clinical Samples in Jambi Using Polymerase Chain Reaction (PCR). *Indonesian Journal of Tropical and Infectious Disease*, 8(2), 1–8

## INTRODUCTION

*S. aureus* is a common bacterial pathogen that causes minor to serious disease in human. *S. aureus* can be treated with methicillin (MSSA) and resistant to methicillin (MRSA). Infection of MRSA becomes an important concern throughout the world and associated with infection in both Hospital-acquired Methicillin-Resistant *Staphylococcus aureus* (HA-MRSA) and Community-acquired Methicillin-Resistant *Staphylococcus aureus* (CA-MRSA).<sup>1,2,3</sup> Infection caused by MRSA keeps increasing year to year. According to research in Indonesia, the prevalence of MRSA is approximately 30–40%. The prevalence of MRSA in Cipto Mangunkusumo Hospital on 2010 and Abdul Moeloek Hospital Lampung on 2013 were 32% and 38%, respectively.<sup>4,5</sup>

The resistance of MRSA against beta-lactam antibiotic is encoded by the *mecA* gene. *MecA* gene is a part of the conserved MRSA genetic elements of the Staphylococcal cassette chromosome *mec* (SCC*mec*), encoding PBP2a or PBP2 mutants.<sup>6,7</sup> *MecA* gene is located in a genetic element called the Staphylococcal Cassette Chromosome (SCC*mec*). SCC*mec* is integrated into the chromosome of *S. aureus* at a unique site located near the *S. aureus* origin of replication. SCC*mec* is a mobile genetic element that carries many resistance genes and allows SCC*mec* to move from one bacterium to another.<sup>8</sup> Thirteen types of SCC*mec* have been identified throughout the world.<sup>9</sup>

The components of SCC*mec* are recombinase genes (*ccr* complexes), *mec* complex genes, additional resistant genes, and insertion sequences (IS).<sup>8,10</sup> Differences between SCC*mec* are determined by variations in the *ccr* complex and the *mec* complex. SCC*mec* type I about 39 kb, in the 1960s era, has a composition of type 1 *ccr* complex and class B *mec* complex. SCC*mec* type II about 52 kb, dominant in the 1980s era, has a

composition of type 2 *ccr* complex and the class A *mec* complex. SCC*mec* type III about 67 kb, dominant in the 1980s, has the composition of the type 3 *ccr* complex and the class A *mec* complex. SCC*mec* type IV (a and b) about 20.9–24.3 kb, found in 2002, has a composition of type 2 *ccr* complex and class B *mec* complex.<sup>4,5,6</sup>

Various findings of MRSA patterns in the last decade have shown the changes in distribution, sensitivity to various antibiotics, and possible changes in the SCC*mec* type.<sup>11,12</sup> Identification of SCC*mec* type is needed to determine the pattern of MRSA resistance in a particular region. Based on the previous description, it is important to identify the type of SCC*mec* MRSA from clinical samples.

## MATERIALS AND METHODS

This study was a cross-sectional study. This study was conducted in the Biomolecular Laboratory of the Faculty of Medicine and Health Sciences in Jambi University from June 2018 to February 2019.

A hundred samples of swabs from festering wound were collected from three secondary referral hospitals in Jambi (Raden Mattaher hospital, dr. Bratanata hospital, and Kambang hospital). The swabs were incubated at 30 °C on Mannitol Salt Agar (MSA) for 18–24 hours, the yellowish colony would be confirmed by Gram staining. Gram-positive coccus bacteria were tested using Staphytest plus Test DR 850 M (Oxoid) to detect clumping factor, protein A and type 5 and 8 capsules of polysaccharide.

Positive samples were tested for resistance to cefoxitin antibiotics by using the disc diffusion method in Mueller Hinton (MH) Agar. The susceptibility testing was conducted as a standard of CLSI 2011.<sup>13</sup> Identification of *MecA* gene and the type of SCC*mec* were using Polymerase Chain Reaction (PCR). Primers used are shown in Table 1.

### Preparation of Bacterial DNA Samples, PCR Mec A and PCR SCCmec

DNA samples 5 µl of bacterial suspension (0.5 Mc Farland) from yellowish colonies were incubated at 30°C 18-24 hours on MSA. PCR was performed in a final volume of 25 µl consisting of 5 µl of DNA samples, 10 µl of 2x GoTaq green master mix (Promega), 2 µl 1mM forward primer (Mec A1), 2 µl 1mM reverse primer (Mec A2) and 6 µl of nuclease-free water. Positive control and negative control were *S. aureus* ATCC 43300 and *S. aureus* ATCC 25923. The mixture was denatured at 94°C for 5 minutes followed by 30 cycles, 94°C for 45 seconds, 72°C for 90 seconds, and 72°C for 10 minutes. DNA was amplified with a thermocycler (Thermo scientific, USA).

Multiplex PCR SCCmec was carried out on positive samples of MecA gene to detect SCCmec chromosomes. Primers used are shown in Table 1. PCR was performed in a final volume of 25 µl consisting of 5 µl of DNA samples, 12.5 µl of 2x GoTaq green master mix (Promega), 0.5 µl 1 mM of forward primer, 0.5 µl 1 mM of reverse primer (SCC mec primers type I, II, III, IVa, and IVb) and 2.5 µl nuclease-free water. PCR to identify the type of SCCmec began with an initial denaturation at 94°C for 5 minutes followed by 10 cycles of denaturation at 94°C for 45 seconds, annealing at 55°C for 45 seconds, extension at 72°C for 90 seconds, then continued with 25 cycles of denaturation at 94°C for 45 seconds,

annealing at 50°C for 45 seconds, extension at 72°C for 90 seconds, and final extension 72°C 10 minutes. The amplicons were visualized in 0.8% agarose stained using Sybr safe DNA (Invitrogen), and images were obtained using a gel documentation system.

### RESULTS AND DISCUSSION


A total of 100 festering wound swab samples were obtained from hospitalized patients in Raden Mattaher hospital, dr. Bratanata hospital, and Kambang hospital. Thirty-two samples were positive *S. aureus* through staphylect plus test. There were 14 isolates of MRSA based on cefoxitin resistance in disc diffusion method and PCR mecA positive (Figure 1).

Multiplex PCR was performed on 14 MRSA isolates to identify the type of SCCmec in the samples. There were 12 samples (85.72%) of SCCmec type III, 1 sample (7.14%) of SCCmec type II, and 1 sample (7.14%) of SCCmec type IVb (Figure 2).


The SCCmec types distribution were depended on geographical manner. Most MRSA isolates from Eastern and Middle Eastern countries hospitals contain SCCmec type III.<sup>15</sup> This SCCmec type is common in some South East Asia countries hospitals such as Thailand, Singapore, Indonesia and Malaysia.<sup>16</sup> Different with some South East Asian countries, MRSA isolates from

**Table 1.** Sequence of oligonucleotide primers.14

Target Gene	Primer	Nucleotide sequence (5'-3')	Amplicon (bp)
MecA gene	MecA1	GTA GAA ATG ACT GAA CGT CCG ATA A	310
	MecA2	CCA ATT CCA CAT TGT TTC GGT CTA A	
SCCmec I	I-F	GCT TTA AAG AGT GTC GTT ACA GG	613
	I-R	GTTCTCTCATAGTATGACGTCC	
SCCmec II	II-F	CGTTGAAGATGATGAAGCG	398
	II-R	CGAAATCAATGGTTAATGGACC	
SCCmec III	III-F	CCATATTGTGTACGATGCG	280
	III-R	CCTTAGTTGTCGTAACAGATCG	
SCCmec IVa	IVa-F	GCCTTATTCGAAGAAACCG	776
	IVa-R	CTACTCTTCTGAAAAGCGTCG	
SCCmec IVb	IVb-F	TCTGGAATTACTTCAGCTGC	493
	IVb-R	AAACAATATTGCTCTCCCTC	



**Figure 1.** Agarose gel electrophoresis of PCR product amplified from *MecA* gene (310 bp). M is DNA marker; K(+) is positive control, Lane 1-14 are *MecA* fragments.



**Figure 2.** Agarose gel electrophoresis of PCR product amplified from *SCCmec* type. M is DNA marker; Lane 1-3,5,7-14 are *SCCmec* type III fragments (280 bp). Lane 4 is *SCCmec* type II fragment (398 bp). Lane 6 is *SCCmec* type IVb fragment (493 bp).

Korea and Japan predominantly contain *SCCmec* type II.<sup>16</sup> While some European countries MRSA isolates contain *SCCmec* type IV.<sup>17</sup>

In this study, the majority of *SCCmec* types was type III (85.72%). These results were consistent with studies conducted in seven countries in Asia including Indonesia and studies conducted in Iran where *SCCmec* type III was the most common in MRSA isolates.<sup>16,18,19</sup>

In addition to *SCCmec* type III, this study also found a small proportion of MRSA isolates contained *SCCmec* type II and type IVb. *SCCmec* type I, II, and III were the commonly found types in hospitals (HA-MRSA), while *SCCmec* type IV and V were the commonly found types in communities (CA-MRSA).<sup>20,21,22</sup>

*SCCmec* type II also found in Jakarta, a study mentioned that the majority of MRSA isolates in hospitals were *SCCmec* type II.<sup>23</sup> While

*SCCmec* type IV also found in Denpasar (12.5%) and Malaysia (3.18%) among MRSA isolates in hospitals.<sup>24,25</sup> This means that there has been a shift in the content of *SCCmec* in MRSA isolates in Indonesia. The discovery of *SCCmec* type IV in the Hospital raises concerns because this type is more mobile, generally causes more severe clinical symptoms, and is more difficult in the selection of suitable antibiotics.<sup>21,24</sup> In comparison to other *SCCmec* elements, *SCCmec* IV is small in size and more variable, which has possibly enabled it to spread easily within *S. aureus*.

## CONCLUSIONS

Based on the results revealed in this study, there has been a change in the type of *SCCmec* in MRSA isolates from hospitals. Therefore, it is

recommended to conduct further research with a larger sample size, both from hospitals and communities to identify the SCCmec type and its relationship to patterns of sensitivity to antibiotics. Keeping in view, the finding of SCCmec type IV in Jambi should be investigated, whether it is a circulator or a persisting invader. Further molecular analysis of these MRSA isolates by pulsed-field gel electrophoresis or MLST (Multi Locus Sequence Typing) may provide much useful information regarding the origin and the epidemiology of local isolates.

## ACKNOWLEDGEMENT

This project was funded by a grant from the Faculty of Medicine and Health Sciences, Jambi University.

## CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

## REFERENCES

1. Alrabiah K, Al Alola S, Al Banyan E, Al Shaalan M, Al Johani S. Characteristics and risk factors of hospital acquired e Methicillin-resistant *Staphylococcus aureus* (HA-MRSA) infection of pediatric patients in a tertiary care hospital in Riyadh, Saudi Arabia. *Int J Pediatr Adolesc Med.* 2016; 3(2): 71–7.
2. Thomas R, Ferguson J, Coombs G, Gibson PG. Community-acquired methicillin-resistant *Staphylococcus aureus* pneumonia: A clinical audit. *Asian Pacific Soc Respiriol.* 2011; 16: 926–31.
3. Bukharie HA. A review of community-acquired methicillin-resistant *Staphylococcus aureus* for primary care physicians. *J Fam Community Med.* 2010; 17(3): 117–20.
4. Mahmudah R, Soleha TU, Ekowati C. Identifikasi Methicillin Resistant *Staphylococcus aureus* (MRSA) pada Tenaga Medis dan Paramedis di Ruang Intensive Care Unit (ICU) dan Ruang Perawatan Bedah Rumah Sakit Umum Daerah Abdoel Moeloek. *Med J Lampung Univ.* 2013; 2(4): 70–8.
5. Liana P. Gambaran Kuman Methicillin-Resistant *Staphylococcus Aureus* (MRSA) di Laboratorium Mikrobiologi Departemen Patologi Klinik Rumah Sakit Dr. Cipto Mangunkusumo (RSCM) Periode Januari-Desember 2010. *MKS.* 2014; 46(3): 171–5.
6. Hill-cawthorne GA, Hudson LO, Fouad M, El A, Piepenburg O, *et al.* Recombinations in Staphylococcal Cassette Chromosome mec Elements Compromise the Molecular Detection of Methicillin Resistance in *Staphylococcus aureus*. *PLoS One.* 2014; 9(6).
7. Paterson GK, Harrison EM, Holmes MA. The emergence of mecC methicillin-resistant *Staphylococcus aureus*. *Trends Microbiol.* 2014; 22(1): 42–7.
8. Ito T, Hiramatsu K, Oliveira DC, De Lencastre H, Zhang K, Westh H, *et al.* Classification of staphylococcal cassette chromosome mec (SCCmec): Guidelines for reporting novel SCCmec elements. *Antimicrob Agents Chemother.* 2009; 53(12): 4961–7.
9. Kaya H, Hasman H, Larsen J, Stegger M, Johannesen B. SCCmecFinder, a Web-Based Tool for Typing of Staphylococcal Cassette Chromosome mec in *Staphylococcus aureus* Using Whole-Genome Sequence Data. *Am Soc Microbiol.* 2018; 3(1): 1–9.
10. Nitschke H, Pfohl K, Monecke S, Jatzwauk L, Mu E, *et al.* Diversity of SCC mec Elements in *Staphylococcus aureus* as Observed in South-Eastern Germany. *PLoS One.* 2016; 11(9): 1–24.
11. Yuwono, Sunarjati S, Masria S, Supardi I. *Staphylococcus aureus* dengan Polymerase Chain Reaction Identification of Staphylococcal Cassette Chromosome Mec Methicillin Resistant *Staphylococcus aureus* Using Polymerase Chain Reaction. *Maj Kedokt Bandung.* 2009; 43(2): 60–5.
12. Sudigdoadi S. Analisis Tipe Staphylococcal Cassette Chromosome mec (SCCmec) Isolat Methicillin Resistant *Staphylococcus aureus* (MRSA). *Maj Kedokt Bandung.* 2014; 42(4): 149–54.
13. Cockerill FR, Wikler MA, Bush K, Craig WA, Dudley MN, Eliopoulos GM, *et al.* Performance Standards for Antimicrobial Susceptibility Testing ; Twenty-First Informational Supplement. Vol. 31, CLSI document. 2011. M100-S21 p.
14. McClure-Warnier J-A, Conly JM, Zhang K. Multiplex PCR Assay for Typing of Staphylococcal Cassette Chromosome Mec Types I to V in Methicillin-resistant *Staphylococcus aureus*. *J Vis Exp.* 2013; (79).
15. Holden MTG, Hsu L, Kurt K, Weinert LA, Mather AE. A genomic portrait of the emergence, evolution, and global spread of a methicillin-resistant *Staphylococcus aureus* pandemic. *Cold Spring Harb Lab Press.* 2013; 23: 653–64.
16. Asghar AH. Molecular characterization of methicillin-resistant *Staphylococcus aureus* isolated from tertiary care hospitals. *Pak J Med Sci.* 2014; 30(4): 698–702.
17. Kinnevey PM, Shore AC, Brennan GI, Sullivan DJ. Extensive Genetic Diversity Identified among Sporadic Methicillin-Resistant *Staphylococcus aureus* Isolates Recovered in Irish Hospitals between 2000 and 2012. *Antimicrob Agents Chemother.* 2014; 58(4): 1907–17.

18. Ghanbari F, Saberianpour S, Ghanbari N. Staphylococcal Cassette Chromosome mec (SCC mec) Typing of Methicillin-Resistant *Staphylococcus aureus* Strains Isolated from Community-and Hospital-Acquired Infections. *Avicenna J Clin Microb Infection*. 2017; 4(2).
19. Peters B, Liu J, Chen D, Peters BM, Li L, *et al.* Staphylococcal chromosomal cassettes mec (SCCmec): A mobile genetic element in methicillin-resistant *Staphylococcus aureus* *Microbial Pathogenesis Staphylococcal chromosomal cassettes mec (SCCmec): A mobile genetic element in methicillin-resistant* *Staphylococcus aureus* *Microb Pathog*. 2016; 101 (July 2018): 56–67.
20. Ahmad N, Ruzan IN, Kamel M, Ghani A, Hussin A, Nawi S, *et al.* Characteristics of community- and hospital- acquired methicillin-resistant *Staphylococcus aureus* strains carrying SCC mec type IV isolated in Malaysia. *J Med Microbiol*. 2009; 58: 1213–8.
21. Ouchenane Z, Smati F, Rolain J, Raoult D. Molecular characterization of methicillin-resistant *Staphylococcus aureus* isolates in Algeria. *Pathol Biol*. 2011; 59: e129–32.
22. Monecke S, Schwarz S, Hotzel H, Ehrlich R. Rapid Microarray-Based Identification of Different mecA Alleles in. *Antimicrob Agents Chemother*. 2012; 56(11): 5547–54.
23. Sabir M, Dwiyanti R, Hatta M, Buntaran L, Sultan AR. Scmec type II gene is common among clinical isolates of methicillin-resistant *Staphylococcus aureus* in Jakarta, Indonesia. *BMC Res Notes*. 2013; 6(1): 110.
24. Santosaningsih D, Santoso S, Setijowati N, Rasyid HA, Budayanti NS, *et al.* Prevalence and characterisation of *Staphylococcus aureus* causing community-acquired skin and soft tissue infections on Java and Bali, Indonesia. *Tropical Medicine and International Health*. 2018; 23(1): 34–44.
25. Hannan A, Javed F, Saleem S, Tahira K, Jahan S. Frequency of Staphylococcal Cassette Chromosome mec Type IV and Type V in Clinical Isolates of Methicillin Resistant *Staphylococcus aureus*. *Open J Med Microbiol*. 2015; 5(June): 69–75.