


Original Article

# IJTID

(INDONESIAN JOURNAL OF TROPICAL AND INFECTIOUS DISEASE)

Scientific Journal of Tropical and Infectious Disease



## Profile of Nontuberculous Mycobacteria and *Mycobacterium Tuberculosis* Detected in the Sputum of Pulmonary Tuberculosis Retreatment Patients at Dr. Soetomo General Hospital

Mochammad Afif Ziaulhaq<sup>1</sup>, Ni Made Mertaniasih<sup>2,3\*</sup>, Resti Yudhawati Meliana<sup>4,3</sup>, Ariani Permatasari<sup>4,3</sup>

<sup>1</sup>Medical Program, Faculty of Medicine, Universitas Airlangga, Surabaya Indonesia

<sup>2</sup>Department of Medical Microbiology, Faculty of Medicine, Universitas Airlangga, Surabaya Indonesia

<sup>3</sup>Dr. Soetomo Academic Hospital, Surabaya Indonesia

<sup>4</sup>Department of Pulmonology and Medical Respirology, Faculty of Medicine, Universitas Airlangga, Surabaya Indonesia



### Abstract

#### ARTICLE INFO

Received: November 6, 2024

Accepted: December 13, 2024

Published: April 30, 2025

Available online: April 30, 2025

\*) Corresponding author:

E-mail: [ni-made-m@fk.unair.ac.id](mailto:ni-made-m@fk.unair.ac.id)

#### Keywords:

Nontuberculous mycobacteria

*Mycobacterium tuberculosis*

Pulmonary tuberculosis

Retreatment patients

Retrospective design

Tuberculosis (TB) remains one of the leading infectious diseases worldwide. Despite global efforts to control TB, it remains a major public health issue, affecting 10.6 million people annually in 2021, with significant morbidity and mortality, particularly in resource-limited settings. Effective treatment of TB requires strict adherence to long-term medication, but challenges such as treatment failure, relapse, and loss to follow-up complicate outcomes. This is especially concerning for patients with comorbidities such as diabetes, HIV, or hypertension, which not only increase the risk of TB but also hinder its treatment and elevate the likelihood of nontuberculous mycobacteria (NTM) infections. This study aimed to analyze 326 pulmonary TB retreatment cases at Dr. Soetomo General Academic Hospital from October 2023 to April 2024. The retrospective design identified that 323 cases involved MTB and 3 involved NTM. The findings show that loss to follow-up was the most common reason for retreatment, particularly among males and older adults. Comorbidities were found to exacerbate treatment challenges, with some retreatment cases lasting up to 24 months. The study concludes that loss to follow-up remains a major risk factor for TB retreatment, particularly in MTB cases, and highlights the importance of managing comorbidities to improve treatment outcomes.



This is an open access article under the CC BY-NC-SA license

(<https://creativecommons.org/licenses/by-nc-sa/4.0/>)

Cite this as: Ziaulhaq, M. A., Mertaniasih, N. M., Meliana, R. Y., and Permatasari, A. (2025). Profile of Nontuberculous Mycobacteria and *Mycobacterium Tuberculosis* Detected in the Sputum of Pulmonary Tuberculosis Retreatment Patients at Dr. Soetomo General Hospital. *Indonesian Journal of Tropical and Infectious Disease*, 13(1) : 79–88. <https://doi.org/10.20473/ijtid.v13i1.64176>

## INTRODUCTION

Tuberculosis (TB), caused by *Mycobacterium tuberculosis* (MTB), remains one of the most critical public health issues worldwide.<sup>1</sup> In 2021, the World Health Organization (WHO) reported 10.6 million TB cases globally, with 969,000 cases occurring in Indonesia, marking an increase from 2020's estimate of 10 million cases. TB treatment requires strict adherence to a long-term regimen of antituberculosis drugs, and any interruptions in treatment can lead to complications such as treatment failure, relapse, or loss to follow-up.<sup>2</sup> When these outcomes occur, patients must undergo retreatment to achieve complete bacterial eradication. In 2021, the treatment success rate (TSR) in East Java was 89.13%, indicating that 10.87% of cases still required retreatment.<sup>3</sup>

Comorbid conditions such as HIV, malignancies, diabetes, and other chronic diseases further complicate TB treatment by weakening the immune system. HIV-1 targets CD4+ T cells and macrophages, while *Mycobacterium tuberculosis* mainly impacts macrophages, which rely on CD4+ T cells to help eliminate intracellular pathogens. Consequently, the decline in CD4+ T cells due to HIV-1 infection is believed to significantly contribute to the heightened risk of tuberculosis in individuals infected with HIV-1.<sup>4</sup> Associated mechanisms of TB and malignancy lead to a persistent and intense inflammatory response contributes to the development and advancement of cancer through various mechanisms. Multiple theories connect DNA damage in lung epithelial cells to free radical-induced harm and the sustenance of a cytokine network that is both pro-inflammatory and immunosuppressive.<sup>5</sup> Several biological factors can weaken the immune system and increase the risk of tuberculosis in

people with diabetes (DM). High insulin levels are linked to reduced T helper 1 (Th1) immunity, lowering the T helper 1 to T helper 2 cell ratio and the interferon-gamma (IFN- $\gamma$ ) to interleukin-4 (IL-4) ratio. People with diabetes also have lower IFN- $\gamma$  levels, which are negatively correlated with HbA1c, a blood sugar measure. Additionally, neutrophils in diabetics show decreased movement and bacteria-killing ability.<sup>6</sup> Studies have shown that TB patients with comorbid diabetes mellitus are ten times more likely to have poor TB treatment outcomes than people without diabetes mellitus due to diminished treatment responses.<sup>7,8</sup> Moreover, patients with a history of comorbidities such as chronic obstructive pulmonary disease (COPD), bronchiectasis, tumors, or immunosuppressant use are at a heightened risk of developing nontuberculous mycobacteria (NTM) infections.<sup>9</sup>

Nontuberculous mycobacteria (NTM) are becoming recognized as a crucial opportunistic pathogen for humans. NTM mostly affect the lungs and may lead to progressive disease in susceptible hosts, mainly in people with chronic lung diseases, including COPD, bronchiectasis, cystic fibrosis, and a history of tuberculosis.<sup>10</sup> The most prevalent nontuberculous mycobacterial species includes slow-growing such as *Mycobacterium avium* complex and *Mycobacterium kansasii*, as well as fast-growing such as *Mycobacterium abscessus* and *Mycobacterium chelonae*.<sup>11</sup>

Despite advances in TB treatment, there remains a significant gap in understanding the relationship between comorbidities, NTM infections, and treatment outcomes in TB retreatment cases. Current research has not fully explored the profile of NTM and MTB in TB retreatment patients, particularly in

Indonesia. Addressing this gap is crucial to improving treatment success rates and reducing retreatment cases.

This research provides valuable insights into the challenges faced by patients, particularly those with comorbidities like diabetes and HIV. Analyzing 326 cases revealed that loss to follow-up was the primary reason for retreatment, especially among older males. The study underscores the significant impact of comorbidities on treatment outcomes, demonstrating that effective management of these conditions is crucial for improving recovery rates in TB patients. This research highlights the need for targeted interventions to enhance treatment adherence and ultimately reduce TB-related complications.

## MATERIALS AND METHODS

### Population and Sample

This study utilizes a comprehensive dataset derived from secondary data, specifically the medical records of pulmonary tuberculosis (TB) retreatment patients at Dr. Soetomo General Academic Hospital, covering the period from January 2020 to March 2023. The dataset includes detailed records for patients aged 18 years and older who underwent sputum examination and met specific inclusion criteria. This approach ensures a robust and complete dataset for analysis. Cases with incomplete data were excluded to enhance the reliability and validity of the findings.

### Methods

A descriptive retrospective design was employed to investigate the profiles of nontuberculous mycobacteria (NTM) and *Mycobacterium tuberculosis* (MTB) identified in the sputum samples of retreatment TB patients. Data were systematically organized and analyzed

using Microsoft Excel (version 2406), facilitating a detailed breakdown of key variables, including demographic information, sputum culture results, and treatment outcomes. Statistical analyses were performed to identify trends and patterns within the patient population, allowing for a comprehensive understanding of the prevalence of NTM and MTB in the studied cohort.

## RESULTS AND DISCUSSION

Table 1 shows that 28.4% of pulmonary TB cases required retreatment, largely due to relapse, treatment failure, or loss to follow-up, consistent with global studies.<sup>12</sup> Various factors contribute, including limited healthcare access, socioeconomic challenges, non-adherence to treatment, comorbidities like HIV or diabetes, and exogenous reinfections.<sup>13</sup>

The majority (99.07%) of retreated TB patients were infected by MTB, with only 0.93% by NTM, similar to findings by Limo (2016) in Kenya. NTM infections mimic MTB symptoms, such as chronic cough, fever, and weight loss, complicating diagnosis, especially in TB-endemic regions.<sup>14</sup> Additionally, NTM's slow growth makes diagnosis challenging.<sup>15</sup>

Drug resistance was found in 42.11% of the patients, presenting a significant challenge, as noted by Adé et al. (2016). The occurrence of drug resistance is often due to biological and clinical interactions, including poor compliance, and molecular mechanisms.<sup>16,17</sup>

Of those with NTM, 100% were drug sensitive (DS TB) patients, with previous TB history increasing susceptibility.<sup>18</sup> Co-infection with NTM and MTB poses a greater risk, particularly in those with pulmonary cavitation, which may act as a reservoir for NTM.<sup>19</sup>

**Table 1.** Proportion of new and retreated pulmonary TB patients, the mycobacterium, and the drug classification

	Amount	Percentage
<b>Proportion</b>		
TB	822	71.6
TB retreatment	326	28.4
<b>Mycobacterium</b>		
MTB	323	99.07
NTM	3	0.93
<b>Drug sensitive</b>		
MTB	187	57.89
NTM	3	100
<b>Drug resistance</b>		
MTB	136	42.11
NTM	0	0

The analysis of retreated pulmonary TB patients based on Table 2 reveals that the majority of cases occurred in adults aged 19-59 years, with 82.02% of MTB infections and 66.67% of NTM infections found in this age group. These findings align with prior research indicating that TB is most prevalent in individuals of productive age.<sup>20,21</sup> High mobility and socioeconomic factors contribute to the spread and severity of the disease, increasing both morbidity and mortality rates in this demographic.<sup>22</sup>

In contrast, adolescents (under 18 years old) represented only a small proportion of cases, with 0.62% for MTB and 33.33% for NTM. Risk factors like diabetes, smoking, and poor living conditions exacerbate the susceptibility of this age group to TB.<sup>18</sup> Meanwhile, elderly patients (over 60 years old) accounted for 17.67% of MTB infections, with no NTM cases. Elderly individuals, due to weakened immune systems and comorbidities, are particularly vulnerable to TB, leading to higher infection rates.<sup>23</sup>

Regarding gender, men represented 59.4% of MTB cases and 66% of NTM cases. This data are consistent with global trends, where TB and NTM infections are more prevalent in males.<sup>24</sup> Alcohol abuse is much higher in men than women.

Furthermore, smoking is the most significant risk factor for COPD and lung cancer and is related to pulmonary TB. Overall, men smoke more than women. As a result, alcohol abuse and smoking are greater contributors to the TB disease burden in men.<sup>25,26</sup>

**Table 2.** Retreated pulmonary TB patients by age and gender

	Amount	Percentage
<b>Age (MTB)</b>		
Teenager (18 years)	2	0.62
Adult (19-59 years)	264	82.02
Elderly (60+ years)	57	17.67
<b>Age (NTM)</b>		
Teenager (18 years)	1	33.33
Adult (19-59 years)	2	66.67
Elderly (60+ years)	0	0
<b>Gender (MTB)</b>		
Male	192	59.4
Female	131	40.6
<b>Gender (NTM)</b>		
Male	2	66
Female	1	34

The data in Table 3 highlight common comorbidities in pulmonary TB patients. Diabetes mellitus affected 18.2% of MTB patients, which impairs immune function, leading to higher TB retreatment rates. This is consistent with findings from Siddiqui et al. (2016) and Tan et al. (2021), emphasizing how hyperglycemia disrupts immune responses and facilitates TB progression.<sup>27</sup>

Additionally, HIV/AIDS was present in 8.9% of MTB cases, a well-known risk factor due to its detrimental effect on T-cells and macrophages. Research by Mahtab and Coetzee (2017) and Tan et al. (2021) confirms that co-infection accelerates disease progression, making TB harder to control.<sup>28</sup> In HIV-1



infection, it targets CD4+ T cells and macrophages. *Mycobacterium tuberculosis* primarily impacts macrophages, which depend on CD4+ T cells to help eliminate intracellular pathogens. As a result, the decrease in CD4+ T cells due to HIV-1 infection is believed to significantly contribute to the increased risk of tuberculosis in HIV-1-infected individuals.<sup>4</sup> Hypertension was less common but still significant, as immune response disruptions can lead to pulmonary hypertension, supported by findings from Huamán et al. (2015) and Bhattacharyya et al. (2016). It is hypothesized that tuberculosis can lead to hypertension by triggering immune responses that impair endothelial function, increasing the risk of cardiovascular disease (CVD) and hypertension. TB may cause damage to lung tissue, affecting vascular structures and leading to conditions like vasculitis and endarteritis, which can reduce the cross-sectional area of pulmonary blood vessels and result in pulmonary hypertension. Additionally, if TB infects the kidneys, it can damage renal tissue, decrease kidney function, and impair blood pressure regulation, potentially causing hypertension.<sup>29</sup>

For NTM patients, comorbidities like hepatitis B and pituitary gland tumors were reported in 33% of cases. Studies by Khan et al. (2021) highlight the risks of drug-induced hepatotoxicity in co-infected patients. Tumors, meanwhile, worsen TB reactivation due to immune system suppression.<sup>30</sup> A persistent and intense inflammatory response contributes to the onset and advancement of cancer through various mechanisms. Multiple theories connect DNA damage in lung epithelial cells to free radical-induced harm and the establishment of a cytokine network that is both pro-inflammatory and immunosuppressive.<sup>5</sup> Lastly, the

relationship between tumors and TB highlights a bi-directional link where both diseases facilitate each other's progression.<sup>31</sup>

**Table 3.** Comorbidities of pulmonary TB patients on retreatment

	Amount	Percentage
<b>Comorbidities (MTB)</b>		
Diabetes Mellitus	59	18.2
HIV	29	8.9
Hypertension	3	0.92
<b>Comorbidities (NTM)</b>		
Hepatitis B	1	33.33
Tumor	1	33.33

Based on Table 4, the majority of pulmonary TB patients on retreatment due to MTB faced loss to follow-up, accounting for 49.5%, followed by 37.5% experiencing treatment failure and 13% relapse. For patients with pulmonary TB caused by NTM, 60% encountered treatment failure, and 40% faced loss to follow-up, with no recorded relapses.

These findings align with research by Lin et al. (2021), who reported 15.9% loss to follow-up and 0.3% treatment failure. Additionally, research by Dedefo et al. (2019) in Ethiopia found 72.6% relapse, 19.6% treatment failure, and 7.8% loss to follow-up cases.

**Table 4.** TB treatment status of the patient

	Amount	Percentage
<b>TB Status (MTB)</b>		
Treatment failure	159	37.5
Relapse	54	13
Loss to follow up	207	49.5
<b>TB Status (NTM)</b>		
Treatment failure	3	60
Relapse	0	0
Loss to follow up	2	40

Loss to follow-up in tuberculosis cases can result from factors such as limited healthcare access, socioeconomic status, lack of family support, and non-compliance with treatment.<sup>10</sup> Risk factors for treatment failure and relapse include HIV infection, diabetes, malnutrition, and drug resistance.<sup>32,33</sup>

The data in Table 5 highlight that the longest duration of treatment for pulmonary TB was 24 months. Most drug sensitive (DS TB) patients had retreatment durations of either six months (45.26%) or 12 months (37.36%). For DR-TB (drug-resistant TB), most patients had retreatment durations of 11 months (56.61%) or 20 months (24.26%).

**Table 5.** Duration of treatment for pulmonary TB patients

	Amount	Percentage
<b>Duration (Drug sensitive TB)</b>		
6 months	86	45.26
7 months	9	4.73
8 months	10	5.26
9 months	1	0.52
11 months	2	1.05
12 months	71	37.36
13 months	4	2.10
15 months	1	0.52
16 months	1	0.52
18 months	2	1.05
20 months	1	0.52
22 months	1	0.52
23 months	1	0.52
<b>Duration (Drug-resistant TB)</b>		
6 months	1	0.73
9 months	5	3.67
10 months	9	6.61
11 months	77	56.61
16 months	1	0.73
18 months	8	5.88
20 months	33	24.26
21 months	1	0.73
24 months	1	0.73

Treatment duration varies due to the need for adjustments based on how patients respond to therapy. Soeroto et al. (2022) found that 43.08% of DR-TB patients required shorter treatments (9-11 months). But, the next research found

91.62% of patients with long-standing DR-TB needed longer treatment due to complications like culture conversion time or comorbidities such as malnutrition, HIV, and CKD. These factors significantly reduce the treatment success rate.<sup>34</sup>

For NTM infections caused by *Mycobacterium avium* complex, treatment typically includes three drugs—macrolides, ethambutol, and rifampicin—and must continue for 12 months after culture conversion.<sup>35</sup> These findings underline the importance of tailoring TB treatment based on specific patient factors to optimize recovery outcomes.

## STRENGTH AND LIMITATION

The strength of this study lies in its focus on retreatment patients affected by *Mycobacterium tuberculosis* and nontuberculous mycobacteria, providing valuable insights into both types of infections. However, the research was limited by the inability to determine the specific NTM species, types of drug resistance, and the total number of patient fatalities.

## CONCLUSIONS

The primary risk factor for retreated pulmonary tuberculosis, particularly in patients infected with MTB, is a high rate of loss to follow-up. Additionally, comorbid conditions exacerbate the outcomes, further complicating treatment success.

## ACKNOWLEDGMENT

This research was supported by the Faculty of Medicine University Airlangga, Surabaya and Dr. Soetomo Academic Hospital, Surabaya.

## ETHICAL CLEARANCE

The research protocol and use of medical record data was approved by Dr. Soetomo Surabaya General Hospital ethics committee (Ref. No.: 1459/LOE/301.4.2/IX/2023).

## FUNDING

This research received no funding

## CONFLICT OF INTEREST

The authors declare no conflicts of interest in relation to this study.

## AUTHOR CONTRIBUTION

All authors were actively involved in the study design, drafting, and approval for publication.

## REFERENCES

1. WHO. Global Tuberculosis report. WHO, 2022.
2. Getnet F, Sileshi H, Seifu W, Yirga S, Alemu AS. Do retreatment tuberculosis patients need special treatment response follow-up beyond the standard regimen? Finding of five-year retrospective study in pastoralist setting. *BMC Infect Dis*, 2017;17(1). doi:10.1186/s12879-017-2882-y
3. Dinas Kesehatan Provinsi Jawa Timur. Profil Kesehatan 2021.
4. Bell L, Noursadeghi M. Pathogenesis of HIV-1 and *Mycobacterium tuberculosis* co-infection. *Nat Rev Microbiol*. 2018;16(2):80–90. doi: 10.1038/nrmicro.2017.128
5. Molina-Romero C, Arrieta O, Hernández-Pando R. Tuberculosis and lung cancer. *Salud Publica Mex*. 2019 May-Jun;61(3):286–91. doi: 10.21149/10090
6. Al-Rifai RH, Pearson F, Critchley JA, Abu-Raddad LJ. Association between diabetes mellitus and active tuberculosis: A systematic review and meta-analysis. *PLoS One*. 2017 Nov 21;12(11):e0187967. doi: 10.1371/journal.pone.0187967
7. Adane HT, Howe RC, Wassie L, Magee MJ. Diabetes mellitus is associated with an increased risk of unsuccessful treatment outcomes among drug-susceptible tuberculosis patients in Ethiopia: A prospective health facility-based study. *J Clin Tuberc Other Mycobact Dis*. 2023 Apr 11;31:100368. doi: 10.1016/j.jctube.2023.100368
8. Gopalaswamy R, Shanmugam S, Mondal R, Subbian S. Of tuberculosis and non-tuberculous mycobacterial infections - A comparative analysis of epidemiology, diagnosis and treatment. *J Biomed Sci*. 2020; 27(1). doi:10.1186/s12929-020-00667-6
9. Loebinger MR, Quint JK, van der Laan R, Obradovic M, Chawla R, Kishore A, et al. Risk factors for nontuberculous mycobacterial pulmonary disease/ *Chest*, 2023;164(5): 1115–24. doi:10.1016/j.chest.2023.06.014
10. Cowman S, van Ingen J, Griffith DE, Loebinger MR. Non-tuberculous mycobacterial pulmonary disease. *Eur Respir J*. 2019;54(1):1900250. doi: 10.1183/13993003.00250-2019
11. Drummond WK, Kasperbauer SH. Nontuberculous *Mycobacteria*: Epidemiology and the impact on pulmonary and cardiac disease.

- Thorac Surg Clin. 2019 Feb;29(1):59-64. doi: 10.1016/j.thorsurg.2018.09.006
12. Sarpal SS, Goel NK, Kumar D, Janmeja AK. Gender disparities in pengobatan ulang patients of tuberculosis: A north Indian study. J Nat Sci Biol Med. 2015 Jan-Jun;6(1):63-6. doi: 10.4103/0976-9668.149087
13. Santos E, Felgueiras Ó, Oliveira O, Duarte R. Factors associated with loss to follow-up in Tuberculosis treatment in the Huambo Province, Angola. Pulmonology 2019;25(3):190-2. doi:10.1016/j.pulmoe.2019.03.003
14. Ryu YJ, Koh W J, Daley CL. Diagnosis and Treatment of Nontuberculous Mycobacterial Lung Disease: Clinicians' Perspectives. Tuberc Respir Dis, 2016; 79(2):, 74. doi:10.4046/TRD.2016.79.2.74
15. Gnanadurai R, Ninan MM, Arul AO, Sam AS, James P, Gupta R, et al. Challenges in the management of slowly growing non-tuberculous mycobacteria causing pulmonary disease: Perspectives from a high burden country. Indian J Med Microbiol. 2021 Oct-Dec;39(4):446-50. doi: 10.1016/j.ijmmb.2021.07.005
16. Khawbung JL, Nath D, Chakraborty S. Drug resistant Tuberculosis: A review. Comp Immunol Microbiol Infect Dis. 2021 Feb; 74:101574. doi: 10.1016/j.cimid.2020.101574
17. Strydom N, Gupta SV, Fox WS, Via LE, Bang H, Lee M, et al. 'Tuberculosis drugs' distribution and emergence of resistance in patient's lung lesions: A mechanistic model and tool for regimen and dose optimization,' PLoS Medicine, 2019; 16(4): e1002773. doi:10.1371/journal.pmed.1002773
18. Tan Y, Deng Y, Yan X, Liu F, Tan Y, Wang Q, et al. Nontuberculous mycobacterial pulmonary disease and associated risk factors in China: A prospective surveillance study. J Infect.2012; 83(1): 46-53. doi:10.1016/j.jinf.2021.05.019
19. Chu, H, Zhao L, Xiao H, Zhang Z, Zhang J, Gui T, et al. (Systematic review/Meta-analysis Prevalence of nontuberculous mycobacteria in patients with bronchiectasis: a meta-analysis/ Arch Med Sci. 2014; (10(4): 661-8. doi:10.5114/aoms.2014.44857
20. Adé S, Adjibodé O, Wachinou P, Toundoh N, Awanou B, Agodokpessi G, et al. Characteristics and treatment outcomes of pengobatan ulang tuberculosis patients in Benin. Tuberc Res Treat, 2016; 1-7. doi:10.1155/2016/1468631
21. Saptawati L, Kusumo H, Suryawati B. Prevalence of non-tuberculous mycobacteria (NTM) in Surakarta, Indonesia: higher than expected. KnE Life Sci. (2019); 4(12):132. doi:10.18502/cls.v4i12.4166
22. Nidoi J, Muttamba W, Walusimbi S, Imoko JF, Lochoro P, Ictho J, et al. Impact of socio-economic factors on Tuberculosis treatment outcomes in north-eastern Uganda: A mixed methods study. BMC Pub Health. 2021 Nov 26;21(1):2167. doi: 10.1186/s12889-021-12056-1
23. Xu J, Li P, Zheng S, Shu W, Pang Y. Prevalence and risk factors of pulmonary nontuberculous mycobacterial infections in the Zhejiang Province of China. Epidemiol Infect. 2019 Sep



- 11;147:e269. doi: 10.1017/S0950268819001626
24. Karthikeyan V, Ganapathy K. Determinants of Categories of TB Pengobatan ulang with Special Reference to Sources of Primary Anti-TB Treatment. *J Health Allied Sci NU*, 2020; 10(3): 116–21. doi:10.1055/s-0040-1716313
  25. Marçôa R, Ribeiro AI, Zão I, Gomes M, Gaio AR, Duarte R. Erratum to: Tuberculosis and gender – factors influencing the risk of tuberculosis among men and women by age group. *Pulmonology*. 2019;25(4):258. doi: 10.1016/j.pulmoe.2018.10.002
  26. Smith GS, Van Den Eeden SK, Baxter R, Shan J, Van Rie A, Herring AH, et al. Cigarette smoking and pulmonary tuberculosis in northern California. *J Epidemiol Community Health*. 2015;69:568-73.
  27. Krishna S, Jacob JJ. Diabetes mellitus and tuberculosis. . In: Feingold KR, Ahmed SF, Anawalt B, Blackman MR, Boyce A, Chrousos G, et al. (eds), *Endotext*. South Dartmouth (MA): 2021.
  28. Bell L, Noursadeghi M. Pathogenesis of HIV-1 and *Mycobacterium tuberculosis* co-infection. *Nat Rev Microbiol*. 2018; 16(2): 80–90. doi:10.1038/nrmicro.2017.128
  29. Seegert AB, Rudolf F, Wejse C, Neupane D. Tuberculosis and hypertension-a systematic review of the literature. *Int J Infect Dis*. 2017 Mar;56:54–61. doi: 10.1016/j.ijid.2016.12.016
  30. Lamplugh ZL, Fan Y. Vascular microenvironment, tumor immunity and immunotherapy,' *Front Immunol*. 2021; 12. doi:10.3389/fimmu.2021.811485
  31. Preda M, Tănase BC, Zob DL, Gheorghe AS, Lungulescu CV, Dumitrescu EA, et al. The bidirectional relationship between pulmonary tuberculosis and lung cancer. *Int J Environ Res Public Health*. 2023 Jan 10;20(2):1282. doi: 10.3390/ijerph20021282
  32. Boadu AA, Asare P, Mosi L, Mantey EA, Gbadamosi KM, Eleeza JA, et al. Tuberculosis and diabetes mellitus: The complexity of the comorbid interactions. *Int J Infect Dis*. 2024;146:107140. doi: 10.1016/j.ijid.2024.107140
  33. Wotale TW, Lelisho ME, Negasa BW, Gelan LG, Adugna AG. Identifying risk factors for recurrent multidrug resistant tuberculosis based on patient's record data from 2016 to 2021: Retrospective study. *Sci Rep*. 2024;14:23912. doi: 10.1038/s41598-024-73209-x
  34. Soeroto AY, Pratiwi C, Santoso P, Lestari BW. Factors affecting outcome of longer regimen multidrug-resistant tuberculosis treatment in West Java Indonesia: A retrospective cohort study. *PloS One*, 2021; 16(2): e0246284. doi:10.1371/journal.pone.0246284
  35. Daley CL, Iaccarino JM, Lange C, Cambau E, Wallace RJ Jr, Andrejak C, et al. Treatment of Nontuberculous mycobacterial Pulmonary Disease: An Official ATS/ERS/ESCMID/IDSA Clinical Practice Guideline. *Eur Respir J*. 2020 Jul 7;56(1):2000535. doi: 10.1183/13993003.00535-2020
  36. Mahtab S, Coetzee D. Influence of HIV and other risk factors on tuberculosis. *SAMJ. South African*

- Med J. 2017; 107(5): 428. doi:/10.7196/samj.2017.v107i5.11271
37. Bhattacharyya P, Saha D, Bhattacharjee PD, Das SK, Bhattacharyya PP, Dey R. Tuberculosis associated pulmonary hypertension: The revelation of a clinical observation. Lung India. 2016 Mar-Apr;33(2):135-9. doi: 10.4103/0970-2113.177433.
38. Huamán MA, Henson D, Ticona E, Sterling TR, Garvy BA. Tuberculosis and cardiovascular disease: linking the epidemics,' Trop Dis. Travel Med Vaccines 2015;1(1).doi: 10.1186/s40794-015-0014-5
39. Khan AF Sajjad A, Mian DA, Tariq MM, Jadoon UK, Abbas M, et al. Co-infection with hepatitis B in tuberculosis patients on anti-tuberculosis treatment and the final outcome. Cureus. 2021 Apr 12;13(4):e14433. doi: 10.7759/cureus.14433
40. Lin Y, Lin H, Xiao L, Chen Y, Meng X, Zeng X, Chang C, Brigden G. (.Tuberculosis recurrence over a 7-year follow-up period in successfully treated patients in a routine program setting in China: a prospective longitudinal study. Int J Infect Dis. 2021 Sep;110:403-409. doi: 10.1016/j.ijid.2021.07.057.
41. Siddiqui A, Khayyam KU. Sharma M. (Effect of Diabetes Mellitus on Tuberculosis Treatment Outcome and Adverse Reactions in Patients Receiving Directly Observed Treatment Strategy in India: A Prospective Study. Biomed Res Int. 2016;2016:7273935.doi: 10.1155/2016/7273935
42. Dedefo MG, Sirata MT, Ejeta BM, Wakjira GB, Fekadu G, Labata BG. Treatment Outcomes of Tuberculosis Retreatment Case and Its Determinants in West Ethiopia. Open Respir Med J. 2019 Dec 31;13:58-64. doi: 10.2174/1874306401913010058