

Jurnal Berkala EPIDEMIOLOGI

PERIODIC EPIDEMIOLOGY JOURNAL

ORIGINAL ARTICLE

ANALYSIS OF TUBERCULOSIS PATIENT CHARACTERISTICS OF GORONTALO CITY HOSPITAL USING K-MEANS CLUSTERING METHOD

Analisis Karakteristik Pasien Tuberkulosis Rumah Sakit Kota Gorontalo Menggunakan Metode K-Means Clustering

Made Hariadi Wijaya¹, Siti Nur Rahmatiya Abas², Ahmad Fahrian Hipmi³, Endang Darmawan⁴, Woro Supadmi⁵, Sugiyarto Surono⁶

¹Department of Pharmacy, Faculty of Pharmacy, Ahmad Dahlan University, Yogyakarta, Indonesia, 55164, <u>madewijaya2301@gmail.com</u>

²Department of Pharmacy, Faculty of Pharmacy, Ahmad Dahlan University, Yogyakarta, Indonesia, 55164, rahmatyaabas.ra@gmail.com

³Department of Pharmacy, Faculty of Pharmacy, Ahmad Dahlan University, Yogyakarta, Indonesia, 55164, <u>fahrianahmad98@gmail.com</u>

⁴Department of Pharmacy, Faculty of Pharmacy, Ahmad Dahlan University, Yogyakarta, Indonesia, 55164, endang.darmawan@pharm.uad.ac.id

⁵Department of Pharmacy, Faculty of Pharmacy, Ahmad Dahlan University, Yogyakarta, Indonesia, 55164, woro.supadmi@pharm.uad.ac.id

⁶Department of Mathematics, Faculty of Mathematics and Natural Sciences, Ahmad Dahlan University, Yogyakarta, Indonesia, 55166, sugiyarto@math.uad.ac.id

Corresponding Author: Sugiyarto Surono, <u>sugiyarto@math.uad.ac.id</u>, Faculty of Mathematics and Natural Sciences, Ahmad Dahlan University, Yogyakarta, 55166, Indonesia

ARTICLE INFO

Article History: Received December, 31th, 2024 Revised form February, 6th, 2025 Accepted March, 19th, 2025 Published online May, 31th, 2025

Keywords:

Tuberculosis; K-Means; Clustering; Patient Characteristics

Kata Kunci: Tuberkulosis; K-Means; Klustering; Karakteristik Pasien

ABSTRACT

Background: Tuberculosis (TB) is a major health problem in Indonesia, especially in Gorontalo, with high spread due to poor ventilation, overcrowding, and unhealthy lifestyles. Purpose: To analyze the characteristics of TB patients in one of Gorontalo City's hospitals using K-Means Clustering. Methods: Data including age, gender, TB history, HIV status, diabetes history, hypertension, drug resistance, drug side effects, and treatment results were analyzed for the number of clusters using the K-Means method because it is effective in grouping data based on similarity, easy to implement, and works well on large datasets. Results: The analysis resulted in three clusters. Cluster 0 (219 individuals): majority female (63.50%), mean age 45.37 years, low address score (0.49), low resistance and therapy (6.40%), no comorbidities, all experienced side effects (100%), and survival rate 4.10%. Cluster 1 (150 individuals): mean age 52.21 years, higher address score (0.77), resistance 7.30%, therapy 5.30%, comorbidities 100%, all experienced adverse events, and survival rate 4.70%. Cluster 2 (98 individuals): mean age 48.58 years, address score 0.65, very low resistance and therapy (2%), no side effects, 42.90% had comorbidities, and the highest survival rate (12.20%). Conclusion: Three clusters were obtained from the analysis using K-Means. Clustering supports specific How to Cite: Wijaya, M. H., Abas, S. N. R., Hipmi, A.F., Darmawan, E., Supadmi, W., & Surono, S. (2025). Analysis of tuberculosis patient characteristics of Gorontalo city hospital using k-means clustering method. *Jurnal Berkala Epidemiologi*, *13*(2), 147–155. https://dx.doi.org/10.20473/jbe.v13i 22025.147–155 interventions such as comorbidity management or intensive surveillance, improving TB control programs in Gorontalo.

©2025 Jurnal Berkala Epidemiologi. Published by Universitas Airlangga. This is an open access article under CC-BY-SA license

ABSTRAK

Latar Belakang: Tuberkulosis (TB) adalah masalah kesehatan utama di Indonesia, terutama di Gorontalo, dengan penyebaran tinggi akibat ventilasi buruk, kepadatan, dan gaya hidup tidak sehat. Tujuan: Menganalisis karakteristik pasien TB di salah satu rumah sakit kota gorontalo menggunakan K-Means Clustering. Metode: Data meliputi usia, jenis kelamin, riwayat TB, status HIV, riwayat diabetes, hipertensi, resistensi obat, efek samping obat, dan hasil pengobatan di analisis jumlah cluster dengan menguakan metode k-means karena efektif dalam mengelompokkan data berdasarkan kemiripan, mudah diterapkan, dan bekerja dengan baik pada dataset berukuran besar. Hasil: Analisis menghasilkan tiga klaster. Klaster 0 (219 individu): mayoritas perempuan (63,50%), usia rata-rata 45,37 tahun, skor alamat rendah (0,49), resistensi dan terapi rendah (6,40%), tanpa komorbid, semua mengalami efek samping (100%), dan survival rate 4,10%. Klaster 1 (150 individu): usia rata-rata 52,21 tahun, skor alamat lebih tinggi (0,77), resistensi 7,30%, terapi 5,30%, komorbid 100%, semua mengalami efek samping, dan survival rate 4,70%. Klaster 2 (98 individu): usia rata-rata 48,58 tahun, skor alamat 0,65, resistensi dan terapi sangat rendah (2%), tanpa efek samping, 42,90% memiliki komorbiditas, dan survival rate tertinggi (12,20%). Simpulan: Terdapat tiga cluster yang di dapatkan dari hasil analisis mengunakann K-Means. Clustering mendukung intervensi spesifik seperti pengelolaan komorbiditas atau pengawasan intensif, meningkatkan program pengendalian TB di Gorontalo.

©2025 Jurnal Berkala Epidemiologi. Penerbit Universitas Airlangga. Jurnal ini dapat diakses secara terbuka dan memiliki lisensi CC-BY-SA

INTRODUCTION

Tuberculosis is the leading cause of death from infectious diseases among adults worldwide (1). An estimated 10.6 million people worldwide fell ill with TB in 2021, an increase of 4.5% from the previous year (2). The disease is a chronic infection caused by Mycobacterium tuberculosis, Basalingappa, K. Symptoms of TB can include a severe cough lasting three weeks or more, chest pain, coughing up blood or phlegm, weakness or fatigue, weight loss, no appetite, chills, fever, and night sweats (3).

Indonesia is one of the countries that have tuberculosis. Indonesia aims to eradicate tuberculosis by 2030, with the theme "Find Tuberculosis, Treat Until Cured" as the main message of TB control efforts (4) About 845,000 Indonesians had tuberculosis in 2019, and about 92,700 people died from it each year(5). About 17.3% of TB patients in Indonesia are 45-54 years old, 16.8% are 25-34 years old, and 16.7% are 15-24 years old (6). Factors such as age, education, stress, depression, stigma, treatment adherence, and social support influence the high number of tuberculosis cases in Indonesia. This is also exacerbated by home environmental conditions such as poor ventilation, overcrowding, and inadequate lighting (7).

One of the provinces with the highest burden of tuberculosis in Indonesia is Gorontalo. Gorontalo province reported 2,873 TB cases, of which Gorontalo City accounted for 643 cases (69.51% of the estimated 925 cases), with Puskesmas Kota Tengah recording the highest number of 92 cases, while Puskesmas Piloloda'a and Hulonthalangi recorded the lowest number of 29 and 36 cases, respectively (8) Similar to the factors affecting tuberculosis cases in Indonesia, age, education, economic status, smoking, and household contact are associated with the incidence of pulmonary tuberculosis in Gorontalo (9).

Precision medicine maximizes healthcare quality by individualizing the healthcare process based on each patient's unique health status, supporting evidence-based decision-making. Differences in patient characteristics can affect response to treatment, so clustering is necessary to identify more effective and personalized treatment patterns. Precision medicine focuses on the individual patient based on their characteristics, genomic information, proteomics, and social environment, aiding in diagnosis and treatment (10). Precision medicine improves health by accounting for individual variability in genes, environment, and lifestyle (11). Patient clustering using clinical and digital data can improve clustering coherence and recommendation accuracy in healthcare (12). One of the clustering features that are often used is K-Means due to its advantages of being efficient in handling large datasets, having a fast computational process, being easy to implement, and producing clear clusters based on data similarity, making it suitable for pattern analysis in patient clustering. Grouping patients into different strata based on risk factors and clinical outcomes is essential for providing appropriate healthcare.

Through this analysis, it is expected to find groups of patients with similar characteristics, which can be used as a basis for designing more appropriate and effective precision medicine treatment strategies for TB patients with comorbidities. The results of this study are expected to make a significant contribution to improving the treatment effectiveness and prognosis of patients with more complex conditions.

METHODS

This study was designed with an exploratory quantitative approach to analyze data on TB patients with comorbidities in Gorontalo Provincial Hospital using the K-Means algorithm. The data used included gender variables divided into male and female, age categorized into less than or equal 50 years and more than 50 years, and classification based on treatment history, included new patients, those who had never undergone TB therapy before; failed patients, those who had undergone treatment but showed no improvement or failed therapy; relapsed patients, those who had recovered or completed treatment but returned to active TB; and treatment withdrawal patients, those who stopped treatment before completion. In addition, the drug resistance history variable was categorized into patients with a history of resistance, who had confirmed resistance to anti-TB drugs previously, and patients without a history of resistance, who had never been diagnosed with resistance to TB drugs.

The next stage is a data cleaning process to ensure the quality and consistency of the information used. Once the cleaning process is complete, the next step is the application of the K-Means algorithm to group patients based on similar characteristics. Determination of the optimal number of clusters is done by the Elbow or Silhouette Score method. This algorithm aims to divide patients into several groups that have similar treatment patterns and responses to therapy. To facilitate the visualization of clustering results, principal component analysis (PCA) is used to reduce the high dimensionality of data into two dimensions. PCA works by transforming the original variables into principal components (PCs) that represent the main patterns in the data. Its function in cluster analysis is to reduce data complexity, remove correlations between variables, and facilitate visualization in two-dimensional form (PC1 and PC2). PCA allows visualization of clustering results on a two-dimensional graph, where each point represents a patient and each cluster is assigned a different color. The platform used to analyze the data is Google Colab, which was chosen for its ease of access, support for Python libraries such as Scikit-learn for the K-Means (K-Means groups data by finding a cluster center (centroid) and updating it until the data in each cluster are most similar to each other) and PCA algorithms, and its ability to efficiently process large data using GPUs or TPUs. The formula for Kmeans is:

$$d_{ik}: \sqrt{\Sigma_j^m} (c_{ij} - c_{kj})$$

Information:

C_ij : Cluster center C_{ki} : Data Figure

In addition, Google Colab supports real-time team collaboration, making it easier to jointly analyze and share research results. This research has obtained ethical permission from the Ahmad Dahlan University Ethics Committee with the number 012405106.

RESULTS

Clustering using K-Means can improve the efficiency and precision of treatment by using clustering methods based on patient characteristics. Characteristics of patients (See Table 1), in the study consisted of female patients (62.75%), with most of them being over 50 years old (62.75%). Based on treatment history, most were new patients (61.66%), followed by relapse patients (30.40%), while treatment failure and dropout cases amounted to 1.92% and 5.9%, respectively. A total of 94.2%

of patients had no history of resistance, and 96% received first-line therapy. Side effects were experienced by 79% of patients, while 41% had comorbidities. Survival rates were very high, with 94% of patients surviving, while 6% did not. These data reflect the predominant profile of patients and the pattern of treatment received.

Table 1

Description	of the	he	charact	eristics	of	TΒ	patients	in
Hospital							-	

Characteristics	Frequency	Percent		
Characteristics	(n = 467)	(%)		
Gender				
Male	293	62.75		
Female	174	37.25		
Age				
≤50 Year	174	37.25		
>50 Year	293	62.75		
Class Based on				
Treatment History				
New	288	61.66		
Failed	9	1.92		
Relapse	142	30.40		
discontinued	28	5.9		
Resistance History				
No	440	94.20		
Yes	27	5.80		
Comorbidities				
No Comorbidities	275	59		
Had	192	41		
Comorbidities				
Tuberculosis				
Therapy Received				
Line 1	443	96		
Line 2	24	4		
Side Effect				
No Side Effects	98	21		
Had Side Effects	369	79		
Survival				
Not Survive	28	6		
Survive	439	9		

There are several characteristics that each cluster possesses. Cluster 0 consisted of 219 individuals with a female majority (63.50%), a mean age of 45.37 years, and a predominantly low address category (mean score 0.49). Classes based on drug use history were predominantly in the early category (mean 1.56), with a history of resistance and low therapy (6.40%), all experienced side effects (100%), no comorbidities, and a survival rate of 4.10%. Cluster 1 contained 150 individuals, with a higher mean age (52.21 years) and a greater

address score (0.77). Resistance and therapy history remained low (7.30% and 5.30%), all experienced adverse events and comorbidities (100%), and the survival rate was 4.70%. Cluster 2, with 98 individuals, had a mean age of 48.58 years, an address score of 0.65, a very low history of resistance and therapy (2%), and no adverse events, but 42.90% had comorbidities, with the highest survival rate (12.20%).

The figure below is a visualization of the clustering results using K-Means, with PCA for dimension reduction, showing three clusters of patients with unique characteristics (see Figure 1). Cluster 0 (blue) has an average age of 45 years, predominantly female (63.50%), low drug resistance (6.40%), and no comorbidities, but all experience side effects with the lowest survival rate (4.10%). Cluster 1 (orange) contained older patients (mean 52 years), also predominantly female (62.00%), all had comorbidities and side effects, with drug resistance of 7.30% and a slightly higher survival rate (4.70%). Cluster 2 (green) showed the best results, with a mean age of 48 years, the lowest drug resistance (2%), 42.90% had comorbidities, no side effects, and the highest survival rate (12.20%). This analysis provides important insights for understanding patient needs and designing more effective therapies according to the characteristics of each cluster.



Figure 1. Visualization of Cluster Analysis Results

In the mean section, Gender can be categorized as majority male if the mean > 0.5, and majority female if \leq 0.5. Age can be divided into young (<30), adult (30-50), and elderly (>50) categories. Address was classified as city-dominant if the mean was close to 0, and county-dominant if higher. History of Drug Use showed new patients if the mean \leq 1, and patients with a history if more than 1. History of Resistance and Therapy Administered could be categorized as the majority did not experience resistance/therapy if the mean ≈ 0 , and some experienced if the mean > 0. Side Effects and Comorbidities showed that if the mean = 1, all patients experienced it, while if the mean = 0,

Table 2

Description of Analysis Cluster

patients did not experience it. Meanwhile, Survival is categorized as low if the mean is close to 0 if many survive and 1 if many do not survive (see Table 2).

Cluster	Variable	Count	Mean	Std	Min	25%	50%	75%	Max
0	Gender	219	0.635	0.483	0	0	1	1	1
	Age	219	45.37	18.31	9	28	50	58	79
	Address	219	0.493	0.875	0	0	0	1	4
	Impression	219	0.0	0.0	0	0	0	0	0
	Grade Based on Drug Use History	219	1.566	0.834	0	1	1	2	4
	Resistance History	219	0.064	0.245	0	0	0	0	1
	Therapy Administered	219	0.064	0.245	0	0	0	0	1
	Side Effects	219	1.0	0.0	1	1	1	1	1
	Comorbidities	219	0.0	0.0	0	0	0	0	0
	Survival	219	0.041	0.199	0	0	0	0	1
1	Gender	150	0.620	0.487	0	0	1	1	1
	Age	150	52.21	13.02	18	44.25	53.5	62	78
	Address	150	0.773	1.216	0	0	0	2	5
	Impression	150	0.0	0.0	0	0	0	0	0
	Grade Based on Drug	150	1.533	0.849	0	1	1	2	4
	Resistance History	150	0.073	0.262	0	0	0	0	1
	Therapy Administered	150	0.053	0.225	0	0	0	0	1
	Side Effects	150	1.0	0.0	1	1	1	1	1
	Comorbidities	150	1.0	0.0	1	1	1	1	1
	Survival	150	0.047	0.212	0	0	0	0	1
2	Gender	98	0.622	0.487	0	0	1	1	1
	Age	98	48.58	16.40	0	38.25	52	61	78
	Address	98	0.653	1.066	0	0	0	1.75	5
	Impression	98	0.0	0.0	0	0	0	0	0
	Grade Based on Drug Use History	98	1.520	0.911	0	1	1	2	4
	Resistance History	98	0.020	0.142	0	0	0	0	1
	Therapy Administered	98	0.020	0.142	0	0	0	0	1
	Side Effects	98	0.0	0.0	0	0	0	0	0
	Comorbidities	98	0.429	0.497	0	0	0	1	1
	Survival	98	0.122	0.329	0	0	0	0	1

DISCUSSION

Tuberculosis is a life-threatening infectious disease caused by Mycobacterium tuberculosis, and

timely diagnosis and effective treatment are essential for its control (13). Treatment of tuberculosis requires adjustment in special circumstances, such as Crohn's disease and other circumstances (14). Grouping patients into subgroups can guide prioritization and determine the provision of intensive care(15). From the research conducted in the hospital, 467 patient data were obtained from the patient's medical record of 467 and processed using the K-Means algorithm.

In Cluster 0, which was dominated by younger patients (mean <45 years, including adolescents) without comorbidities but experiencing adverse effects of therapy, tuberculosis infection in children and adolescents is influenced by factors such as characteristics. contact traits. case and environmental determinants, and TB preventive treatment is recommended to reduce the risk of disease progression (16). Individuals of productive age (ages 15-49) are particularly susceptible to tuberculosis, with 96% affected by the disease, and high school education is associated with higher risk (17). Children most commonly acquire TB infection after exposure to adults with infectious TB disease, with a high risk of disease progression and death (18). Children can contract TB due to poor implementation of preventive therapy in those with latent infection, contacts, and people living with HIV and AIDS in most developing countries (19). Those with immunodeficiency have a higher risk of progression of TB infection to TB disease, and concurrent infections with viruses, bacteria, fungi, and protozoa can influence this risk (20). Prevention can focus on optimizing drug side effect monitoring through intensive patient education. Information on the early signs of serious side effects needs to be clearly communicated, followed by a rapid reporting system to healthcare professionals. In addition, early screening to adjust drug doses based on the patient's weight and clinical status can help reduce side effects.

In Cluster 1, patients tend to be older (mean age >52 years) with a high prevalence of comorbidities and continue to experience adverse effects of Elderly patients with pulmonary therapy. tuberculosis are vulnerable, with high mortality and challenging diagnosis, and tertiary referral hospitals with expertise in TB management can improve treatment outcomes (21). Elderly people are more susceptible to TB due to factors such as decreased lung function, premature aging, inflammation, drug side effects, low tolerance to anti-TB drugs, and age-related comorbidities (22). Elderly patients with pulmonary tuberculosis experience increased mortality during treatment and more frequent adverse effects, even when pyrazinamide was often avoided (23). This is compounded by the presence of comorbidities such as diabetes, hypertension, and HIV in patients. Comorbidities, especially diabetes and chronic respiratory diseases, may delay

treatment response and require continued treatment for a successful outcome in TB patients (24). In addition, multimorbidity is common among TB patients (25). Early detection of comorbidities of diabetes, hypertension, and HIV in TB patients is important, as the prevalence of these conditions is high and their associated risk factors are significant. TB patients with diabetes mellitus or depression have a higher risk of death, relapse, and recurrence, and their comorbidities may negatively accrue in a synergistic manner (26). In addition, high blood pressure in TB patients leads to increased mortality. People with TB and HIV need to be vigilant because HIV increases the risk of TB infection and disease progression, and TB slows CD4 recovery and increases progression to AIDS and death among the HIV-infected (27). Preventive measures include comprehensive comorbidity management, such as control of chronic diseases (hypertension or diabetes) to reduce complications. Drug therapy should be selected with consideration of drug interactions, and a multidisciplinary approach involving internal medicine specialists or clinical pharmacologists is required. Patient education regarding the importance of medication adherence also needs to be strengthened to prevent drug resistance.

In Cluster 2, which includes patients with a mean age of 48 years with moderate comorbidities and low risk of drug resistance, prevention can focus on early detection of comorbidities through routine screening programs. Tuberculosis screening can improve diagnosis, treatment outcomes (28). Pharmacist intervention in screening tests for comorbidities in at-risk patients allows early detection and prompt treatment, preventing serious health consequences (29). Education on healthy lifestyles, such as a balanced diet, exercise, and cessation of bad habits such as smoking, can reduce the burden of future comorbidities. In addition, supervision of appropriate drug use should be strengthened to prevent the emergence of resistance in the future. Multilevel interventions are needed to address the unique challenges of TB treatment adherence among children and adolescents/young adults, which are influenced by structural, community, health system, household, and individual factors (30).

Based on the analysis of the three patient clusters, the recommendations for health workers, especially pharmacists, involve a specific approach for each group. In Cluster 0, focus should be on intensive education on monitoring drug side effects and early detection of serious signs to prevent complications. In Cluster 1, priority is given to comorbidity management through multidisciplinary collaboration, such as diabetes, hypertension, and HIV control, and therapy selection that considers drug interactions. For Cluster 2, prevention through routine screening programs and healthy lifestyle education needs to be enhanced to reduce the risk of future comorbidities and resistance. This approach aims to improve patients' quality of life and overall therapeutic success.

Intervention strategies obtained can be carried out with an approach to each cluster. In Cluster 1, which includes older patients with high comorbidities, chronic disease management is a priority with regular comorbidity checks by doctors, close monitoring of treatment by nurses, and pharmacist consultations on potential drug interactions. Meanwhile, Cluster 2, which has the highest mortality rate and moderate comorbidity levels, requires an intensive risk-based approach, such as closer supervision of therapy by doctors, monitoring of side effects and clinical conditions by nurses, and special education by pharmacists to ensure patients understand the importance of adherence. This multidisciplinary collaboration can improve the effectiveness of treatment and optimize care outcomes in each cluster.

In the future, researchers may look into ways to lower the side effects of TB drugs in young patients using pharmacogenetic approaches and non-drug interventions. They may also look into how to manage TB and other health problems in older patients. In addition, exploration of socioeconomic influences, digital monitoring technologies, and predictive biomarkers may provide new insights to improve the success of TB therapy in a holistic manner.

CONCLUSION

K-Means clustering analysis shows that TB patients with comorbidities in Gorontalo city hospital have diverse characteristics, requiring a customized approach for each group. Cluster 0 requires attention to side effect monitoring and intensive education, Cluster requires 1 comprehensive comorbidity management and strengthening of treatment adherence, while Cluster 2 focuses on prevention through routine screening and promotion of healthy lifestyles. These recommendations highlight the importance of a multidisciplinary role, including pharmacists, in supporting more effective therapy and improving TB patient outcomes.

CONFLICT OF INTEREST

All authors declare there is no conflict of interest.

AUTHOR CONTRIBUTIONS

The contributions of each author are as follows: MHW was responsible for conceptualization, methodology, software, and writing; SNA contributed to data curation and writing; AFH handled visualization; ED focused on reviewing and editing; WS contributed to reviewing; and SS was involved in reviewing and editing.

ACKNOWLEDGMENTS

This study was supported by the Directorate of Research, Technology, and Community Service, Ministry of Education, Culture, Research, and Technology, Indonesia, under the grant No. 107/ES/PG.02.00.PL/2024, 069.12/LL5.INT/AL.04/2024,115 / PTM / LPPM-UAD/VI/2024, 15 June 2024. The author also expresses his gratitude to the Prof. Dr. Aloei Saboe Hospital in Gorontalo City for its assistance in the process of this research support.

REFERENCES

- Furin J, Cox H, Pai M. Tuberculosis. Vol. 393, The Lancet. Lancet Publishing Group; 2019. p. 1642–56.
- 2. Falzon D, Zignol M, Bastard M, Floyd K, Kasaeva T. The impact of the COVID-19 pandemic on the global tuberculosis epidemic. Vol. 14, Frontiers in Immunology. Frontiers Media SA; 2023.
- 3. Amanda Perkins. Tuberculosis [Internet]. Available from: www.NursingMadeIncrediblyEasy.com
- 4. Nadjamuddin M, Junita N. TEMUKAN TUBERCULOSIS OBATI SAMPAI SEMBUH. Vol. 1. 2023.
- 5. Nurany H, Raharjo M, Sakundarno Adi M. Environmental Quality Factors with The Incidence of Pulmonary Tuberculosis: A Literature Review. Serambi Engineering. 2022;VII(3).
- H, Tosepu R, Savitri Effendy D. Tuberculosis Epidemiology and Medical Treatment Efforts in Indonesia in the Year 2020. KnE Life Sciences. 2022 Sep 13;
- 7. Indri Yosua M, Ningsih F, Ovany R, Eka Harap Palangka Raya Stik, Raya P, Tengah

K. Relationship With House Environmental Conditions Event Of Tuberculosis (TB) Lungs. 2022 [Internet]. Available from: http://journal.umpalangkaraya.ac.id/index.p hp/jsm

- 8. Dinkes Provinsi Gorontalo. Profil Kesehatan Provinsi Gorontalo 2022. 2023.
- Pakaya R. Factors related to Pulmonary Tuberculosis Incident in Public Health Center of Limboto Year 2018. KESMAS UWIGAMA: Jurnal Kesehatan Masyarakat. 2020 Jul 9;6(1):1–13.
- Wang Y, Zheng D. The importance of precision medicine in modern molecular oncology. Vol. 100, Clinical Genetics. John Wiley and Sons Inc; 2021. p. 248–57.
- Denny JC, Collins FS. Precision medicine in 2030—seven ways to transform healthcare. Cell. 2021 Mar 18;184(6):1415–9.
- Choi D, Shrestha D, Xiang A, Drake B, Ozturk O, Haidarian H, et al. Patient Clustering via Integrated Profiling of Clinical and Digital Data. In: International Conference on Information and Knowledge Management, Proceedings. Association for Computing Machinery; 2023. p. 3818–22.
- Dong B, He Z, Li Y, Xu X, Wang C, Zeng J. Improved Conventional and New Approaches in the Diagnosis of Tuberculosis. Vol. 13, Frontiers in Microbiology. Frontiers Media S.A.; 2022.
- Kafle MP. Treating tuberculosis in special situations. Bangladesh Journal of Medicine. 2024 Jun 6;141–2.
- Li X, van Giessen A, Altunkaya J, Slieker RC, Beulens JWJ, 'T Hart LM, et al. Potential Value of Identifying Type 2 Diabetes Subgroups for Guiding Intensive Treatment: A Comparison of Novel Data-Driven Clustering With Risk-Driven Subgroups. Diabetes Care. 2023 Jul 1;46(7):1395–403.
- 16. Tchakounte Youngui B, Tchounga BK, Graham SM, Bonnet M. Tuberculosis Infection in Children and Adolescents. Vol. 11, Pathogens. MDPI; 2022.
- Marlina L, Darmansyah D. Descriptive Study of Certainty Level Towards Risk of Tuberculosis Disease in Productive Ages. KnE Life Sciences. 2021 Mar 15;1063–8.
- Togun T. Childhood tuberculosis in high burden settings. Vol. 63, EBioMedicine. Elsevier B.V.; 2021.
- 19. Mustapha MG, Ashir GM, Rabasa AI, Farouk AG, Elechi HA, Alhaji MA.

Childhood tuberculosis: Characteristics and peculiarities. Niger J Paediatr. 2020 Aug 6;47(3):190–200.

- Whittaker E, Lopez-Varela E, Broderick C, Seddon JA. Examining the complex relationship between tuberculosis and other infectious diseases in children: A review. Vol. 7, Frontiers in Pediatrics. Frontiers Media S.A.; 2019.
- Di Gennaro F, Vittozzi P, Gualano G, Musso M, Mosti S, Mencarini P, et al. Active pulmonary tuberculosis in elderly patients: A 2016–2019 retrospective analysis from an Italian referral hospital. Antibiotics. 2020 Aug 1;9(8):1–10.
- 22. Olmo-Fontánez AM, Turner J. Tuberculosis in an Aging World. Vol. 11, Pathogens. MDPI; 2022.
- Hase I, Toren KG, Hirano H, Sakurai K, Horne DJ, Saito T, et al. Pulmonary Tuberculosis in Older Adults: Increased Mortality Related to Tuberculosis Within Two Months of Treatment Initiation. Drugs Aging. 2021 Sep 1;38(9):807–15.
- Kunoor A, Reddy S, Gopalakrishnan V, Rakesh P, Kadayara A, Haridas N, et al. Burden of comorbidity and treatment outcome in tuberculosis – A descriptive study from a tertiary care center, Kerala, India. PULMON. 2023;25(2):44.
- 25. Chen Q, Che Y, Xiao Y, Jiang F, Chen Y, Zhou J, et al. Impact of Multimorbidity Subgroups on the Health Care Use and Clinical Outcomes of Patients With Tuberculosis: A Population-Based Cohort Analysis. Front Public Health. 2021 Oct 8:9.
- 26. Cáceres G, Calderon R, Ugarte-Gil C. Tuberculosis and comorbidities: treatment challenges in patients with comorbid diabetes mellitus and depression. Vol. 9, Therapeutic Advances in Infectious Disease. SAGE Publications Ltd; 2022.
- Vázquez García JC, Sada Díaz E, Rivera Martínez E, Narváez Porras O, Salazar Lezama MA. Tuberculosis associated with HIV infection. Rev Invest Clin. 1994;46(6):473–7.
- Telisinghe L, Ruperez M, Amofa-Sekyi M, Mwenge L, Mainga T, Kumar R, et al. Does tuberculosis screening improve individual outcomes? A systematic review. EClinicalMedicine. 2021 Oct 1;40.
- 29. Lewicki J, Religioni U, Merks P. Evaluation of the community pharmacy comorbidities screening service on patients with chronic

diseases. Patient Prefer Adherence. 2021;15:1611–9.

30. Leddy AM, Jaganath D, Triasih R, Wobudeya E, Bellotti De Oliveira MC, Sheremeta Y, et al. Social Determinants of Adherence to Treatment for Tuberculosis Infection and Disease Among Children, Adolescents, and Young Adults: A Narrative Review. J Pediatric Infect Dis Soc. 2022 Oct 1;11:S79–84.