RURAL-URBAN DISPARITIES IN DETERMINANTS OF BCG IMMUNIZATION COVERAGE FOR CHILDREN AGED 0-3 MONTHS IN INDONESIA

Nur Mutiara Husnah Hidayatullah BW¹, *Erni Astutik¹, Arief Hargono¹ ¹Faculty of Public Health, Universitas Airlangga, 60115 Surabaya, East Java, Indonesia *Corresponding Author: Erni Astutik ; Email: <u>erniastutik@fkm.unair.ac.id</u>

ABSTRACT

Bacillus Calmette-Guérin (BCG) immunization plays a crucial role in preventing tuberculosis, especially in developing countries like Indonesia. Its coverage is influenced by various factors that differ between urban and rural areas. This study aimed to identify factors associated with BCG immunization among infants aged 0-3 months in urban and rural Indonesia. This study used data from the 2017 Indonesia Demographic and Health Survey (IDHS), including 562 rural and 476 urban respondents. Variables analyzed included maternal age, education, household wealth, contraceptive use, and ANC visits. Chi-square tests and multiple logistic regression were conducted separately for urban and rural areas. The findings showed that 64.2% of urban and 48.7% of rural children received BCG immunization. In rural areas, children from the richest households were more likely to be immunized compared to those from the poorest households (OR = 2.29; 95%CI = 1.1–4.6). Children of mothers with higher education in urban areas were more likely to receive BCG compared to those with lower education (OR = 3.43; 95%CI = 1.3-9.0), as were those with secondary education (OR = 2.28; 95%CI = 1.1-4.4). Mothers who used contraception were also more likely to immunize their children compared to those who did not (OR = 2.71; 95%CI = 1.5-4.8). BCG coverage remains lower in rural communities. Improving coverage requires expanding access to health services in rural areas and strengthening education and contraceptive programs in urban settings.

Keywords: BCG immunization, children, disparities, Indonesian Demographic Health Survey, rural-urban

ABSTRAK

Imunisasi Bacillus Calmette–Guérin (BCG) berperan penting dalam mencegah tuberkulosis, terutama di negara berkembang seperti Indonesia. Cakupannya dipengaruhi oleh berbagai faktor yang berbeda antara daerah perkotaan dan perdesaan. Studi ini bertujuan mengidentifikasi faktor yang berhubungan dengan imunisasi BCG pada bayi usia 0-3 bulan di daerah perkotaan dan perdesaan di Indonesia. Studi ini menggunakan data dari Survei Demografi dan Kesehatan Indonesia (SDKI) 2017, dengan 562 responden dari perdesaan dan 476 dari perkotaan. Variabel yang dianalisis meliputi usia ibu, pendidikan, kekayaan rumah tangga, penggunaan kontrasepsi, dan kunjungan ANC. Uji chi-square dan multiple logistic regression dilakukan secara terpisah untuk daerah perkotaan dan perdesaan. Hasil menunjukkan 64,2% anak di perkotaan dan 48,7% anak di perdesaan menerima imunisasi BCG. Di perdesaan, anak dari rumah tangga terkaya lebih mungkin diimunisasi dibandingkan anak dari rumah tangga termiskin (OR = 2,29; 95%CI = 1,1-4,6). Di perkotaan, anak dari ibu dengan pendidikan tinggi lebih mungkin menerima BCG dibandingkan yang berpendidikan rendah (OR = 3,43; 95%CI = 1,3-9,0), demikian juga dengan pendidikan menengah (OR =2,28; 95%CI = 1,1-4,4). Ibu yang menggunakan kontrasepsi juga lebih mungkin mengimunisasi anaknya dibandingkan yang tidak (OR = 2,71; 95%CI = 1,5-4,8). Cakupan BCG tetap lebih rendah di komunitas perdesaan. Peningkatan cakupan memerlukan perluasan akses layanan kesehatan di perdesaan dan penguatan program pendidikan serta kontrasepsi di perkotaan.

Kata kunci: Imunisasi BCG, anak-anak, kesenjangan, Survei Demografi dan Kesehatan Indonesia, perdesaan-perkotaan

INTRODUCTION

(TB), Tuberculosis caused bv various species of Mycobacteria, persists as a major threat to health on a global scale, despite being both preventable and treatable. With a staggering 1.5 million deaths annually, TB continues to be the most lethal infectious illness globally (1). Recognizing the urgency, World Health Organization (WHO) and United Nations (UN) member states launched the End TB Strategy alongside the the Sustainable Development Goals (SDGs) in 2014-2015, aiming to combat the epidemic. The plan aims for significant reductions in tuberculosis incidence, targeting a decrease of 4-5% annually by 2020, 10% by 2025, and 17% by 2035 (2).

Approximately 10 million individuals across the globe were affected by tuberculosis during 2020, with 5.6 million men, 3.3 million women, and 1.1 million children. The disease transcends geographic boundaries and affects all age demographics (3). A total of 87% of all estimated global cases came from the 30 countries with the highest TB rates, with eight nations alone contributing to over two-thirds of the worldwide total. Within these countries, Indonesia ranks as the second-largest contributor, responsible for 10% of the global cases (2).

The projected occurrence rate of tuberculosis in Indonesia in 2021 is 969,000 or 354 per 100,000 population, while the death rate from TB is estimated at 144,000 deaths (4). The Indonesian government has launched a comprehensive national program for tuberculosis control that encompasses a series of strategic measures, including health promotion, TB surveillance, risk factor management, case detection and treatment, immunization, and preventive medication (5). Included in these measures, the BCG vaccine is a mandatory immunization that has shown high success rates, especially in children aged 12-23 months (6).

The only approved vaccine for preventing tuberculosis is Bacillus Calmette-Guérin (BCG), which provides protection against various forms of the disease such as pulmonary, meningeal and disseminated tuberculosis (7)(8). Across high incidence of countries with а tuberculosis, the BCG vaccine is strongly recommended for children and serves as a strategy primary in TΒ prevention. Therefore. BCG immunization is administered to infants as part of the national tuberculosis control program (5). BCG immunization rates among children aged 12–23 months were reported to be high in the 2017 IDHS data—92.9% in urban areas and 89.3% in rural ones (9). However, these aggregate figures may mask inequalities in early coverage, particularly among infants in their first three months of life when BCG should ideally be administered.

Several studies have identified key factors influencing childhood vaccination, including maternal age (10), education (11), household wealth (12), antenatal care utilization (13), contraceptive use (14), and place of residence (15)(10). While research in Indonesia has explored the determinants of BCG immunization (14), few studies in Indonesia have explored how these factors differ between urban and rural areas, especially for infants aged 0–3 months. This gap is particularly concerning given Indonesia's geographic and socioeconomic diversity, which may impact access to timely immunization services.

A study on immunization determinants in Indonesia emphasized that geographic location significantly affects immunization coverage, with place of residence identified as one of the key predictors (16). Regarding the Indonesian context, urban areas often benefit from better infrastructure, shorter distances to health facilities, and more healthcare providers, advantages that may not be available in rural settings.

Building on this understanding of geographic disparities, this study aims to examine the disparities BCG in immunization coverage among infants aged 0-3 months in urban and rural areas of Indonesia, and it also explores the role of maternal factors such as age, education level, household wealth, antenatal care utilization. and contraceptive use as potential contributors to immunization inequalities in both settings. Maternal factors are a central focus because mothers are typically the primary decision-makers in child healthcare. Moreover, previous evidence indicates that maternal characteristics play a significant role in supporting children's access to immunization services (17).

METHODS

Data Sources

This study used secondary data from the 2017 IDHS.

Population and Sample

This research collected data from the 2017 IDHS, which included a total of 17,848 mothers aged 15 to 49 who were successfully surveyed. The sampling process was carried out in two stages: 1) choosing groups from every stratum and creating a directory of households located in the selected groups, and 2) subsequently selecting households for interviews (18). This study concentrated on children aged 0 to 3 months, leading to a total of 1,073 mothers with children in that age range. Within this group, 580 mothers were from rural areas, while 493 were from urban areas. The analysis was performed after filtering the data to include only complete observations, resulting 1.038 in

participants, with 562 residing in rural areas and 476 in urban areas. The population and sample are described as follows:



Figure 1. Flowchart of Sample Selection

Research Variables`

This study's dependent variable focused on BCG immunization, categorized into two groups: not received and received. A child was considered vaccinated if the vaccination date was noted on the card and confirmed by the mother. Conversely, if there was no such record, the child was classified as not having received BCG immunization.

The independent variables examined included maternal age, maternal education level, wealth index, contraceptive use, and antenatal care (ANC). Maternal age was categorized into ≤ 24 , 25 - 34, and ≥ 35 vears. Maternal education was classified into three tiers: low (no formal schooling or completion of only primary education), medium (junior high school), and high (senior high school through college). The wealth index was categorized into poorest, poorer, average, richer, and richest. For contraceptive use, we distinguished between those who used contraceptives and those who did not. ANC visits were classified as incomplete if they were less than three and as complete if they were four or more.

The choice of these variables was based on Andersen's Behavioral Model of Health Services Use (19), a framework frequently used to identify factors affecting healthcare utilization, including immunization (20)(21). Based on this model, grouped the variables were into predisposing, enabling, and environmental components. Maternal age and education categorized predisposing were as characteristics, wealth index. while and ANC contraceptive visits use. represented enabling resources. The place of residence served an external as environmental factor, aligning with the study's focus on rural-urban disparities. Recognizing these contextual drivers is essential to inform policies and design equitable immunization programs that meet the needs of both rural and urban communities in Indonesia.

Data Analysis

The data analysis followed a threestage approach. Initially, descriptive statistics were employed outline to respondent characteristics. Next, chi-square tests were performed to analyze the association between individual predictors and the dependent variable. Variables demonstrating a p value below 0.25 in the bivariate analysis were subsequently entered into a multivariable logistic regression model to determine the factors significantly associated with the outcome. All statistical procedures incorporated the complex sampling structure of the IDHS and were conducted separately for urban and rural populations.

Research Ethics

The IDHS received ethical approval from both the ICF Institutional Review Board (IRB) and the IRB in the host country (22).

RESULT

The analysis included 1,038 respondents, with 562 residing in rural areas and 476 in urban areas. A marked difference in BCG immunization coverage was observed, with 64.2% of children aged 0–3 months vaccinated in urban areas, compared to 48.7% in rural areas. Conversely, a larger proportion of rural children (51.2%) had not received BCG compared to urban children (35.7%).

Most mothers in both areas were aged 25–34 years, followed by those aged \leq 24 years. The proportion of mothers aged 35 and older was the smallest in both settings. Urban mothers had high levels of education, with 61.7% having medium education compared to 50.5% in rural areas. Low education was more prevalent in rural areas (32.2%) than in urban (15.3%).

Rural respondents were more likely to fall into the poorest wealth category (30.7%), while the richest category had proportions in similar both areas. Contraceptive use was slightly higher among urban mothers (74.1%) than rural (72.6%). ANC visit completion was generally high, but slightly greater in urban (89.0%) than rural areas (82.5%), while incomplete ANC visits were more common in rural communities (17.4% vs. 10.9%).

Table 2 shows that in rural areas, BCG immunization in children aged 0-3 months was associated with wealth index, contraceptive use, and ANC visits with BCG immunization in children aged 0-3 months. Meanwhile, in urban areas, maternal age, maternal education, wealth index, contraceptive use, and ANC were associated with BCG immunization in children 0-3 months of age.

Table 3 shows that in rural areas, household wealth had a significant influence on BCG vaccination. Children from wealthier families were 2.29 times more likely to receive the vaccine compared to those from the poorest families (OR=2.29; 95% CI: 1.1–4.6). Meanwhile in urban areas, maternal education and contraceptive use were important predictors. Mothers with high education were 3.43 times more likely to vaccinate their children compared to those with low education (OR = 3.43; 95% CI: 1.3–9.0), while mothers with medium education were 2.28 times more likely (OR = 2.28; 95% CI: 1.1–4.4). Additionally, mothers who used contraceptives were 2.71 times more likely to vaccinate their children compared to non-users (OR=2.71; 95% CI: 1.5-4.8).

DISCUSSION

BCG immunization uptake among children aged 0-3 months was higher in urban than in rural areas. A previous study supports this finding, indicating that immunization coverage, including BCG, tends to be higher in urban areas in Indonesia (23). Access to healthcare services in rural regions is often constrained by a shortage of

Table 1. Description of Study Participants

adequately trained medical personnel and limited transportation infrastructure. Several studies have identified distance to health facilities as a key barrier to completing childhood immunization, particularly in Indonesia, an archipelago comprising more than 17,000 islands (24)(25). Other research conducted in developing nations has similarly indicated that immunization rates are more significant in urban area than in rural regions (10)(26). Prior investigations demonstrated that geographical have proximity significantly influences mothers' ability to access healthcare services (27). The higher immunization rates in urban areas result from improved access to healthcare facilities offering the Essential Immunization Program and variations in vaccination awareness and health-seeking behaviors (26).

Variables		Rural (N = 562)	Urban (N = 476)	Total (N=1038)	
		N (%)	N (%)	N (%)	
BCG immunization status	Not received	288 (51.2)	170 (35.7)	458 (44.1)	
	Received	274 (48.7)	306 (64.2)	580 (55.9)	
Mother's Age	≤ 24	160 (28.4)	138 (28.9)	398 (28.7)	
	25 - 34	300 (53.3)	236 (49.5)	536 (51.6)	
	≥ 35	102 (18.1)	102 (21.4)	204 (19.7)	
Mother's Education	Low	181 (32.2)	73 (15.3)	254 (24.5)	
	Medium	284 (50.5)	294 (61.7)	578 (55.7)	
	High	97 (17.2)	109 (22.9)	206 (19.8)	
Wealth Index	Poorest	173 (30.7)	132 (27.7)	305 (29.4)	
	Poorer	106 (18.8)	91 (19.1)	197 (19.0)	
	Average	94 (16.7)	105 (22.0)	199 (19.2)	
	Richer	91 (16.1)	74 (15.5)	165 (15.9)	
	Richest	98 (17.4)	74 (15.5)	1/2 (16.6)	
Contraceptive Use	No	154 (27.4)	123 (25.8)	277 (26.7)	
	Yes	408 (72.6)	353 (74.1)	761 (73.3)	
ANC Visits	Incomplete	98 (17.4)	52 (10.9)	150 (14.5)	
	Complete	464 (82.5)	424 (89.0)	888 (85.5)	

Source: IDHS 2017

Conversely, individuals in rural areas often face significant challenges in accessing immunization services. with limited transportation options and long distances to health facilities being major barriersparticularly in Indonesia, an archipelagic nation consisting of over 17,000 islands (24)(25). The geographical complexity of the country makes it difficult for rural populations to reach health centers, thereby hindering timely vaccination. This finding is reinforced by evidence that distance to health facilities remains one of the key obstacles to completing childhood immunization. However, this contrasts with findings from other studies (28), which revealed that rural communities achieved higher immunization rates than their urban counterparts, although it still remained far below the WHO-recommended target.

Table 2. Distribution of Maternal Socio-Demographics, Contraceptive Use, and ANC Visits Based on The BCG Immunization Status of Children Aged 0-3 Months in Rural and Urban Areas in Indonesia

		Rural			Urban							
	BCG			BCG								
	ImmunizationVariablesNoYes		χ^2 p		Immunization					p		
Variables			Yes			value	No		Yes		χ^2	value
	Ν	%	Ν	%	I		N	%	N	%		
Mother's Age												
≤ 24	87	54.3	73	45.6			58	42.0	80	57.9		
25 - 34	144	48.0	156	52.0	2.8	0.42	73	30.9	163	69.0	13.0	< 0.001
\geq 35	57	55.8	45	44.1			39	38.2	63	61.7		
Mother's Educa	tion											
Low	100	55.2	81	44.7			33	45.2	40	54.7		
Medium	141	49.6	143	50.3	2.5	0.40	106	36.0	188	63.9	12.1	0.02
High	47	48.4	50	51.5			31	28.4	78	71.5		
Wealth Index												
Poorest	105	60.6	68	39.3			57	43.1	75	56.8		
Poorer	58	54.7	48	45.2			30	32.9	61	67.0		
Average	46	48.9	48	51.6	15.2	0.04	34	32.3	-71	67.6	8.6	0.21
Richer	33	36.2	58	63.7			22	29.7	52	70.2		
Richest	46	46.9	52	53.6			27	36.4	47	63.5		
Contraceptive U	Jse											
No	94	61.4	60	38.9	2.8	0.18	59	47.9	64	52.0	16.9	< 0.001
Yes	194	47.5	214	52.4			111	31.4	242	68.5		
ANC Visits												
Incomplete	64	65.3	34	34.6	00	<0.001	23	44.2	29	55.7	56	0.03
Complete	224	48.2	240	51.7	9.0	<0.001	147	34.6	277	65.3	5.0	0.05
Source: IDHS 201	7											

Source: IDHS 2017

BCG immunization is more likely among children from wealthier families in rural areas. Previous studies have shown that as household wealth rises, the likelihood of children being unimmunized decreases, with children from the poorest households in Indonesia significantly facing a 2.9 times higher risk of not receiving vaccinations (21). This research further supports findings from other studies indicating that children born to families with a higher wealth index are more inclined to get vaccinated (12)(29)(30). Research from Nigeria also highlights persistent health disparities, with children from wealthier households more likely to receive full immunization. These inequalities are considered unfair yet preventable, and call for policies tailored to local socio-cultural contexts to reduce gaps in coverage (31). This study supports their view that household wealth plays a role in immunization, and public health efforts should address economic barriers, especially in rural areas. Families with greater economic means often have improved access to immunization information and resources, which enhances their capacity to fulfill the recommended childhood vaccinations.

Although essential vaccinations are offered at no cost in Indonesia, wealthier families tend to be more proactive in seeking and obtaining vaccinations due to better access to health-related information and services. Other countries, such as Ghana, demonstrate that even when health insurance and services are provided for free, women often still face out-of-pocket expenses for transportation and additional services costs that tend to favor wealthier mothers (32)(33). As a result, economically disadvantaged women, despite recognizing the importance of immunization, often face financial barriers (29). This study opposes previous findings of no association between wealth and BCG coverage in 0–2-month-old infants (14).

Table 3. Relationship Between Maternal Socio-Demographics, Contraceptive Use, and ANCVisits with BCG Immunization among Children Aged 0-3 Months in Rural and Urban Areas inIndonesia

	Rural		Urban		
Variables	OR	95% CI	OR	95% CI	
Mother's Age					
\leq 24	Ref		Ref		
25 - 34	1.26	0.75 - 2.12	1.60	0.91 - 2.79	
≥ 35	0.92	0.49 - 1.74	0.87	0.44 - 1.72	
Mother's Education					
Low	Ref		Ref		
Medium	1.05	0.62 - 1.76	2.28**	1.17 - 4.44	
High	1.22	0.58 - 2.53	3.43**	1.30 - 9.03	
Wealth Index					
Poorest	Ref		Ref		
Poorer	1.04	0.55 - 1.95	1.15	0.59 - 2.23	
Average	1.27	0.68 - 2.36	1.41	0.71 - 2.77	
Richer	2.29**	1.14 - 4.61	1.34	0.61 - 2.94	
Richest	1.52	0.76 - 3.04	0.83	0.34 - 2.04	
Contraceptive Use					
No	Ref		Ref		
Yes	1.34	0.81 - 2.20	2.71*	1.51 - 4.85	
ANC Visits					
Incomplete	Ref		Ref		
Complete	1.64	0.93 - 2.86	1.41	0.65 - 3.02	
Source : IDHS 2017					

Note : * = p < 0.01; ** = p < 0.05

Maternal education and contraceptive use are key factors influencing BCG immunization in urban areas, unlike rural regions. This research indicates that more educated mothers are more likely to vaccinate their children with BCG. The category of 'high education' referred to in this study encompasses high school to university levels. These results are in line with a study conducted in urban India (34)

and are further supported by research in various developing countries, including Afghanistan (35), Spain (36), Laos (37), Pakistan (38), and several others such as Nepal, Senegal, and Zambia (39). A systematic review and meta-analysis also revealed regional variations, with the impact of maternal education on immunization uptake being stronger in Asia and Africa compared to Europe, likely because education in low-income countries is more closely linked to access to healthcare services (40). Additionally, mothers with higher education tend to have better and awareness capability to access immunization services (41)(42)(43). They are also more effective in reducing missed opportunities for vaccination (MOV) (44). In general, educated mothers are expected to have greater access to health information, which promotes positive health-seeking behaviors and contributes to higher childhood vaccination rates (45).

This research also revealed that urban mothers who utilized contraceptives were more likely to vaccinate their children aged 0-3 months with BCG. A prior study in Indonesia reported similar findings, associating contraception with BCG immunization in infants under 3 months. An increase in the number of children in a household is associated with divided parental attention, making it more challenging to meet each child's needs and decreasing the likelihood of timelv vaccination (14). Parents who use contraception tend to have fewer children. allowing them to provide more attention to each child. Conversely, families with a larger number of children frequently face financial challenges, which diminishes the likelihood of vaccinating the youngest child (46)(47). Additionally, phenomenological research examining barriers to basic immunization coverage in children suggests that having closely spaced siblings may hinder mothers from accessing immunization services for their children (48). The adoption of contraceptive methods may facilitate improved family

planning, enabling parents to focus more on the health and immunization of each child.

The disparities in factors influencing BCG immunization between rural and urban areas highlight the need for context-specific strategies in immunization program design. Rural areas face economic constraints as а dominant barrier. particularly among low-income families who often struggle with indirect costs such as transportation and opportunity costs. Addressing this requires policies that reduce financial burdens through targeted economic empowerment programs, such as conditional cash transfers or transportation subsidies, while simultaneously strengthening community-based education on the importance of immunization. These combined efforts can help increase both awareness and access, ultimately improving immunization coverage in underserved rural populations. In contrast, urban areas face different challenges. While physical access to health services is generally better, gaps persist in maternal knowledge and engagement with immunization services. Therefore, improving maternal education through health literacy programs is crucial. Such programs should emphasize not only the benefits of immunization but also promote contraceptive use as part of a broader child health strategy. As maternal education levels increase. so does contraceptive use (49), which can lead to better family planning and more focused attention on each child's health needs. Integrating immunization education into family planning and maternal health services can serve as a synergistic approach, equipping mothers with the knowledge and resources to ensure their children receive timely and complete vaccinations. Tailoring interventions to the unique barriers of each setting is essential to close the rural-urban gap in immunization coverage.

Limitations of the Study

This study is subject to several limitations that should be acknowledged when interpreting the findings. First, the analysis relies on secondary data derived from the 2017 IDHS, which, although methodologically rigorous, mav be considered outdated. Second, the scope of the study is confined to a limited set of variables, with a primary focus on maternal characteristics. This constraint precludes the examination of other potentially influential determinants, such as the availability and quality of healthcare infrastructure, cultural and religious beliefs surrounding immunization, paternal involvement, and broader household or community-level factors.

Nonetheless, a key strength of this research is its use of the IDHS dataset, which is nationally representative and follows standardized sampling and data collection methods. These features enhance the generalizability of the findings and support broader conclusions about immunization disparities across rural and urban populations.

CONCLUSION AND SUGGESTION

Conclusions

A disparity in BCG immunization coverage was observed, with infants aged 0–3 months in urban areas showing higher coverage than those in rural areas. The observed urban-rural disparity is associated with maternal characteristics, as identified within the scope of this research. In rural settings, higher household wealth increases the likelihood of immunization. Meanwhile, in urban areas, mothers with higher education levels and those who use contraceptives are more likely to have their children vaccinated.

Suggestions

Targeted policies and contextspecific intervention programs are needed to improve BCG immunization coverage. Economic empowerment programs in rural areas have the potential to enhance access to health services. Meanwhile, in urban settings, strengthening health literacy and maternal education—particularly through counseling on immunization and contraception—plays crucial а role. Collaboration between government and civil society organizations will be key to addressing access gaps. Further research on socioeconomic factors and health policies is also important to formulate more effective strategies.

ACKNOWLEDGEMENT

The Authors would like to thank the DHS Program for access to the data used in this study.

AUTHOR CONTRIBUTIONS

NMHH was responsible for study design, data collection, analysis, and manuscript drafting. EA and AH supervised the research process and provided critical input on data interpretation.

REFERENCES

- 1. WHO. Tuberculosis Geneva: World Health Organization [Internet]. 2020. Available from: https://www.who.int/healthtopics/tuberculosis#tab=tab 1.
- 2. WHO. Global Tuberculosis Report [Internet]. 2023. Available from: https://www.who.int/publications/i/i tem/9789240083851
- 3. WHO. Tuberculosis [Internet]. 2022. Available from: https://www.who.int/indonesia/news /campaign/tb-day-2022/fact-sheets
- 4. Ministry of Health. Laporan Program Penanggulangan Tuberkulosis Tahun 2022 [Report on the Tuberculosis Control Program 2022] Indonesia: [Internet]. 2023. Available from: https://www.tbindonesia.or.id/pusta ka---program-la/laporan-programpenanggulangan-tuberkulosis-tahun-2022/
- 5. Ministry of Health. Minister of

Health Regulation No. 67 on Tuberculosis Control, 2016 [Internet]. 2016. Available from: https://peraturan.bpk.go.id/Details/1 14486/permenkes-no-67-tahun-2016

- 6. Ministry of Women's Empowerment and Child Protection. Profil Anak Indonesia Tahun 2019 [Indonesian Child Profile 2019] [Internet]. 2019. Available from: https://www.kemenpppa.go.id/page/ view/MjUyMg==
- 7. Lange C, Aaby P, Behr MA, Donald PR, Stefan H E Kaufmann MGN, Mandalakas AM. 100 Years of Mycobacterium Bovis Bacille Calmette-Guérin. Lancet Infect Dis [Internet]. 2022;22(1). Available from: https://doi.org/10.1016/S1473-3099(21)00403-5
- Setiabudiawan TP, Reurink RK, Hill 8. PC, Netea MG, Crevel R van, Koeken VACM. Protection against tuberculosis by Bacillus Calmette-Guérin (BCG) vaccination: Α perspective. historical Med 2022;3(1):6-24. [Internet]. Available from: https://doi.org/10.1016/j.medj.2021. 11.006
- 9. Ministry of Health. Indonesia Demographic and Health Survey (IDHS) [Internet]. Jakarta; 2017. Available from: https://ia802800.us.archive.org/30/it ems/LaporanSDKI2017/Laporan SDKI 2017.pdf
- Tsehay, A. K., Worku, G. T., & Alemu YM. Determinants of BCG vaccination coverage in Ethiopia: A cross-sectional survey. BMJ Open [Internet]. 2019;9(2):1–7. Available from: https://doi.org/10.1136/bmjopen-2018-023634
- Ahmed Ali H, Hartner AM, Echeverria-Londono S, Roth J, Li X, Abbas K, et al. Vaccine equity in low and middle income countries: a

systematic review and meta-analysis. Int J Equity Health [Internet]. 2022;21(1):1–30. Available from: https://doi.org/10.1186/s12939-022-01678-5

- 12. Ijarotimi IT, Fatiregun AA, Adebiyi OA, Ilesanmi OS, Ajumobi O. differences Urban-rural in immunisation status and associated demographic factors among children 12-59 months in a southwestern state, Nigeria. Uthman O, editor. PLoS One [Internet]. 2018;13(11):1-11. Available from: https://doi.org/10.1371/journal.pone. 0206086
- 13. Ibrahim J AS. Survey on Coverage and Factors Influencing Delays in BCG Immunization in Hayin Mallam Zango, Zaria, North Western Nigeria. Trop Med Surg [Internet]. 2017;03(03):1–4. Available from: https://doi.org/10.4172/2329-9088.1000188
- 14. Kusnanto K, Arifin H, Kurniawati Y. Determinant of BCG vaccine coverage among Indonesian children aged 0–2 months. Child Youth Serv Rev [Internet]. 2020;116. Available from: https://doi.org/10.1016/j.childyouth. 2020.105238
- 15. Hu Y, Chen Y, Liang H, Wang Y. An overview of coverage of BCG vaccination and its determinants based on data from the coverage survey in Zhejiang Province. Int J Environ Res Public Health [Internet]. 2018;15(6). Available from: https://doi.org/10.3390/ijerph15061 155
- 16. Sinuraya RK, Alfian SD, Abdulah R, Postma Suwantika MJ. AA. Comprehensive Childhood Vaccination and Its Determinants: Insights From The Indonesia Family Life Survey (IFLS). J Infect Public Health [Internet]. 2024;17(3):509-Available 17. from: https://doi.org/10.1016/j.jiph.2024.0

1.007

- Demi L. 17. Analysis of Factors Influencing Mothers' Compliance in Providing Complete Basic Immunization in Kaledupa Subdistrict, Wakatobi Regency [Internet]. Universitas Hasanuddin; 2022. Available from: https://repository.unhas.ac.id/id/epri nt/18576/2/R011191082_skripsi_26 -08-2022 1-2.pdf
- Demographic Health Survey. Guide to DHS Statistics DHS-7 [Internet].
 2017. Available from: https://www.dhsprogram.com/pubs/ pdf/DHSG1/Guide_to_DHS_Statisti cs_DHS-7_v2.pdf
- Andersen RM. Revisiting the Behavioral Model and Access to Medical Care: Does it Matter? J Health Soc Behav [Internet]. 1995;36(1):1. Available from: https://pubmed.ncbi.nlm.nih.gov/77 38325/
- 20. Sarker AR, Akram R, Ali N, Chowdhury ZI, Sultana M. Coverage Determinants of Full and Immunization: Vaccination Coverage Senegalese among Children. Medicina **(B** Aires) 2019;55(8):480. [Internet]. Available from: https://doi.org/10.3390/medicina550 80480
- 21. Herliana P, Douiri A. Determinants of immunisation coverage of children aged 12-59 months in Indonesia: A cross-sectional study. BMJ Open [Internet]. 2017;7:1–14. Available from: https://doi.org/10.1136/bmjopen-2016-015790
- 22. Demographic Health Survey. The DHS Program. 2017. Protecting the Privacy of DHS Survey Respondents. Available from: https://www.dhsprogram.com/Meth odology/Protecting-the-Privacy-of-DHS-Survey-Respondents.cfm
- 23. Hardhantyo M, Chuang YC. Urban-

rural differences in factors associated with incomplete basic immunization among children in Indonesia: A nationwide multilevel study. Pediatr Neonatol [Internet]. 2021;62(1):80– 9. Available from: https://doi.org/10.1016/j.pedneo.202 0.09.004

- 24. Central Bureau of Statistics. Statistik Indonesia 2019 [Indonesian Statistics 2019] [Internet]. Jakarta, Indonesia; 2019. Available from: https://www.bps.go.id/publication/2 019/07/04/daac1ba18cae1e90706ee 58a/statistik-indonesia-2019.html
- 25. Ministry of Health. Profil Direktorat Surveilans dan Karantina Kesehatan Tahun 2016 [Profile of The Directorate of Health Survey and Quarantine 2016] [Internet]. 2017. Available from: http://p2p.kemkes.go.id/wpcontent/uploads/2018/04/Profil-2016.pdf
- 26. Asmare G, Madalicho M, Sorsa A. Disparities in full immunization coverage among urban and rural children aged 12-23 months in southwest Ethiopia: A comparative cross-sectional study. Hum Vaccines Immunother [Internet]. 2022;18(6). Available from: https://doi.org/10.1080/21645515.20 22.2101316
- 27. Asm Shahabuddin . Christiana Nöstlinger, Thérèse Delvaux, Malabika Sarker, Alexandre Delamou, Azucena Bardají JEWB, Brouwere V De. Exploring Maternal Health Care-Seeking Behavior of Married Adolescent Girls in Bangladesh: A Social-Ecological Approach. PLoS One [Internet]. 2017;12(1). Available from: https://doi.org/10.1371/journal.pone. 0169109
- 28. Gelaye SS, Yenit MK, Baraki AG. Rural Vaccination Coverage Among Children Aged 12-23 Months Was Higher Than the Urban

Counterparts: A Comparative Cross-Sectional Study in Pawi District, Ethiopia. Pediatr Heal Med Ther [Internet]. 2021;12:119–127. Available from: https://doi.org/10.2147/PHMT.S299 064

- Ameyaw EK, Kareem YO, Ahinkorah BO, Seidu AA, Yaya S. Decomposing the rural-urban gap in factors associated with childhood immunisation in sub-Saharan Africa: Evidence from surveys in 23 countries. BMJ Glob Heal [Internet]. 2021;6(1):1–11. Available from: https://doi.org/10.1136/bmjgh-2020-003773
- 30. Kundu S, Kundu S, Seidu AA, Okyere J, Ghosh S, Hossain A, et al. Factors influencing and changes in childhood vaccination coverage over time in Bangladesh: a multilevel mixed-effects analysis. BMC Public Health [Internet]. 2023;23:1–13. Available from: https://doi.org/10.1186/s12889-023-15711-x
- 31. Adebowale A, Obembe T, Bamgboye E. Relationship between household wealth and childhood immunization in core-North Nigeria. Afr Health Sci [Internet]. 2019;19(1):1582–93. Available from:
- https://doi.org/10.4314/ahs.v19i1.33
- 32. Agbanyo R. Ghana's national health insurance, free maternal healthcare and facility-based delivery services. African Dev Rev [Internet]. 2020;32(1):27–41. Available from: https://doi.org/10.1111/1467-8268.12412
- 33. N'doh Ashken Sanogo SY. Wealth Status, Health Insurance, and Maternal Health Care Utilization in Africa: Evidence from Gabon. Biomed Res Int [Internet]. 2020;(1). Available from: https://doi.org/10.1155/2020/403683 0

- 34. Awasthi A, Pandey CM, Singh U, Kumar S, Singh TB. Maternal determinants of immunization status of children aged 12-23 months in urban slums of Varanasi, India. Clin Epidemiol Glob Heal [Internet]. 2015;3(3):110–6. Available from: http://dx.doi.org/10.1016/j.cegh.201 4.07.004
- 35. Farzad F, Reyer JA, Yamamoto E, Hamajima N. Socio-economic and demographic determinants of full immunization among children of 12-23 months in Afghanistan. Nagoya J Med Sci [Internet]. 2017;79:179–88. Available from: https://doi.org/10.18999/nagjms.79. 2.179
- 36. Mora T, Trapero-Bertran M. The influence of education on the access to childhood immunization: the case of Spain. BMC Public Health [Internet]. 2018;18. Available from: https://doi.org/10.1186/s12889-018-5810-1
- 37. Xeuatvongsa A, Hachiya M, Miyano S, Mizoue T KT. Determination of factors affecting the vaccination status of children aged 12-35 months People's in Lao Democratic Republic. Heliyon [Internet]. 2017;3(3). Available from: https://doi.org/10.1016/j.heliyon.201 7.e00265
- Asif AM, Muhammad Akbar MRT, Arshad IA. Role of Maternal Education and Vaccination Coverage: Evidence From Pakistan Demographic and Health Survey. Asia Pacific J Public Heal [Internet]. 2019;31(8):679–88. Available from: https://doi.org/10.1177/1010539519 889765
- 39. Castillo-Zunino F, Hester KA, Keskinocak P, Nazzal D, Freeman of childhood MC. Association vaccination with family planning, healthcare access, and women analysis education: of Nepal. Senegal and Zambia. medRxiv

[Internet]. 2022; Available from: https://doi.org/10.1101/2022.11.26.2 2282789

- 40. Forshaw J, Gerver SM, Gill M, Cooper E, Manikam L, Ward H. The global effect of maternal education on complete childhood vaccination: A systematic review and metaanalysis. BMC Infect Dis [Internet]. 2017;17(1):1–16. Available from: https://doi.org/10.1186/s12879-017-2890-y
- 41. Onsomu EO, Abuya BA, Okech IN, Collins-McNeil Moore D. J. Maternal education and immunization status among children in Kenya. Matern Heal Child J 2015:19:1724-1733. [Internet]. Available from: https://doi.org/10.1007/s10995-015-1686-1
- 42. Saitoh A, Saitoh A, Sato I, Tomohiro Shinozaki SN. Current practices and needs regarding perinatal childhood immunization education for Japanese mothers. Vaccine [Internet]. 2015;33(45):6128–33. Available from: https://doi.org/10.1016/j.vaccine.20

15.08.069

- 43. Third SJ, Elshafei M, Buttenheim A. Social determinants of communitylevel human papillomavirus vaccination coverage in aschoolbased vaccination programme. Sex Transm Infect [Internet]. 2018;94(4):248–53. Available from: https://doi.org/10.1136/sextrans-2017-053357
- 44. Hargono A, Syahrul F, Indriani D, Chalidyanto D, Megatsari H, Artanti KD, et al. Parents' Knowledge about Immunization with Missed Opportunity for Vaccination in Children. Malaysian J Med Heal Sci [Internet]. 2022;18:101–6. Available from:

https://medic.upm.edu.my/upload/d okumen/2022110210162915_1481. pdf

45. Balogun SA, Yusuff HA, Yusuf KQ, Al-Shenqiti AM, Balogun MT, Tettey P. Maternal education and child immunization: the mediating roles of maternal literacy and socioeconomic status. Pan Afr Med J [Internet]. 2017;26:1–8. Available from: https://doi.org/10.11604/pamj.2017.

26.217.11856

- 46. Samir A Abadura WTL, Kulkarni U, Mekonnen ZA. Individual and community level determinants of childhood full immunization in Ethiopia: a multilevel analysis. BMC Public Health [Internet]. 2015;15. Available from: https://doi.org/10.1186/s12889-015-2315-z
- 47. Holipah M, Kuroda Y. Determinants of immunization status among 12 to 23 month old children in Indonesia (2008–2013): A multilevel analysis. BMC Public Health [Internet]. 2018;18:1–11. Available from: https://doi.org/10.1186/s12889-018-5193-3
- 48. Edayani S, Suryawati I. Obstacles Of Immunization Coverage In Children In Aceh Utara District. Idea Nurs J [Internet]. 2019;10(3):50–7. Available from: https://doi.org/10.52199/inj.v10i3.1 9759
- 49. Suza DE, Efendi F, Hadisuyatmana S, Astutik E. Factors associated with the use of contraception among women age 15 24 years in indonesia. Sys Rev Pharm [Internet]. 2020;11(5):234–40. Available from: https://doi.org/10.31838/srp.2020.5. 35