



***In Vitro* Evaluation of Antidiabetic and Anti-Inflammatory Activities of Five Selected *Syzygium* Leaves Ethanolic Extract as Alpha-Glucosidase Inhibitors and Anti-denaturation of Bovine Serum Albumin (BSA)**

Meiliza Ekayanti*, Feri Setiadi, Siti Aminah, Alam Almaaz Afandi
Department of Pharmacy, STIKes Prima Indonesia, Bekasi, Indonesia

*Corresponding author: meilizaekayanti@gmail.com

Orcid ID: 0000-0002-5541-8610

Submitted: 10 September 2024

Revised: 13 April 2025

Accepted: 24 April 2024

Abstract

Background: Diabetes mellitus (DM) has become the main health problem in the world with a continuous increase in mortality due to complications caused by hyperglycemia. The chronic hyperglycemia is often associated with inflammation due to increase production of free radicals. **Objective:** This study's main objective is to assess antidiabetic and anti-inflammatory properties in vitro of five particular *Syzygium* leaves extract (*S. cumini*, *S. aqueum*, *S. malaccense*, *S. polyanthum*, and *S. aromaticum*) using alpha-glucosidase and Bovine Serum Albumin (BSA). **Methods:** The five of selected *Syzygium* leaves were macerated by using ethanol 96%, each extract was assessed in vitro for antidiabetic activity by analyzing the inhibitory of alpha-glucosidase using acarbose as strandard, and anti-inflammatory activity by analyzing the inhibitory denaturation of BSA Heat-induced and BSA induced by 2,2-diphenyl-1-picrylhydrazine (DPPH) with Sodium diclofenac as standard. **Result:** The result of the greater IC_{50} of α -glucosidase inhibition was *S. malaccense* 76.235 $\mu\text{g/mL}$ (strong) and acarbose standard was 0.241 $\mu\text{g/mL}$ (very strong). The greater IC_{50} of antidenaturation of BSA with heat-induced was *S. polyanthum* (95.7 $\mu\text{g/mL}$) and sodium diclofenac standard (59.25 $\mu\text{g/mL}$) both were strong inhibitor. Along with greater antidenaturation of BSA DPPH-induced was *S. malaccense* (90.320 $\mu\text{g/mL}$) and sodium diclofenac standard (43.301 $\mu\text{g/mL}$) both were strong inhibitor. **Conclusion:** Ethanol extraxt of *Syzygium* leaves were potential to be developed as an antidiabetic and anti-inflammatory herbal medicine, particularly *S. malaccense* and *S. polyanthum* leaves extract which provide greater activity on this study.

Keywords: Antidiabetic, *Syzygium*, α -glucosidase, Anti-denaturation, Bovine Serum Albumin.

How to cite this article:

Ekayanti, M., Setiadi, F., Aminah, S. & Afandi, A. A. (2025). *In Vitro* Evaluation of Antidiabetic and Anti-Inflammatory Activities of Five Selected *Syzygium* Leaves Ethanolic Extract as Alpha-Glucosidase Inhibitors and Anti-denaturation of Bovine Serum Albumin (BSA). *Jurnal Farmasi dan Ilmu Kefarmasian Indonesia*, 12(1), 85-94. <http://doi.org/10.20473/jfiki.v11i32025.85-94>

INTRODUCTION

The increasing prevalence and continuity of Diabetes Mellitus (DM) has become the main health problem in the world with a consistent increase in mortality due to complications of the disease (Zimmet *et al.*, 2016). The IDF was reported 463 million adults are diabetes and the projection in 2045 will estimated 700 million (International Diabetes Federation, 2019). The DM complications were predicted to be the cause of death of 4.2 million adults and its comparable to one mortality every eight seconds (Zimmet *et al.*, 2016). The progress of Type-2 diabetes mellitus (T2DM) complications in chronic hyperglycemia induces oxidative stress and inflammation, its simultaneously promoting complications including renal and cardiovascular diseases (Charlton *et al.*, 2021; Yuan *et al.*, 2019).

The increase in global spending on DM is reported to be more than 700 billion USD (International Diabetes Federation, 2019). The side effects of prolonged consumption of conventional drugs as well as the less invasive prevention and therapeutics using natural medicine are one of the problems in the success of DM therapy. The COVID-19 pandemic has also contributed to the increase in T2DM, so it is necessary to develop easy and inexpensive natural ingredient therapies to overcome the increased inflammation of T2DM so as to reduce diabetes complications (Salleh *et al.*, 2021). Most T2DM therapies are reported to have side effects of gastrointestinal disorders including the use of α -glucosidase Inhibitor (AGI) drugs due to the degradation of undigested carbohydrates by colon bacteria, causing excessive gas formation with a case percentage of 78% (Kumar *et al.*, 2018). The result of Riset Kesehatan Dasar (Riskesdas) (2018) reported that based on the doctor's diagnosed the prevalence of DM in Indonesia at the aged over 15 years was 2%, an increase from the previous research results in 2013 of 1.5%. Based on blood sugar test findings, the prevalence of diabetes mellitus increased from 6.9% (2013) to 8.5% (2018). The pattern of increase of DM showed higher in the age ranged 56-64 years and 66-74 years (Kemenkes RI, 2020).

Increased production of free radicals, particularly reactive oxygen species (ROS), can result from hyperglycemia and lead to oxidative stress. The disproportion between ROS and antioxidant defenses not only causes direct cell damage but also inflammation that results in tissue damage. Diabetes is often associated with inflammation when biochemical changes in diabetes affect the increase of TNF- α and IL-1 β which leads to an increase in ROS by mitochondria. Emerging therapeutic strategies address this pathway in a different way, ranging from enhancing free radical scavenging

(antioxidants and Nrf2 activators) to reduce ROS production such as NADPH oxidase inhibitors and XO inhibitors or inhibiting associated inflammatory pathways (NLRP3 inflammation inhibitors, lipoxins, GLP, receptor -1 agonists and AT-1 receptor antagonists) (Asmat *et al.*, 2016; Oguntibeju, 2019; Zhang *et al.*, 2021). The oxidative stress reported to trigger the development of micro- and macro-vascular damage complications in T2DM and hyperglycemia, as well as being responsible for DNA, lipid and protein damage associated with ROS production (Oguntibeju, 2019).

Natural ingredient therapy is currently in great demand because it is relatively cheaper and easier to obtain to become one of the options in the treatment of DM, especially T2DM. Natural medicine is projected to play a role in overcoming DM complications and the problem of side effects of conventional drug consumption, especially if used for a long period of time. *Syzygium*, a genus of plants from the *Myrtaceae* family is one of the largest genera of flowering plants with a total of 1800 species and is distributed in areas that mainly have a tropical climate including Indonesia (Abdullah *et al.*, 2021; Kavitha & Poonguzhali, 2021). Plants of the *Myrtaceae* family are reported to be the 20 largest ethnomedicinal families in Indonesia with 5 (five) species of the *Syzygium* genus (Hidayat, 2021). Some of the species of the genus *Syzygium* studied are *S. polyanthum*, *S. aromaticum*, *S. aqueum*, *S. cumini* and *S. malaccense*. The *Syzygium* group of plant species is reported to have been used for generations in traditional Ayurvedic medicine in India (Cock & Cheesman, 2018).

Studies conducted by Zaen & Ekayanti (2022) and Aklimah & Ekayanti (2022) on antioxidants from leaf extracts of several *Syzygium* genus plants showed very strong antioxidants in ethanol extracts of *Syzygium* plant leaves with IC₅₀ values *Syzygium aromaticum* 3.026 \pm 1.699 μ g/ml, *Syzygium polyanthum* 3.555 \pm 2.776 μ g/ml, *Syzygium aqueum* 5.416 \pm 2.588 μ g/mL, *Syzygium malaccense* 3.297 \pm 2.595 μ g/mL and *Syzygium cumini* 2.416 \pm 1.543 μ g/mL (Aklimah & Ekayanti, 2022; Zaen & Ekayanti, 2022). The phenetic study of each five *Syzygium* was conducted as an initial screening for the further investigation of pharmacological activities with ROS-related mechanisms. The purpose of this study was to analyze the α -glucosidase inhibition test and anti-denaturation of Bovin Serum Albumin (BSA) protein of several *Syzygium* leaf extracts as a therapy for T2DM and inflammation through the mechanism of increasing antioxidant defense, regulating carbohydrate metabolism and inhibiting inflammatory pathways. The novelty of this study is anti-denaturation BSA induced 2,2-diphenyl-1-picrylhydrazil (DPPH) of five *Syzygium* leaves extract. This study is expected to support and

improve references in the development of herbal medicine therapy as antidiabetes and anti-inflammatory. The α -glucosidase enzyme inhibition and protein anti-denaturation of *Syzygium* genus plant species that has been done before is limited to one activity test while diabetes is associated with inflammation so it is necessary to analyze the potential of natural medicine with antidiabetic and anti-inflammatory activities.

MATERIALS AND METHODS

Materials

Alpha-glucosidase from *Bacillus stearothermophilus*, 4-Nitrophenyl α -D-glucopyranoside and Bovine Serum Albumin (BSA) purchased from Sigma-Aldrich; Ethanol, Aluminum trichloride, Potassium acetate, Methanol, Citric acid, Ethyl acetate, Formic acid, silica gel F254 TLC Plates were from Merck; and DPPH (2,2-Diphenyl-1-picryl hydrazine) was from Himedia. Leaves of five grounded of *Syzygium* (*S. aromaticum*, *S. polyanthum*, *S. aqueum*, *S. malaccense*, and *S. cumini*) were collected from Balai Penelitian Tanaman Rempah dan Obat (Bogor, Indonesia). Each five plants of *Syzygium* were conducted for the identification of plant species at the Research Center of Biosystematics and Conservation (Bogor, Indonesia).

Methods

Plant extraction

Five *Syzygium* leaves powder were weighed 500 g and extracted by maceration method with 96% ethanol (1:10 w/v), soaked for three days, and stirred periodically. The filtrate of the first maceration was filtered and repeated two replication. The filtrate from maceration and repetition was evaporated at 50 °C by using a vacuum rotatory evaporator (DLAB Rotary Evaporator RE-100 Pro) (Wahyulianingsih *et al.*, 2016).

Determination of anti-diabetic activity by inhibition of α -glucosidase (Akmal & Roopma, 2023)

Initial enzyme activity assay: A total of 60 μ L of 0.1M phosphate-buffered saline (pH6.8) was added to 50 μ L of 0.07 U/mL enzyme solution and incubated in a 96 well micro plate at 37 °C for 20 minutes. After preincubation 50 μ L of 2mM pNPG was added into the microplate and then incubated again at 37 °C. The last stage was stopped by adding 160 μ L of sodium carbonate solution to the micro-plate well and evaluating the absorbance with a wavelength of 425 nm. Enzyme activity

assay - *Syzygium* leaf extract: A total of 60 μ L of 0.1M of each *Syzygium* leaves ethanol extract were added with 50 μ L of 0.07 U/mL enzyme solution and incubated in a 96 well micro plate at 37 °C for 20 minutes. After preincubation 50 μ L of 2mM pNPG was added into the microplate and then incubated again at 37 °C. The last stage was stopped by adding 160 μ L of sodium carbonate solution to microplate and reading the absorbance with a wavelength of 425 nm.

Acarbose-enzyme activity assay: A total of 60 μ L acarbose was added to 50 μ L of 0.07 U/mL enzyme solution and incubated in a 96 well micro plate at 37 °C for 20 minutes. After preincubation 50 μ L of 2mM pNPG was added into the micro plate and incubated again at 37 °C. The last stage was stopped by adding 160 μ L of sodium carbonate solution to the cuvette and reading the absorbance with a wavelength of 425 nm. The assay of initial enzyme blank activity and enzyme inhibition (BEA and BIE): A total of 60 μ L of phosphate solution of each *Syzygium* leaves extract and acarbose standard was added to 50 μ L of 0.07 U/mL enzyme solution and incubated in a 96 well micro plate at 37 °C for 20 minutes. After preincubation 50 μ L of 2mM PNPNG was added into the micro plate and then incubated again at 37 °C. The last stage was stopped by adding 160 μ L of sodium carbonate solution and reading the absorbance with a wavelength of 425 nm by using Biotek Epoch Microplate Spectrophotometer.

Determination of anti-inflammatory activity by inhibition denaturation of heat-induced bovine serum albumin (Williams *et al.*, 2008)

The assay of anti-inflammatory activity five selected *Syzygium* extracts were conducted using a heat-induced Bovine Serum Albumin (BSA) denaturation assay. Tris-buffer Saline 0.05 M was used to create the stock solutions of BSA 5% (w/v), which was then adjusted to pH 6.8 using glacial acetic acid. 100 μ L-aliquots of the BSA stock solution and distilled water were mixed with varying test tube of the extract to create 1.0 mL sample that included the extract at different dilutions. The samples were heated in a waterbath for seven minutes at 70°C to cause protein denaturation. The solutions cooled to room temperature. Spectrophotometry UV-Vis (PG-Instrument) was used to quantify the turbidity of the solutions at wavelength of 517 nm. As controls, solutions with distilled water in place of the extract were used; values obtained with these

preparations were interpreted as indicating 100% protein denaturation. BSA-free samples served as blanks and Diclofenac's anti-denaturing properties served as positive control in concurrent studies. The following formula was used to determine the percentage inhibition of denaturation, where Abs control is the absorbance of the controls, Absorbance (Abs) of sample of the *Syzygium* leaves extract or diclofenac samples, and Abs blank of the blank.

$$\% \text{ inhibition of denaturation} = \left[\frac{\text{Abscontrol} - (\text{Abssample} - \text{Absblank})}{\text{Abscontrol}} \right] \times 100\%$$

Determination of anti-inflammatory activity by inhibition denaturation of DPPH-induced bovine serum albumin (Alam *et al.*, 2022)

The anti-inflammatory activity of five selected *Syzygium* extracts was analyzed by a modified method using a DPPH-induced BSA denaturation assay. The radical scavenging activity (RSA) adopted to measure anti-inflammatory activity using the DPPH method. Briefly, 2 mL of extract solution (1–100 µg/mL) in methanol and BSA stock solutions (prepared as the same method above) was added to 2 mL of DPPH (0.1 mM) solution. The mixtures were kept aside in a dark area for 30 min and absorbance was measured at λ_{max} 517 nm against an equal amount of DPPH and methanol as a blank. The percentage of DPPH• scavenging (% Scavenging of DPPH) was estimated using the equation:

$$\% \text{ Scavenging of DPPH} \bullet = \left[\frac{(A_0 - A_1)/A_0}{1} \right] \times 100$$

Statistical Analysis

All the experiments for the determination of inhibitory activity of alpha-glucosidase and denaturation of BSA Heat-induced, and denaturation of BSA by using DPPH-induced have been conducted triplet (n=3). The values are expressed as the mean \pm standard deviation (SD).

RESULTS AND DISCUSSION

Syzygium is a genus of plants from Myrtaceae family which is one of the largest genera of flowering plants with total 1800 species and distributed in regions mainly have a tropical climate including Indonesia (Abdullah *et al.*, 2021). The five fresh leaves of *Syzygium* were determined at the Research Center of Biosystematics and Conservation (Bogor, Indonesia) and it was validated as *Syzygium aromaticum*, *S. polyanthum*, *S. aqueum*, *S. malaccense*, and *S. cumini*. The extract of five leaves of *Syzygium* was done by cold extraction method (maceration). The crude extract of five leaves of *Syzygium* was obtained green-brownish with a distinctive aroma of leaves and the result of the higher yield percentage was *Syzygium aromaticum* (32.90%) and *Syzygium polyanthum* (30.19%) shown in Table 1. The yield percentages of five *Syzygium* extracts obtained fulfilled the quality requirements of the extraction standard as values above 10% (Kementrian Kesehatan RI, 2017).

Table 1. Yield Percentages of Five *Syzygium* Leaves Extracts

Sample	Yield (%)
<i>Syzygium cumini</i>	24.380
<i>Syzygium aqueum</i>	18.556
<i>Syzygium malaccense</i>	23.756
<i>Syzygium polyanthum</i>	30.19
<i>Syzygium aromaticum</i>	32.90

Several species of *Syzygium*, including *S. cumini*, *S. polyanthum*, *S. aqueum*, *S. aromaticum*, and *S. malaccense*, have been shown to exhibit enzymatic inhibitory action (Zulcafli *et al.*, 2020). The absorbance measurement results then calculate to percentage inhibition value. The activity of the *Syzygium* extracts mostly inhibited 50% of the enzyme activity at 100–200 µg/mL unless *S. malaccense* inhibited <100 µg/mL. The result of the α -glucosidase enzyme inhibition study showed *S. malaccense* was the greater inhibitory activity with IC₅₀ value 76.235 µg/mL (Table 2). The

concentration 200 µg/mL of *Syzygium cumini* obtained percentage of inhibition 63.261 \pm 0.178% (Figure 1). The inhibitory activity of acarbose is greater compared to each *Syzygium* extracts with IC₅₀ values 0.241 µg/mL. Acarbose acts by competitive and reversible inhibition of α -amylase and α -glucosidase from the pancreas (Glittenberg, 2012). It is well known that flavonoids derived from plants have antidiabetic effects (Najafian *et al.*, 2012; Yoshikawa *et al.*, 1998). Ethanolic extract of *Syzygium* leaves reported have high flavonoid content (Aklimah & Ekayanti, 2022;

Zaen & Ekayanti, 2022). Flavonoid and other polyphenols are reported potentially inhibit α -amylase and α -glucosidase without adverse gastrointestinal effects and useful to T2DM therapy (Barber *et al.*, 2021). Many flavonoids have a higher inhibition of α -glucosidase and leading to slow-release effect that of α -amylase which may be favoured over acarbose to decrease postprandial glucose without uncomfortable side effects (Barber *et al.*, 2021). The active substances myricetin-3-O-rhamnoside and europetin-3-O-rhamnoside, which were separated from *S. aqueum*, inhibited α -glucosidase (Manaharan *et al.*, 2012). Maslinic acid (MA) and Oleanolic acid (OA) produced from *S. aromaticum* were observed to decrease the expression of α -glucosidase, and α -amylase in the small intestines of STZ-induced diabetic rats (Khathi *et al.*, 2013). The active component of *S. cumini* and *S. malaccense*, myricitrin, was found to inhibit α -glucosidase and α -amylase (Khathi *et al.*, 2013).

The strong antioxidant activity of some *Syzygium* leaves extract reported in the previous study (Aklimah & Ekayanti, 2022; Zaen & Ekayanti, 2022), a problem-solving approach as well as a therapeutic strategy for DM and inflammation through the mechanism of increasing antioxidant defenses, improving carbohydrate metabolism profiles and inhibiting

inflammatory pathways so as to reduce the risk of hyperglycemia complications (Akmal & Roopma, 2023; Shaw *et al.*, 2017). According to Ekayanti *et al.* (2018) Phenolic components in natural materials are known to bind to proteins of enzymes and form enzyme-inhibitor bonds, thereby reducing enzyme activity (Ekayanti *et al.*, 2018). Polyphenols are also reported to have therapeutic effects on DM vascular dysfunction (Nor *et al.*, 2022). The antioxidant activity of ethanol extract of *Syzygium* plant leaves is expected to overcome the ROS imbalance that triggers oxidative stress in T2DM (Oguntibeju, 2019). Further studies in the development of T2DM and inflammation therapy need to be carried out by analyzing the α -glucosidase enzyme inhibitory activity and anti-denaturation of Bovine Serum Albumin (BSA) protein in ethanol extracts of the leaves of several *Syzygium* plants. Alpha Glucosidase Inhibitor (AGI) is one of the effective therapeutic groups used as T2DM treatment in improving metabolic profiles and potentially reducing the risk of long-term hyperglycemia complications (Akmal & Roopma, 2023). The α -glucosidase is an enzyme secreted from the epithelium of the small intestine that is responsible for carbohydrate degradation by hydrolyzing complex carbohydrates into simple glucose that can eventually be absorbed (Syabana *et al.*, 2022).

Table 2. Inhibitory percentages and IC₅₀ values of α -glucosidase

Sample	Inhibitor Percentages (%)					IC ₅₀ values (μ g/mL)
	1	2	3	4	5	
Acarbose	36.057	61.305	70.687	93.731	96.930	0.241 \pm 0.478
<i>S. cumini</i>	6.667	17.609	29.457	44.094	63.261	123.239 \pm 0.224
<i>S. aqueum</i>	28.659	37.391	43.804	48.225	56.087	169.676 \pm 0.237
<i>S. malaccense</i>	35.000	43.841	49.638	54.565	60.870	76.235 \pm 0.234
<i>S. polyanthum</i>	31.486	38.261	43.768	48.949	52.790	198.222 \pm 0.392
<i>S. aromaticum</i>	2.826	17.283	27.971	40.109	58.370	144.698 \pm 0.186

Values are mean \pm SD (n=3). Acarbose concentrations (0.1, 0.5, 1.0, 5.0, and 10.0 ppm), Sample extract concentrations (25, 50, 75 100, 200 ppm)




Figure 1. α -Glucosidase Inhibitor of Five Leaves *Syzygium* Extract

The production of free radicals leads to protein denaturation in the body, which triggers the release of inflammatory mediators and triggers inflammatory pathways (Shaw *et al.*, 2017). The method used to reduce BSA volume and stock solution of compound or extract for assessing anti-denaturation (anti-inflammatory) activity. The results presented at Table 3 and Figure 2 are representation of *Syzygium* ethanolic extracts and Sodium diclofenac as a control positive. Anti-denaturation protein assay using heat induction was increase kinetic energy and cause the molecules vibrating and move quickly, disrupting hydrogen bond and non-polar hydrophobic interactions of the protein. Sodium diclofenac is a non-steroidal anti-inflammatory drug works non-selectively and has better solubility in water and organic solvents. The concentration of sodium

diclofenac is 5, 15, 25, 50, and 75 $\mu\text{g/mL}$. The concentrations of *Syzygium* ethanolic extracts is 25, 50, 75, 100, and 200 $\mu\text{g/mL}$. The concentrations of *Syzygium* leaves extract can inhibit protein denaturation by >50% at range 100-200 $\mu\text{g/mL}$. The inhibition activity of protein denaturation due to the presence of bioactive compounds. The inhibition of BSA denaturation induce by heating resulted the higher IC_{50} for *Syzygium polyanthum* (95.7 $\mu\text{g/mL}$) compared to standard (sodium diclofenac) was 59.25 $\mu\text{g/mL}$. *Syzygium* extract was reported rich of polyphenol compound (Sobeh *et al.*, 2018). Polyphenol, phenyl propanoids and the disulphides interacting with the aliphatic regions around the lysine residue on the BSA and reported suitable as an anti-oxidants, anticancer, and anti-glycation (Williams *et al.*, 2008).

Table 3. Inhibitory percentages and IC_{50} values of Anti-denaturation BSA protein heat-induced

Sample	Anti-Denaturation Percentages BSA Protein Heat-Induced (%)					IC_{50} ($\mu\text{g/mL}$)
	1	2	3	4	5	
Sodium diclofenac	22.459	26.089	47.289	63.771	82.720	59.25 \pm 0.557
<i>S. cumini</i>	6.968	17.608	33.380	40.301	50.964	200.270 \pm 1.331
<i>S. aqueum</i>	8.061	13.724	32.054	41.363	51.727	198.26 \pm 0.398
<i>S. malaccense</i>	13.876	28.469	40.072	50.598	55.801	182.73 \pm 0.143
<i>S. polyanthum</i>	20.754	34.330	42.883	58.194	65.670	95.7 \pm 0.448
<i>S. aromaticum</i>	27.213	39.533	43.002	44.797	59.988	164.15 \pm 0.717

Values are mean \pm SD (n=3), Sodium diclofenac concentrations (5, 15, 25, 50, and 75 ppm), Sample extract concentrations (25, 50, 75, 100, 200 ppm)




Figure 2. Anti-Denaturation BSA of Five Leaves *Syzygium* Extract Heat-Induced

Table 3. Inhibitory percentages and IC₅₀ values of Anti-denaturation BSA protein DPPH-induced

Sample	Anti-Denaturation Percentages (%)					IC ₅₀ (μg/mL)
	1	2	3	4	5	
Sodium diclofenac	13.427	31.841	49.916	56.432	61.338	43.301±0.422
<i>S. cumini</i>	20.103	27.890	36.710	49.066	63.886	117.233±1.120
<i>S. aqueum</i>	16.530	29.006	35.322	43.353	56.179	180.320±1.282
<i>S. malaccense</i>	8.782	18.306	40.198	52.628	62.400	90.320±1.076
<i>S. polyanthum</i>	13.417	22.075	28.900	44.150	51.404	196.470±0.405
<i>S. aromaticum</i>	11.817	18.760	32.722	45.008	59.165	171.596±0.479

Values are mean ± SD (n=3), Sodium diclofenac concentrations (5, 15, 25, 50, and 75 ppm), Sample extract concentrations (25, 50, 75, 100, 200 ppm)




Figure 3. Anti-Denaturation BSA of Five Leaves *Syzygium* Extract DPPH-Induced

Low doses of BSA have been found as a channel for broad-spectrum *in vitro* study to analyze possible therapeutic prototypes (Williams *et al.*, 2008). The modified method of anti-denaturation BSA was designed relevant to hyperglycemia chronic conditions which inflammatory caused by increasing free radicals and stress oxidative. In this method, BSA was denaturated by DPPH-inducing and plays role as free radicals and denaturing agent as well. DPPH compound is used as a substrate to evaluate antioxidant activity, it was the stable free radical has an ability to accept one electron or Hydrogen. The greater anti-denaturation BSA DPPH-induced was *Syzygium malaccense* with IC₅₀ values is 90.320 µg/mL (Table 3) and the percentage chart showed at Figure 3. *Syzygium* leaves extract can inhibit BSA denaturation which induced by DPPH at range 100-200 µg/mL >50% inhibition percentage. The IC₅₀ value less than 50 Sodium diclofenac (positive control) is categorized as having a very strong inhibitory effect on protein denaturation. The ability of blocking the Cyclooxygenase (COX) and Lipoxygenase-5 (LOX-5) pathways was one of the pathway to overcome inflammation. Flavonoid substances are known to have anti-inflammatory properties while tannin and saponin compounds maintain membranes by attaching to cations, flavonoid and saponin compounds have been shown to have anti-inflammatory properties via scavenging free radicals. The erythrocyte membranes from hypotonic solutions could be stabilized by free radical inhibitors (Shalihah *et al.*, 2021).

CONCLUSION

Diabetes mellitus (DM) is has become the main health problem in the world with a consistent increase in mortality due to complications of the disease. The chronic hyperglycemia is often associated with inflammation due to increase production of free radicals thus lead to oxidative stress. *Syzygium* leaves were potential to be developed as antidiabetic and anti-inflammatory candidates of drugs. The greater result of IC₅₀ of α-glucosidase inhibition was *Syzygium malaccense* was 76.235 µg/mL (strong) and acarbose standard was 0.241 µg/mL (very strong). The greater antidenaturation of BSA with heat-induced was *Syzygium polyanthum* (95.7 µg/mL) and sodium diclofenac standard was 59.25 µg/mL both were strong inhibitor. Along with antidenaturation of BSA DPPH-induced was *Syzygium malaccense* 90.320 µg/mL and

sodium diclofenac standard was 43.301 µg/mL both were strong inhibitor. Further research is needed to confirm the antidiabetic and anti-inflammatory activities of *Syzygium* extract using different methods *in vitro* and *in vivo*.

ACKNOWLEDGMENT

The authors are grateful to Ministry of Education Culture, Research, and Technology of the Republic of Indonesia whom funded the study. Acknowledgment is also due to all the colleagues in accomplishing this study.

REFERENCES

- Abdullah, M., Priyono, B., Kartijono, N. E. F. & Bodijantoro, P. M. H. (2021). Medicinal Plant of Gunung Prau, Indonesia: Exploration and Ethno-Botanical Study. *Journal of Physics: Conference Series*; 1918; 1–4. doi: 10.1088/1742-6596/1918/5/052046.
- Aklimah, M. & Ekayanti, M. (2022). Penetapan Flavonoid Total dan Uji Aktivitas Antioksidan Ekstrak Etanol Daun Cengkeh (*Syzygium aromaticum* (L). Merr) dan Daun Salam (*Syzygium polyanthum* Thwaites). *Jurnal Kedokteran Universitas Palangka Raya*; 10; 11–14. doi: 10.37304/jkupr.v10i2.5536.
- Akmal, M., Patel, P. & Roopma, W. (2023). Alpha Glucosidase Inhibitors. *StatPearls Publishing LLC*. <https://www.ncbi.nlm.nih.gov/books/>
- Alam, T., Ekayanti, M., Permana, N., Hadissabil, Z. & Prima Indonesia, S. (2022). Potensi Aktivitas Antioksidan Ekstrak Etanol dan Fraksi Ciplukan (*Physalis angulata*) pada DPPH (1,1-difenil-2-pikrihidrazil) Potential Antioxidant Activity of Ethanol Extract and Fraction of Ciplukan (*Physalis angulata*) on DPPH (1,1-diphenyl-2-picrylhydrazyl). *Jurnal Farmasi Indonesia*; 19; 193–199. doi: 10.31001/jfi.v19i1.1490.
- Asmat, U., Abad, K. & Ismail, K. (2016). Diabetes Mellitus and Oxidative Stress a Concise Review. In *Saudi Pharmaceutical Journal*; 24; 547–553. Elsevier B.V. doi: 10.1016/j.jsps.2015.03.013.
- Barber, E., Houghton, M. J. & Williamson, G. (2021). Flavonoids as Human Intestinal α-glucosidase Inhibitors. *Foods*; 10; 1–22. doi: 10.3390/foods10081939.

- Charlton, A., Garzarella, J., Jandeleit-Dahm, K. A. M., & Jha, J. C. (2021). Oxidative Stress and Inflammation in Renal and Cardiovascular Complications of Diabetes. *In Biology*; 10; 1–18. MDPI AG. doi: 10.3390/biology10010018.
- Cock, I. E. & Cheesman, M. (2018). Plants of The Genus *Syzygium* (Myrtaceae): A Review on Ethnobotany, Medicinal Properties and Phytochemistry. In *Bioactive Compounds of Medicinal Plants: Properties and Potential for Human Health*; 35–85.
- Ekayanti, M., Sauriasari, R. & Elya, B. (2018). Dipeptidyl peptidase IV Inhibitory Activity of Fraction from White Tea Ethanolic Extract (*Camellia sinensis* (L.) Kuntze) ex vivo. *Pharmacognosy Journal*; 10; 190–193. doi: 10.5530/pj.2018.1.32.
- Glittenberg, D. (2012). Starch-Based Biopolymers in Paper, Corrugating, and Other Industrial Applications. *Polymer Science: A Comprehensive Reference*; ;1–10; 165–193. doi: 10.1016/B978-0-444-53349-4.00258-2.
- Hidayat, S. (2021). Pemanfaatan Tumbuhan Obat oleh Beberapa Etnis di Indonesia. *Prosiding Seminar Nasional PMEI Ke-V*, 177–185. <https://jte.pmei.or.id/index.php/jte/article/view/141>
- International Diabetes Federation. (2019). IDF Diabetes Atlas 9th Ed. In *Diabetes Atlas* (9th ed.). <https://www.diabetesatlas.org>.
- Kavitha, V. & Poonguzhali, T. V. (2021). Phylogenetic Analysis on Different Species of *Syzygium*. *International Research Journal of Plant Science*; 12; 1–03. doi: 10.14303/irjps.2021.001
- Kemenkes RI. (2020). InfoDatin (Pusat Data dan Informasi Kementerian Kesehatan RI): Tetap Produktif, Cegah, dan Atasi Diabetes Melitus. *Kementerian Kesehatan RI*, 4–5. <https://www.kemkes.go.id/downloads/resources/download/pusdatin/infodatin/Infodatin%202020%20Diabetes%20Melitus.pdf>
- Kementerian Kesehatan RI. (2017). *Farmakope Herbal Indonesia Edisi II*.
- Khathi, A., Serumula, M. R., Myburg, R. B., Van Heerden, F. R. & Musabayane, C. T. (2013). Effects of *Syzygium aromaticum*-Derived Triterpenes on Postprandial Blood Glucose in Streptozotocin-Induced Diabetic Rats Following Carbohydrate Challenge. *PLoS ONE*; 8; 1–8. doi: 10.1371/journal.pone.0081632.
- Kumar, Y., Goyal, R. K. & Thakur, A. K. (2018). Pharmacotherapeutics of Miglitol: An α -Glucosidase Inhibitor. *Journal of Analytical & Pharmaceutical Research*; 7; 617–619. doi: 10.15406/japlr.2018.07.00292.
- Manaharan, T., Appleton, D., Cheng, H. M., & Palanisamy, U. D. (2012). Flavonoids isolated from *Syzygium aqueum* Leaf Extract as Potential Antihyperglycaemic Agents. *Food Chemistry*; 132; 1802–1807. doi: 10.1016/j.foodchem.2011.11.147.
- Najafian, M., Jahromi, M. Z., Nowrozejjad, M. J., Khajeaian, P., Kargar, M. M., Sadeghi, M. & Arasteh, A. (2012). Phloridzin Reduces Blood Glucose Levels and Improves Lipids Metabolism in Streptozotocin-Induced Diabetic Rats. *Molecular Biology Reports*; 39; 5299–5306. doi: 10.1007/s11033-011-1328-7.
- Nor, N. A., Budin, S. B., Zainalabidin, S., Jalil, J., Sopian, S., Jubaidi, F. F. & Anuar, N. N. (2022). Review: The Role of Polyphenol in Modulating Associated Genes in Diabetes-Induced Vascular Disorders. *International Journal of Molecular Sciences*; 23; 6936–6960. doi: 10.3390/ijms23126396.
- Oguntibeju, O. O. (2019). Type 2 Diabetes Mellitus, Oxidative Stress and Inflammation. *Int J Physiol Pathophysiol Pharmacol*; 11; 45–63.
- Oguntibeju, O. O. (2019). Review Article Type 2 Diabetes Mellitus, Oxidative Stress and Inflammation: Examining the Links. *Int J Physiol Pharmacol*; 11; 45–63.
- Salleh, N. H., Zulkipli, I. N., Mohd Yasin, H., Ja' Afar, F., Ahmad, N., Wan Ahmad, W. A. N. & Ahmad, S. R. (2021). Systematic Review of Medicinal Plants Used for Treatment of Diabetes in Human Clinical Trials: An ASEAN Perspective. *Evidence-Based Complementary and Alternative Medicine*; 2021; 1–10. doi: 10.1155/2021/5570939.
- Shalihah, A., Christianty, F. M. & Fajrin, F. A. (2021). Anti-inflammatory Activity of the Ethanol Extract of Cinnamon (*Cinnamomum burmannii*) Bark using Membrane Stabilization Method and Protein Denaturation. In *Indonesian Journal of Pharmaceutical Science and Technology*

- Journal Homepage*; 1; 9-14 doi: 10.24198/ijpst.v1i1.36323.
- Shaw, Y., Williams, L., Green, C., Rodney, S. & Smith, M. (2017). In-Vitro Evaluation of the Anti-Inflammatory Potential of Selected Jamaican Plant Extracts using the Bovine Serum Albumin Protein Denaturation Assay. *Int. J. Pharm. Sci. Rev. Res*; 47; 145–153.
- Sobeh, M., Mahmoud, M. F., Petruk, G., Rezq, S., Ashour, M. L., Youssef, F. S., El-Shazly, A. M., Monti, D. M., Abdel-Naim, A. B. & Wink, M. (2018). *Syzygium Aqueum*: A Polyphenol-Rich Leaf Extract Exhibits Antioxidant, Hepatoprotective, Pain-Killing and Anti-Inflammatory Activities in Animal Models. *Frontiers in Pharmacology*; 9; 1-14. doi: 10.3389/fphar.2018.00566.
- Syabana, M. A., Yuliana, N. D., Batubara, I. & Fardiaz, D. (2022). α -Glucosidase Inhibitors from *Syzygium polyanthum* (Wight) Walp Leaves as Revealed by Metabolomics and in silico Approaches. *Journal of Ethnopharmacology*, 282. 1-13. doi: 10.1016/j.jep.2021.114618
- Wahyulianingsih, W., Handayani, S. & Malik, Abd. (2016). Penetapan Kadar Flavonoid Total Ekstrak Daun Cengkeh (*Syzygium aromaticum* (L.) Merr & Perry). *Jurnal Fitofarmaka Indonesia*; 3; 188–193. doi: 10.33096/jffi.v3i2.221.
- Williams, L., O'Connar, A., Latore, L., Dennis, O., Ringer, S., Whittaker, J., Conrad, J., Volger, B., Rosner, H. & Kraus, W. (2008). The in vitro Anti-denaturation Effects Induced by Natural Products and Non-steroidal Compounds in Heat Treated (Immunogenic) Bovine Serum Albumin is Proposed as a Screening Assay for the Detection of Anti-inflammatory Compounds, without the use of Animals, in the Early Stages of the Drug Discovery Process. *West Indian Medical Journal*; 57; 327–378. doi: 10.1215/9780822388630-010
- Yoshikawa, M., Shimada, H., Nishida, N., Li, Y., Toguchida, I., Yamahara, J. & Matsuda, H. (1998). Antidiabetic Principles of Natural Medicines II. Aldose Reductase and Alpha-Glucosidase Inhibitors from Brazilian Natural Medicine, The Leaves of *Myrcia multiflora* DC. (Myrtaceae): Structures of Myrciacitrins I and II and Myrciaphenones A and B. *Chemical and Pharmaceutical Bulletin*; 46; 113–119. doi: 10.1248/cpb.46.113
- Yuan, T., Yang, T., Chen, H., Fu, D., Hu, Y., Wang, J., Yuan, Q., Yu, H., Xu, W. & Xie, X. (2019). New Insights into Oxidative Stress and Inflammation During Diabetes Mellitus-Accelerated Atherosclerosis. *In Redox Biology*; 20; 247–260. Elsevier B.V. doi: 10.1016/j.redox.2018.09.025.
- Zaen, D. M. & Ekayanti, M. (2022). Penetapan Flavonoid Total dan Uji Aktivitas Antioksidan Ekstrak Etanol Daun Jambu Air (*Syzygium aqueum*), Daun Jambu Bol (*Syzygium malaccense*) dan Daun Jamblang (*Syzygium cumini*). *Jurnal Kedokteran Universitas Palangka Raya*; 10; 15–18. doi: 10.37304/jkupr.v10i2.5531.
- Zhang, C., Wang, X., Du, J., Gu, Z. & Zhao, Y. (2021). Reactive Oxygen Species-Regulating Strategies Based on Nanomaterials for Disease Treatment. *In Advanced Science*; 8; 1-34. doi: 10.1002/advs.202002797.
- Zimmet, P., Alberti, K. G., Magliano, D. J. & Bennett, P. H. (2016). Diabetes Mellitus Statistics on Prevalence and Mortality: Facts and Fallacies. *Nature Reviews Endocrinology*; 12; 616–622. doi: 10.1038/nrendo.2016.105.
- Zulcafli, A. S., Lim, C., Ling, A. P., Chye, S. & Koh, R. (2020). Antidiabetic Potential of *Syzygium* sp.: An Overview. *The Yale Journal of Biology and Medicine*; 93; 307–325.