

A qualitative description of nurses' problems to monitor and supervise the vital signs in covid-19 patients at the isolation room

Tri Yahya Christina¹ , Suhartini Ismail^{2*} , and Meira Erawati³ 

¹ Master Nursing Student Nursing Department Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia.

² Emergency and Critical Care Nursing Division, Nursing Department Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

³ Pediatric Nursing Division, Nursing Department Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

*Correspondence: Suhartini Ismail. Address: Faculty of Nursing, Emergency and Critical Care Nursing Division, Nursing Department Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia. Email: suhartini.ismail@fk.undip.ac.id

Responsible Editor: Yulis Setiya Dewi

Received: 13 February 2022 ◦ Revised: 28 April 2022 ◦ Accepted: 23 May 2022

ABSTRACT

Introduction: Changes and deterioration of the condition of Corona Virus Disease-19 (COVID-19) patients are unknown and can even lead to the death of the patient in the isolation room. This study focuses on the problem of nurses monitoring and supervising the vital signs of COVID-19 patients in the isolation room. This study aimed to identify nurses' problems monitoring and supervising the vital signs of COVID-19 patients in the isolation room.

Methods: A descriptive qualitative study was conducted in January 2022 at a private hospital in Sukoharjo, Central Java. Ten participants who met the inclusion criteria were involved through purposive sampling. The semi-structured interview guidelines were developed by the researcher and they used them to obtain the data. Interviews were recorded using a recorder device. The data from the interviews was recorded verbatim and analysed using inductive content analysis following Elo and Kyngas's method. The member checking and thick description were done to maintain the validity of the data.

Results: Three main themes were found from the emerging data, including: 1) the comparison of nurses and patients is not ideal with patients' conditions that cannot be predicted, 2) personal protective equipment (PPE) and monitoring tools have not reached the appropriateness target, and 3) the demand for high-technology vital sign monitoring devices was not feasible.

Conclusions: Health The problems being faced by the nurses in monitoring and supervising the vital signs of COVID-19 patients in the isolation room need innovative tools. Therefore, it is important for policymakers and related parties to create high-technology solutions for the isolation rooms to improve nursing service effectively and ensure the safety of both the patients and the nurses.

Keywords: COVID-19 patients, isolation room, monitoring, supervision, vital signs.

Introduction

Corona virus disease-19 (COVID-19), which is also known as SARS-CoV-2, was reported to have emerged in December 2019 in Wuhan, China. This epidemic quickly spread outside China, causing the whole world to experience an emergency response, especially in health services (Nishiura *et al.*, [2020](#)). The disease's

transmission runs very easily through aerosols and droplets, which makes the number of cases continue to increase rapidly (Rahman *et al.*, [2020](#)).

Since the COVID-19 virus has not yet found the medication, the health workers can be effective in caring for the patients by conducting continuous monitoring and identifying the deterioration of the patient's

condition (Pimentel *et al.*, 2020; Sharma *et al.*, 2020). In accordance with the recommendation of the Ministry of Health of Indonesia in 2020, patients with confirmed COVID-19 from moderate to severe degrees must be admitted to a hospital for isolation and periodic monitoring, including monitoring of general conditions, vital signs, and laboratory check-up (Indonesia Ministry of Health, 2020). Those monitored were pivotal to predict the patient's conditions and detect the changes in health outcomes during the isolation phase in COVID-19 patients (Connor and Pilar Camargo Plazas, 2021).

One of the things being monitored for the COVID-19 patients is their vital signs. Therefore, the nurses who work in the isolation room should catch up on how to monitor the patients and report the patients' health outcomes (Yuki and Sophia, 2020). The assessment of vital signs gives the signs that the body is functioning in an acceptable condition or has health problems. Moreover, the monitoring of the vital signs also provides information to the health care providers in support of the objective data in the nursing process. Indeed, COVID-19 patients with severe conditions requiring advance monitoring can be seen every one to two hours (Noviestari *et al.*, 2020; Yuki and Sophia, 2020).

Previous research conducted by Rao *et al.* (2021) and Joo and Liu (2021) has explored the experiences, challenges, and obstacles of various health care providers in treating COVID-19 patients from different treatment rooms. They found in advance that caring for the COVID-19 patients is challenging practice, insufficient support, concerns about family, and emotional and psychological stress. Otherwise, they felt focusing on infection risk at the expense of high-quality care, struggling with dynamic and unfamiliar challenges, being overwhelmed and exhausted by personal protective equipment (PPE), information overload and confusion, also being overstretched by additional responsibilities at work. The health care facilities to take care of the COVID-19 patients still lack innovation. The assessment and reporting of the data were not digitally recorded. It is evidence that finding the right solution to the problems is difficult.

The nurse's perspective, who has experience caring for COVID-19 patients in the isolation room of the hospital where the study was conducted, said that the flow of COVID-19 patients due to this outbreak made the nursing services hectic. When the pandemic broke out, healthcare providers were not ready for health services and treatment availability. The capacity of the isolation room was for 24 patients, but the condition should be able to handle up to 29 patients. The ironic situation

when the nurses are taking care of the patients is that there are only three or four nurses on duty per shift. Even with the health facilities available in the patient's room, such as oxygen central, digital blood pressure, oxygen saturation, and more, the time to monitor and supervise the patients still lacks. Due to the limited admission in the Intensive Care Unit (ICU), the isolation room was forced to care for patients with severe conditions. In addition, the transmission of the COVID-19 virus requires the nurses to take care of the patients' with personal protective equipment (PPE). The flow of COVID-19 patients in the isolation room made some patients not get maximum monitoring and supervision, so that changes and deterioration of the patient's condition were difficult to detect early (Pimentel *et al.*, 2020).

Based on the researchers' experience, monitoring and supervision of the vital signs of COVID-19 patients in the isolation room is manual and conventionally recorded. In the setting of the hospital, we found that there were 78% of COVID-19 patients admitted to the isolation room. Some of the patients became severely ill and required advanced nursing care in the ICU. However, since in the ICU there is also no bedside, those patients should be taken care of in the isolation room. Therefore we conducted the study to identify nurses' problems in monitoring and supervising COVID-19 patients' vital signs in isolation room.

Materials and Methods

This study was a descriptive qualitative study that describe the problems facing by the nurses in monitoring and supervision the vital signs of COVID-19 patients at the isolation room. This research was conducted in the COVID-19 isolation room of the private hospital in Sukoharjo, Central Java in January 2022. The researchers ensured that the health protocols applied by wearing masks, hand sanitizers, and keeping a distance from participants when research data collection.

The sampling technique was carried out by purposive sampling (Polit and Beck, 2015). Ten nurses who work in the isolation room participated in this study. To encourage the participants, the researchers communicated with the head nurses and head team to ask for the nurses who met the inclusion criteria. Inclusion criteria in this study were: 1) nurses had work experience of at least 6 months, 2) taking care of the COVID-19 patients in isolation room for at least 2 weeks. The exclusion criteria of the study were that if the participants were not available to complete the

interview process due to their duty to take care of the patients. Informed consent was explained to the participants before the interview. The interview guidelines used in this study were developed by the researcher. We constructed the interview guidelines based on the research questions and objectives as well as the research concepts and phenomena. The interview questions consist of: 1) What do you have done in monitoring COVID-19 patients in the isolation room? 2) What does the manager do in supervising COVID-19 patients in the isolation room? 3) What are the obstacles faced by nurses in monitoring vital signs of COVID-19 patients in the isolation room? 4) What are the obstacles faced by nurses in supervising vital signs of COVID-19 patients in the isolation room? 5) Is there any innovation in monitoring and supervision vital signs other than manually?. All the interview processes require 45-60 minutes for each participant.

The results of the interviews were recorded using a tape recorder and were verbatim. The data verbatim was then analyzed to obtain the themes according to the research objectives. Content analysis with an inductive approach followed by the Elo and Kyngas method was used to analyze the research data. The data analysis consists of an open coding process, coding sheets, grouping, categorization, and abstraction (Elo and Kyngäs, 2008). An example of the data analysis process is shown in [Table 1](#).

To ensure trustworthiness, the researchers conducted data validity by conducting a member check to know the participants' validation of the interview results. Transferability was tested by comparing the results of this study with similar studies to explain the concept of the phenomena. To verify the emergent themes from the data, the researchers consulted an

external expert to justify the theoretical and methodological discussion (Korstjens and Moser, 2018).

To guarantee the ethical consideration for this study, the research ethics approval from Sultan Agung Islamic Hospital with the number of Ethical Clearance 37/KEPK-RSISA/XII/2021 was required. The researchers also followed the rules of ethics for human subject guidance based on the Council for International Organizations of Medical Sciences (CIOMS) in collaboration with the World Health Organization (2016). Anonymity, beneficence, justice, and do not harm were practiced by researchers in the data gathering process. The participants could withdraw from this study at anytime if they felt uncomfortable and there were no consequences required.

Results

Characteristic of the Participants

The characteristics of participants include age, sex, education level, work experience, and time to take care of the COVID-19 in the isolation room. All participants were female nurses, with an age range of 26 to 35 years old (80%) mostly. The participants' education background was 80% graduated from Diploma III, and had work experience as nurses for 10 years (50%). The participants had the time to take COVID-19 patients at the isolation room within 4-6 months (50%). [Table 2](#) explains the characteristics and demographics of the participants.

This study found three main themes in nurses' problems in monitoring and supervision of COVID-19 patients' vital signs in the isolation room. They are: 1) the comparison of nurses and patients is not ideal with patients' conditions that cannot be predicted, 2)

Table 1 Depicts an example of the data analysis process

Participant Statement	Coding	Theme
"The imbalance between the number of patients and the nurses' resources makes nurses not optimal in monitoring the vital signs of COVID-19 patients in the isolation room" (P2)	The comparison of unideal patients and nurses	The comparison of nurses and patients is not ideal with patients' conditions that cannot be predicted
"When monitoring the vital signs, there are unexpected activities, such as a patient who suddenly experiences a deterioration in their condition. The unexpected activities often cause the patient's vital signs to not be monitored properly. Nurses must be in a hurry or not be on time to monitor the vital signs of the patients" (P8)	Unpredictable situations and conditions	
I had an experience when I was on my shift. That day, I did not assess the patient's saturation because a device was running out. I have reported it to the head nurse, but that time cannot be fixed promptly as it is a hospital regulation. We suggest the patient's family bring the oximetry to check their own saturation. "(P6)	There is no monitor of vital signs with high technology.	The demand for high- technology vital signs monitoring devices was not feasible
"In the patient room, there are no vital signs devices that are continuously attached to the patient, so we cannot monitor the patient's vital signs directly from the nurse station. The deterioration of the patient's condition somehow delays response." (P10)	The absence of monitoring patients' vital signs with high technology connected to the nurse station	Sub theme 1. The condition of isolated patients requires personal monitoring device Sub theme 2. The monitor devices that are connected to the nurse station directly

Table 2 Participant characteristics

No	Demographic Characteristic	Frequency (f)	Percentage (%)
1	Sex		
	Female	10	100
2	Age (Year)		
	17-25	2	20
	26-35	8	80
3	Education background		
	D3 Nursing Program	8	80
	Certified Nurse Profession	2	20
4	Work Experience (Year)		
	1-5	5	50
	6-10	5	50
5	Time to take care of COVID-19 patients in the isolation room (months)		
	1-3	3	30
	4-6	5	50
	10-12	2	20

personal protective equipment (PPE) and monitoring tools have not reached the appropriateness target, and 3) the demand for high-technology vital sign monitoring devices was not feasible.

Theme 1: The comparison of nurses and patients is not ideal with patients' conditions that cannot be predicted

The participants explained that the comparison of nurses and patients was not ideal under the conditions. Worsening of the patient's condition could not be predicted since the obstacles in monitoring the vital signs of COVID-19 patients in the isolation room. The nurse-patient ratio is not ideal due to the lack of nurses compared to the high rate of admission of COVID-19 patients and the workload of nursing care. This condition is exacerbated by unpredictable conditions such as unexpected activities or a deterioration of the patient's condition. The statement was expressed by four participants as follows:

"The imbalance between the number of patients and the nurses' resources makes nurses not optimal in monitoring the vital signs of COVID-19 patients in the isolation room."(P2)

"When monitoring the vital signs, there are unexpected activities, such as a patient who suddenly experiences a deterioration in their condition. The unexpected activities often cause the patient's vital signs to not be monitored properly. Nurses must be in a hurry or not be on time to monitor the vital signs of the patients."(P8)

"Here, we (the nurses) work in the isolation room based on the managerial setting. The comparison between nurses and patients is definitely not ideal.

Sometimes we feel that what we do in monitoring the patients does not complete due to the limited sources of information between patients and nurses."(P9)

"Nurses should have approximately 30 minutes in the patient room to complete the treatment for a patient. All monitoring and supervision of patients in the isolation room is done on paper. We can imagine what a load of work this is here."(P10)

Theme 2: Personal protective equipment (PPE) and monitoring tools have not reached the appropriateness target

Participants argued that personal protective equipment (PPE) and monitoring devices had not yet reached the appropriate target. Quality standards, such as comfort and protection, should be met by the target PPE. However, the PPE that was provided for them was still low quality, so they felt uncomfortable wearing it. In addition, the supply of the devices for vital signs monitoring was limited compared to the number of patients who should assess their continuous vital signs. Four participants expressed it as follows:

"Because of the quality standard of PPE, we have problems assessing the respiratory rate of COVID-19 patients. If we use it, we feel discomfort and somehow perspire."(P1)

"For monitoring the vital signs of COVID-19 patients in the isolation room, it is constrained by the equipment of vital signs devices that are still lacking."(P3)

"The problem for nurses in monitoring and supervision of COVID-19 patients' vital signs at the isolation room is that the devices provided are still lacking. We use the old instruments that the hospital provided. In the reality of this situation, we need innovation and technology-based monitoring to monitor the patients."(P9)

"The use of PPE makes it difficult for nurses to feel the pulse and see the patient's breathing. This is due to the use of gloves that must be doubled and google glass often condenses, thus limiting nurses in monitoring the vital signs of COVID-19 patients."(P10)

Theme 3: The demand for high- technology vital signs monitoring devices was not feasible.

The need for a high-technology device to monitor the vital signs is not feasible yet in this setting. Two sub-themes emerge from the data, namely, that isolated patients require personal monitoring devices and those monitoring devices that are connected directly to the nurse station.

Sub-theme 1. The condition of isolated patients requires personal monitoring device

Monitoring of the vital signs of COVID-19 patients in the isolation room should follow the regulation that one device per patient. However, the situation might not be accomplished because the devices were often in error, such as running out of battery and out of service. One of the factors that causes frequent battery runs out and device errors is the sterilization process and its too frequent use, considering that these devices must be used alternately for all patients. Digital blood pressure, thermometer, and oximetry instruments, for example, must be calibrated to ensure the feasibility and accuracy of the data measured. The statement was expressed by three participants as follows:

"For example, we (the nurse) should check the digital thermometer and the oximetry before using it. The devices are out of service due to a low battery charge and high utilization for all patients. Therefore, we need to check two or three times before and after assessing the patient's vital signs to record the right data." (P2)

"I had an experience when I was on my shift. That day, I did not assess the patient's saturation because a device was running out. I have reported it to the head nurse, but that time cannot be fixed promptly as it is a hospital regulation. We suggest the patient's family bring the oximetry to check their own saturation." (P6)

"A digital sphygmomanometer actually needs calibration to ensure the actual data recorded is accurate." Nevertheless, the calibration devices are undone." (P9)

Sub-theme 2. The monitor devices that are connected to the nurse station directly

While caring for COVID-19 patients in the isolation room, there is no monitor that can be connected to the nurse station. It is evidence that the nurses find it difficult to do monitoring and supervision if they enter the patient's bedside. The nurse imagines what would happen if the monitoring and supervision of the patients' vital signs were synchronized in a good system, so the vital signs data would remain objective and accurate. Closed-circuit television (CCTV) is not a basis to check a patient's condition to date. The patient's room even has no CCTV, thus some patients are not monitored optimally and there are even delays in knowing the deterioration of the patient's condition. Four participants said the following:

"The patient's bedside is not equipped with CCTV, so the nurses cannot control the patient's mobility around. Thus, we should enter the room and check the patients." (P1)

"Because there is no integrating monitor installed on the patient's bedside continuously, the vital signs of the

patients somehow failed to be recorded on time and to date." (P3)

"The difficulty for nurses in monitoring the vital signs of COVID-19 patients in this isolation room is that there are no vital sign devices that are continuously installed on the patient and connected to the nurse station's monitor." (P8)

"In the patient room, there are no vital signs devices that are continuously attached to the patient, so we cannot monitor the patient's vital signs directly from the nurse station. The deterioration of the patient's condition somehow delays response." (P10)

Discussions

This study aims to identify nurses' problems in monitoring and supervising the vital signs of COVID-19 patients in the isolation room. The results of the study indicate three main themes that indicate the problems underpinning nursing services in the hospital. The need for technology based to measure the vital signs and record them was also present in this study's results. Therefore, the development of the tools of digital or technology-based care for COVID-19 patients cannot be delayed.

The theme of the comparison of nurses and patients who are not ideal with conditions that cannot be predicted, confirmed that the high ratio between the nurse and the patient the ideal one is being one nurse to one patient. They can't provide more responsibilities when monitoring COVID-19 patients who are being treated in the isolation room (González-Gil *et al.*, 2020; Kang and Shin, 2020; Maben and Bridges, 2020). A shortage of nurse resources and an overflow number of patients result in an inability to perform regular and timely monitoring and reduce the quality of care (Tan MN, RN *et al.*, 2020).

The unpredictable situation is also felt to be a problem for nurses in monitoring and supervising the vital signs of COVID-19 patients in the isolation room. This is due to unexpected responsibilities and the ratio between the nurses and the patients was not ideal. For example, if one of the patients experiences a critical condition, the hectic situation is twisted. Patients with COVID-19 have a higher risk of deterioration and severity than sufferers of other viruses, so many patients experience a sudden deterioration. This increases the nurses' workload in monitoring and supervising COVID-19 patients in the isolation room (Pimentel *et al.*, 2020). The nurses' workload resulted in physiological and psychological responses such as anxiety, stress, and difficulty sleeping (Ismail, Ridlo and

Rochana, [2021](#)). Joo and Liu ([2021](#)) argued that unexpected tasks when caring for COVID-19 patients are the obstacles for nurses in monitoring COVID-19 patients in the isolation room. Nurses have to do other work and even complete the duties other health team members (Lee and Lee, [2020](#); Schroeder *et al.*, [2020](#)). Moreover, this pandemic has made the care of COVID-19 patients increase in direct care time, missed treatment time, and waiting time for nurses' duties by 27%, 311%, and 44%, respectively (Qureshi *et al.*, [2021](#)).

The results of the study also stated that PPE and monitoring devices had not reached the appropriateness target. The result of the study was in line with Tallulembang ([2020](#)) statement that the limited supply of PPE, care devices, or facilities were the obstacles for nurses in implementing nursing care (Tallulembang, Widani and Bandur, [2020](#)). Atay and Cura (2020) also agreed that one of the barriers for nurses in monitoring COVID-19 patients was the use of PPE. As the PPE and monitoring devices were necessary in assessing the vital signs of COVID-19 patients, it was necessary for the manager to provide good quality PPE and supply the vital signs tools appropriately (BoSkoski *et al.*, [2020](#)).

The participants explained that the monitoring of vital signs in COVID-19 patients was still done manually by nurses. It is because there were no vital signs monitoring device that was connected to the nurse's station. These results are in line with previous research which stated that technology-based monitoring of vital signs for the COVID-19 in the isolation room can not be provided automatically and remotely. Nurses must remain present with patients if they wish to monitor the patients' vital signs (Giménez-Espert, Prado-Gascó and Soto-Rubio, [2020](#); Lee and Lee, [2020](#)). Based on the results of the study, the need for high technology to monitor vital signs in the isolation room is important to provide. The technology based on monitoring and supervision of the patient's condition can influence the quality of nursing care. It is better for the health care division to develop high-tech medical devices that can be the right solution in caring for COVID-19 patients (Sera *et al.*, [2020](#)).

This research has several strengths, including: 1) the data taken is from the nurses who were taking care of COVID-19 patients directly, so that they can deliver the real information through monitoring and supervision of the vital signs of those kinds of patients; 2) this study focuses on describing the nurses' problems in monitoring and supervision of the COVID-19 patients' vital signs in the isolation room. Meanwhile, the weakness of this study was noted as this is a descriptive

qualitative approach, which means the phenomena or the problem might not have been explored deeply enough yet. The sample took 10 nurses and was conducted in one hospital where the equipment needs more new stock. Moreover, the obstacles faced during the research were those of the interview process, where both researchers and participants had to keep their distance and use masks. Somehow, its situation was not clear to hear. To limit the barriers of communication, we did the clarification after the interview process, which we explained in trustworthiness.

Conclusion

The COVID-19 patients are challenging for nurses now, particularly to monitor and supervise their vital signs while patients are in the isolation room. The findings of this study have illustrated the problems and what the impact on the entire patient's health outcomes is. The health outcomes of COVID-19 patients are to improve their quality of life. The improvement of the quality of life of the patients should be based on the quality of nursing care that is provided by the nursing team and health care providers. The nurses should know how to solve the problems by doing continuous monitoring and supervision in caring for the COVID-19 patients, and the nursing manager should be aware of the nurses' work thoroughly.

This study highlights that there are exists problems for nurses in monitoring and supervising vital signs of COVID-19 patients in isolation rooms, and that the problems might remain the same with other nurses outside the setting. These problems are related to time, devices, and resources. Therefore, it is necessary to develop a high-technology-based vital signs monitoring device to monitor and supervise the vital signs of COVID-19 patients in the isolation room. Moreover, the monitoring can be completed automatically, and the data is recorded in real time. There is definitely no paper-based method to support a green economy and green digitalization.

Funding

None. We do not have any financial support.

Acknowledgments

We are grateful to Indriati Solo Baru hospital and all the nurses who participated in this study.

Conflict of interest

The authors declare no potential conflict of interest, and all authors contribute equally.

References

- Atay, S. and Cura, Ş. Ü. (2020) 'Problems Encountered by Nurses Due to the Use of Personal Protective Equipment During the Coronavirus Pandemic: Results of a Survey'. doi: 10.25270/wmp.2020.10.1216.
- BoSkoski, I. et al. (2020) 'Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID- 19 . The COVID-19 resource centre is hosted on Elsevier Connect , the company ' s public news and information', (January). doi: 10.1016/j.gie.2020.04.048.
- Connor, D. M. and Pilar Camargo Plazas (2021) 'Reflections on vital sign measurement in nursing practice', *Nursing Philosophy*, 22(1), pp. 1–9. doi: 10.1111/nup.12326.
- Council for International Organizations of Medical Sciences (CIOMS) in collaboration with the World Health Organization (2016) *International Ethical Guidelines for Health-related Research Involving Humans, Biomedical Research*. Available at: <https://cioms.ch/wp-content/uploads/2017/01/WEB-CIOMS-EthicalGuidelines.pdf>
- Elo, S. and Kyngäs, H. (2008) 'The qualitative content analysis process', *Journal of Advanced Nursing*, 62(1), pp. 107–115. doi: 10.1111/j.1365-2648.2007.04569.x.
- Giménez-Espert, M. del C., Prado-Gascó, V. and Soto-Rubio, A. (2020) 'Psychosocial Risks, Work Engagement, and Job Satisfaction of Nurses During COVID-19 Pandemic', *Frontiers in Public Health*, 8(November), pp. 1–10. doi: 10.3389/fpubh.2020.566896.
- González-Gil, M. T. et al. (2020) 'Nurses' perceptions and demands regarding COVID-19 care delivery in critical care units and hospital emergency services', *Intensive and Critical Care Nursing*. doi: 10.1016/j.iccn.2020.102966.
- Indonesia Ministry of Health (2020) *Protocol of COVID-19 treatment*. JAKARTA. Available at: <https://infeksiemerging.kemkes.go.id/index.php/protokol-covid-19/protokol-penanganan-covid-19>
- Ismail, S., Ridlo, M. and Rochana, N. (2021) 'A Qualitative Description of Nurses ' Psychological Responses in Caring for COVID-19 Patients : An Indonesian Context Abstract :', pp. 136–142. doi: 10.2174/1874434602115010136.
- Joo, J. Y. and Liu, M. F. (2021a) 'Nurses' barriers to caring for patients with COVID-19: a qualitative systematic review', *International Nursing Review*, 68(2), pp. 202–213. doi: 10.1111/inr.12648.
- Joo, J. Y. and Liu, M. F. (2021b) 'Nurses' barriers to caring for patients with COVID-19: a qualitative systematic review', *International Nursing Review*. doi: 10.1111/inr.12648.
- Kang, Y. and Shin, K. R. (2020) 'COVID-19: Korean nurses' experiences and ongoing tasks for the pandemic's second wave', *International Nursing Review*, 67(4), pp. 445–449. doi: 10.1111/inr.12644.
- Korstjens, I. and Moser, A. (2018) 'Series: Practical guidance to qualitative research. Part 4: Trustworthiness and publishing', *European Journal of General Practice*. doi: 10.1080/13814788.2017.1375092.
- Lee, N. and Lee, H. J. (2020) 'South korean nurses' experiences with patient care at a covid-19-designated hospital: Growth after the frontline battle against an infectious disease pandemic', *International Journal of Environmental Research and Public Health*, 17(23), pp. 1–22. doi: 10.3390/ijerph17239015.
- Maben, J. and Bridges, J. (2020) 'Covid-19: Supporting nurses' psychological and mental health', *Journal of Clinical Nursing*, 29(15–16), pp. 2742–2750. doi: 10.1111/jocn.15307.
- Nishiura et al. (2020) 'The Extent of Transmission of Novel Coronavirus in Wuhan, China, 2020', *Journal of Clinical Medicine*. doi: 10.3390/jcm9020330.
- Noviestari, E. et al. (2020) *Dasar-Dasar Keperawatan, Elseiver Singapura*. p. 1-635
- Pimentel, M. A. F. et al. (2020) 'Trajectories of vital signs in patients with COVID-19', *Resuscitation*. doi: 10.1016/j.resuscitation.2020.09.002.
- Polit and Beck (2015) *Nursing research: Generating and assessing evidence for nursing practice 10th edition*, Wolters Kluwer Health. p. 814.
- Qureshi et al. (2021) *Exploring the Impact of COVID-19 on Nurse Workload and Quality of Care via chapter Computerized Simulation*. Springer International Publishing. 219. https://doi.org/10.1007/978-3-030-74602-5_104
- Rahman, H. S. et al. (2020) 'The transmission modes and sources of COVID-19: A systematic review', *International Journal of Surgery Open*. doi: 10.1016/j.ijso.2020.08.017.
- Rao, H. et al. (2021) 'Frontline interdisciplinary clinician perspectives on caring for patients with COVID-19 : a qualitative study'. doi: 10.1136/bmjopen-2021-048712.
- Schroeder, K. et al. (2020) 'Nursing perspectives on care delivery during the early stages of the covid-19 pandemic: A qualitative study', *International Journal of Nursing Studies Advances*. doi: 10.1016/j.ijnsa.2020.100006.
- Sera, W. et al. (2020) 'Applications of digital technology in COVID-19 pandemic planning and response', *The Lancet Digital Health*, 2(8), pp. e435–e440. doi: 10.1016/S2589-7500(20)30142-4.
- Sharma, S. K. et al. (2020) 'Clinical nursing care guidance for management of patient with COVID-19', *Journal of the Pakistan Medical Association*, pp. S118–S123. doi: 10.5455/JPMA.29.
- Tallulembang, A., Widani, N. L. and Bandur, A. (2020) 'Experience of Nurses in Providing Nursing Care for Covid-19 Patients in DKI Jakarta', *Tallulembang ,Ni Luh Widani ,Agustinus Bandur*, 4(1), pp. 74–91. doi: <https://doi.org/10.31934/mppki.v4i1.1392>.
- Tan MN, RN, N. R. et al. (2020) 'Experiences of clinical first-line nurses treating patients with COVID-19: A qualitative study', *Journal of Nursing Management*, 28(6), pp. 1381–1390. doi: 10.1111/jonm.13095.
- Yuki, M. F. and Sophia (2020) 'COVID-19 pathophysiology: A review', *Clinical Immunology*, 215(April 2020). doi: 10.1016/j.clim.2020.108427.

How to cite this article: Christina, T. Y., Ismail S., Erawati M. (2022). 'A qualitative description of nurses' problems to monitor and supervise the vital signs in covid-19 patients at the isolation room', *Jurnal Ners*, 17(2), pp. 103-109. doi: <http://dx.doi.org/10.20473/jn.v17i2.33638>