

SUPPLEMENTARY MATERIAL/MATERI TAMBAHAN

Indonesian Longitudinal Survey on Mental Health and Social Factors (INDOLUMEN): Early Findings and Protocol

EDO S. JAYA*, EKO HERMANTO, SHABRINA AUDINIA, SHIERLEN OCTAVIA, SALIMA CARTER, & FADHILAH RAMADHANNISA

[1] Psychosis Studies Research Group, Faculty of Psychology, Universitas Indonesia, Depok, Indonesia

Supplementary Table 1. Psychotic Symptoms Distress in Participants Based on Community Assessment of Psychic Experience ($N = 464$)

Variable	Mean ^a	SD ^a	Sample Range ^a
Positive psychotic symptoms	0.72	0.57	0–2.85
Bizarre experiences	0.78	0.68	0–3
Hallucinations	0.82	0.86	0–3
Paranoia	0.87	0.73	0–3
Grandiosity	0.36	0.63	0–3
Magical thinking	0.59	0.76	0–3
Negative psychotic symptoms	1.03	0.66	0–3
Social withdrawal	1.01	0.73	0–3
Affective flattening	0.70	0.75	0–3
Avolition	1.20	0.76	0–3
Depressive symptoms	1.52	0.76	0–3

Note. ^acomputed by averaging the items.

Tabel Tambahan 1. Gejala Distres Psikotik pada Partisipan Berdasarkan *Community Assessment of Psychic Experience* ($N = 464$)

Variabel	Rata-rata ^a	SD ^a	Rentang Sampel ^a
Gejala psikotik positif	0,72	0,57	0–2,85
Pengalaman aneh	0,78	0,68	0–3
Halusinasi	0,82	0,86	0–3
Paranoia	0,87	0,73	0–3
Grandiositas	0,36	0,63	0–3
Pemikiran magis	0,59	0,76	0–3
Gejala psikotik negatif	1,03	0,66	0–3
Penarikan sosial	1,01	0,73	0–3
Affective flattening	0,70	0,75	0–3
Avolition	1,20	0,76	0–3
Gejala depresi	1,52	0,76	0–3

Catatan. ^adihitung dengan rata-rata butir.