Evaluation of Modified Unilateral Nasoplasty in Patient with Post Operation of Unilateral Labioplasty at Malahayati Hospital Banda Aceh: Cohort Study from 2017 until 2019

Devyana Enggar Taslim^a, Muhammad Taufik^a, Muhammad Jailani^{a*}, Syamsul Rizal^a, Mirnasari Amirsyah^a

^aClinical Research Unit Cleft Lip and Palate Center Aceh, Banda Aceh, Indonesia

^bPlastic Reconstructive and Aesthetic Surgery Division Faculty of Medicine Universitas Syiah Kuala Banda Aceh, Indonesia

*Corresponding author: Muhammad Jailani; Plastic Reconstructive and Aesthetic Surgery Division Faculty of Medicine Universitas Syiah Kuala Banda Aceh, Indonesia; Email: jailani@unsyiah.co.id

ARTICLE INFO

Keywords:
Cleft Nasal, Modified
Unilateral Rhinoplasty,
Gender, Age, Outcome

ABSTRACT

Background : Nasal deformity associated with cleft lip has been viewed as one of the most challenging reconstructive problems in rhinoplasty.(1) The common clinical features associated with cleft lip nasal deformity are its lack of symmetry, alar collapse on the affected side, short nasal length, loss of tip definition, obtuse nasal labial angle, and altered columella show among others.(2) The complexity of cleft lip rhinoplasty is demonstrated by the abundance of technique that is available for its correction.(3) In this case we used the technique that modified by Dr. Muhammad Jailani, SpBP-RE in Aceh.

Methods: We conducted a retrospective study involving 139 patients who is done the rhinoplasty unilateral operation at Malahayati Hospital from January 2017 to November 2019. Demographic information was recorded such as ratio between pre – operation and post – operation, gender and age.

Results : Our results has shown the result of comparison between cleft nose before rhinoplasty and cleft nose after rhinoplasty is $\pm 0.26 : \pm 0.58$, and the patients who came to the hospital to do the rhinoplasty unilateral surgery are 81% from 139 patients. Woman (61.2%) experience rhinoplasty more dominant compare to man (38.8%), and the highest age average is between 1 years old until 7 years old (71%)

Conclusions: Improvement in procedure's duration and better positioning of both nasal tip and nostril. Expected improvements in terms of aesthetic and functions were observed but further documentations still needed.

INTRODUCTION

Bair and Brown first described the cleft nose in 1931, critically identifying the nuances of the pathology.(4) The nose and the lips develop from derivatives of these prominences as follows: frontonasal prominence, lateral nasal process, medial nasal process, maxillary prominences, mandibular prominences, mesenchyme in the facial prominences.(5) The most complicated morphogenic movement occurs between the fourth and eight weeks of embryogenesis.(6)

Nasal deformity associated with cleft lip has been viewed as one of the most challenging reconstructive problems in rhinoplasty. The complexity of cleft lip rhinoplasty is demonstrated by the abundance of technique that is available for its correction.(7) The common clinical features associated with cleft lip nasal deformity are its lack of symmetry, alar collapse on the affected side, short nasal length, loss of tip definition, obtuse nasal labial angle, and altered columella show among others.(3) The surgical approaches

for a cleft lip rhinoplasty can be a closed endonasal or an open approach. The open or external approach is indicated in cases where there is a severe deformity of the nasal tip.(8)

Surgical Technique

All the patient who was undergo the open rhinoplasty surgery was carried out under general anesthesia and placed in the supine position. This is a technique used by Dr. M. Jailani Specialist Plastic Surgery and Reconstruction who modified the technique of open reduction by Djohansjah (doctor's dissertation: Prof. Dr. dr. Djohansjah Marzoeqi, SpBP) which uses an incision according to Rethi who opened the columella in both sidewhreas this modification is only done on one side only(9)(10)

During rhinoplasty surgery, we make an incision design at the cleft area and the normal side as the rule. Release all the nose cartilage such as greater alar cartilage, lateral nasal cartilage, lesser alar cartilage and septum cartilage from the skin on the nose cleft side area. After we separate the skin and the cartilage, the excess skin at this part is removed and the cartilage fixed on the upper side of the nostril area once believed to be nearing the normal point.

Fig. 1- 2 Make an incision design on the cleft nose area with the normal nose as a rule.

Fig. 3-6 Release all the nose cartilage such as greater alar cartilage, lateral nasal cartilage, lesser alar cartilage and septum cartilage from the skin on the nose cleft side area.

Fig. 7-12 After we separate the skin and the cartilage, the excess skin at this part is removed to get the ideal shape of the nostril which is then sewn to the upper nostril area.

On the columella septum section, fixation of the inner nasal area between the cleft and the non-cleft area (transnasal sutures) (Fig. 13) to create the nasal dome for better symmetry and projection (raise the cartilage / repositioning the cartilage) to a rather normal position.(11) It sewn using absorbable thread Proline 6-0 and eventually make the septum cartilage stand straight.

Fig. 13 Transnasal sutures to create the nasal dome for better symmetry and projection, and also make the septum cartilage stand straight to support nasal dome and nasal tip at the center.

METHOD

We conducted a retrospective study involving 139 patients who is done the rhinoplasty unilateral operation at Malahayati Hospital from January 2017 to November 2019. Demographic information was recorded such as ratio between pre – operation and post – operation, gender and age.

RESULT

Fig. 14 Diagram of Total Patient of Rhinoplasty Surgery

Fig. 15. Diagram of Comparison Between Male and Female

Fig. 16. Diagram of Age

Fig. 17. Average Between Before and After Operation of Normal Side and Cleft side 2017

Fig. 18. Average Between Before and After Operation of Normal Side and Cleft side 2018

Fig. 19. Average Between Before and After Operation of Normal Side and Cleft side 2019

From 2017 to 2019, out 172 patients who registered for unilateral rhinoplasty procedure, 81% were followed up. The gender ratio is 0.38 to 0.61 with female predominates. The age group incidence highest for age 1 – 7 years old with 71% from the total samples. Post procedural observations showed slight improvement at normal side 0.060 : 0.66 and 0.26 : 0.58 on the cleft side.

DISCUSSION

A deformed nose that results from unilateral cleft of the lip and palate is likened to a tent whose one side depressed.(12) On the cleft side, the orbicularis oris muscle inserts into nasal base, retracting it laterally and inferiorly. The goal of rhinoplasty surgery include creation of nasal symmetry, definition of nasal base and tip, relief of nasal obstruction, and management of nasal scarring.(13) The literature reports various methods with which to assess the cleft lip nasal deformity.(14) According to Kaufman

Fig. 20A-20J. Some pictures of before and after using the modification technique by dr. Muhammad Jailani, SP.BP-RE

et al, an open rhinoplasty can be performed to expose the bilateral cartilage on the lower lateral sides and directly observe the geometric difference. If there is any lateral vestibular webbing accours through an incision, a V-Y or back-cut type incision can

be used to extend the lateral nasal wall, and advance the lower lateral cartilage forward.

This modified technique that introduced by Dr. Muhammad Jailani, SpBP-RE, even without doing the usual technique, the surgical outcome still come out good. This technique can be applied to unilateral cleft nose post - labioplasty procedure. .Improvement in procedure's duration and better positioning of both nasal tip and nostril (pre - rhinoplasty and post - rhinoplasty ratio is $\pm 0.26 : \pm 0.58$) and expected improvements in terms of aesthetic and functions were observed but further documentations still needed.

ETHICAL CLEARANCE

The authors declare that ethics approval was not required for this case report

CONFLICT OF INTEREST

The authors have no conflicts of interest to disclose.

FUNDING

No financial support was received for this case report.

AUTHOR CONTRIBUTIONS

All authors drafted the manuscript. All authors listed have made substantial, direct, and intellectual contribution to the work and approved the final manuscript.

REFERENCES

1. Baskaran M, Packiaraj I, Arularasan SG, Divakar TK. Cleft rhinoplasty. J Pharm Bioallied Sci [Internet]. 2015 Aug;7(Suppl 2):S691-4. Available from: <https://www.ncbi.nlm.nih.gov/pubmed/26538945>
2. Tiong WHC, Zain MAM, Basiron NH. Augmentation Rhinoplasty in Cleft Lip Nasal Deformity: Preliminary Patients' Perspective. Plast Surg Int. 5 2014;2014:1-7.
3. Van Beek AL, Hatfield AS, Schnepf E. Cleft Rhinoplasty. Plast Reconstr Surg. 2004;114(4):57e-69e.
4. Blair VP, Brown JB. Nasal abnormalities, fancied and real. Int J Orthod Oral Surg Radiogr [Internet]. 1932;18(4):363-401. Available from:

- [http://dx.doi.org/10.1016/S0099-6963\(32\)80076-1](http://dx.doi.org/10.1016/S0099-6963(32)80076-1)
5. Wantia N, Rettinger G. The current understanding of cleft lip malformations. *Facial Plast Surg.* 2002;18(3):147–53.
 6. Clark JM, Skoner JM, Wang TD. Repair of the Unilateral Cleft Lip / Nose Deformity.
 7. Cho BC, Baik BS. Correction of cleft lip nasal deformity in Orientals using a refined reverse- U incision and V-Y plasty. *Br J Plast Surg* [Internet]. 2001 Oct 1 [cited 2020 Jan 29];54(7):588–96. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0007122601936823>
 8. Harold MC. Treatment Of Unilateral Cleft Lip Nose.
 9. Marzoeki Djohansjah ; Muhammad Jailani ; David S. Perdanakusuma. *Tehnik Pembedahan Celah Bibir Dan Langit-langit. Teh Pembedahan Celah Bibir Dan Langit-langit.* 2002
 10. Marzoeki Djohansjah; Repositioning Of The Nose in Primary Unilateral Cleft Lips Closure. 1989;
 11. Kaufman Y, Buchanan E, Wolfswinkel E, Weathers W, Stal S. Cleft nasal deformity and rhinoplasty. *Semin Plast Surg.* 2012;26(4):184–90.
 12. Shih CW, Sykes JM. Correction of the cleft-lip nasal deformity. *Facial Plast Surg.* 2002;18(4):253–62.
 13. Deshpande G, Mehrotra V, Natarajan S, Merchant R. Cleft Rhinoplasty: A Surgical Technique. *J Contemp Dent.* 2017;7(3):188–93.
 14. Surgery M, Unit O. Methods of assessment of cleft-related facial deformity: a review. 2003;