

Primary Open-Angle Glaucoma Patients in the Ophthalmology Outpatient Clinic Dr. Soetomo General Academic Hospital, Surabaya, in 2013–2015

Risnanda Putri Rasyda¹, Indri Wahyuni^{2*}, Dewi Ratna Sari³

¹Faculty of Medicine, Universitas Airlangga, Surabaya, Indonesia.

²Department of Ophthalmology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Academic Hospital, Surabaya, Indonesia.

³Department of Histology, Faculty of Medicine, Universitas Airlangga/Dr. Soetomo General Academic Hospital, Surabaya, Indonesia.

ABSTRACT

Introduction: Primary open-angle glaucoma (POAG) is a progressive, chronic optic neuropathy characterized by distinct optic nerve atrophy and impairment of the visual field. The purpose of this study was to describe the characteristics of primary open-angle glaucoma patients in the ophthalmology outpatient clinic of Dr. Soetomo General Academic Hospital, Surabaya, in 2013–2015.

Methods: This was a descriptive-retrospective study. Samples were patients with a primary open-angle glaucoma (POAG) diagnosis in the ophthalmology outpatient clinic at Dr. Soetomo General Academic Hospital, Surabaya, in 2013–2015.

Results: There were 88 samples of POAG patients in the ophthalmology outpatient clinic at Dr. Soetomo General Academic Hospital, Surabaya, in 2013–2015. POAG was dominated in the age group 60–69 years old, with a percentage of 37.50%. Males were more dominant than females, with a percentage of 55.70%. Many POAG patients were unemployed, with a percentage of 56.82%. Most patients with POAG did not have a history of hypertension or diabetes mellitus, with a percentage of 73.86%. Patients who had a history of hypertension were 11.36%, those with diabetes mellitus were 9.09%, and those who had a history of both hypertension and diabetes mellitus were 5.68%.

Conclusion: POAG was a higher risk in older adults, with males being more dominant than females. Most patients did not have a history of either hypertension or diabetes mellitus.

Highlights:

1. POAG is a common type of glaucoma.
2. POAG patients were predominantly >60 years old, with males being more dominant than females.

ARTICLE INFO

Article history:

Received 27-10-2022

Received in revised form
29-11-2022

Accepted 1-12-2022

Available online 10-01-2024

Keywords:

Age,
Gender,
Human & Disease,
Occupation,
Primary open-angle glaucoma
(POAG).

Cite this as:

Rasyda RP, Wahyuni I, Sari DR. Primary Open-Angle Glaucoma Patients in the Ophthalmology Outpatient Clinic Dr. Soetomo General Academic Hospital, Surabaya, in 2013–2015. *JUXTA J Ilm Mhs Kedokt Univ Airlangga* 2024; 15: 41–43.

* Correspondence: indri-w@fk.unair.ac.id

JUXTA: Jurnal Ilmiah Mahasiswa Kedokteran Universitas Airlangga

p-ISSN: 1907-3623; e-ISSN: 2684-9453

DOI: 10.20473/juxta.V15I12024.41-43

Open access under Creative Commons Attribution-ShareAlike 4.0 International License
(CC-BY-SA)

Introduction

There are several types of glaucoma, namely glaucoma with primary and secondary open angles, glaucoma with primary and secondary closed angles, and congenital glaucoma.¹ Primary open-angle glaucoma (POAG) can be an optic neuropathy characterized by degeneration of retinal ganglion cells and elevated intraocular pressure (IOP). POAG can remain asymptomatic until it becomes severe. Therefore, many people lack awareness of POAG. Most studies reported an association between age and gender with POAG.

Over 60 million people worldwide are estimated with glaucomatous optic neuropathy, of which 8.4 million are blind.² POAG is a common type of glaucoma worldwide. Hence, it becomes the second leading cause of irreversible blindness after cataracts.³ Age is one of the main risk factors for POAG, followed by gender, occupation, and systemic disease.⁴ This study aimed to describe the characteristics of POAG patients in the ophthalmology outpatient clinic Dr. Soetomo General Academic Hospital, Surabaya, in 2013–2015.

Methods

This was a retrospective descriptive design study. This study was conducted at the ophthalmology outpatient clinic Dr. Soetomo General Academic Hospital, Surabaya, from September 2013–2015. The population in this study was medical records of all patients with a diagnosis of glaucoma in the ophthalmology outpatient clinic Dr. Soetomo General Academic Hospital, Surabaya. The sample in this study was medical records of patients diagnosed by POAG in the ophthalmology outpatient clinic Dr. Soetomo General Academic Hospital, Surabaya. The sample was selected using the total sampling method. The variables in this study included age, gender, and occupation.

The inclusion criteria in this study were patients older than 40 with complete medical records data. The exclusion criteria in this study were medical records with incomplete data. Data were analyzed descriptively and presented in tables.

Results

Based on the results of the study, the number of patients with POAG in the ophthalmology outpatient clinic Dr. Soetomo General Academic Hospital, Surabaya, in 2013–2015 was 107 patients. 19 medical records were excluded due to incomplete data. Thus, the number of subjects in this study was 88 patients. The distribution of age, gender, and occupation of POAG are shown in [Table 1](#). It shows that most of the patients were in the age group 60–69 (37.50%) and were dominated by males. According to the occupation of the patients, most (56.82%) patients were unemployment, followed by private employee (17.05%) and government employee (13.64%).

Table 1. Distribution of age, gender, and occupation in POAG patients

Variable	Frequency (f = 88)	Percentage (%)
Age (years old)		
40–49	12	13.63
50–59	29	32.95
60–69	33	37.50
70–79	13	14.77
≥80	1	1.13
Gender		
Female	49	44.30
Male	39	55.70
Occupation		
Unemployment	50	56.82
Government employees	12	13.64
Private employees	15	17.05
Entrepreneur	5	5.68
Farmer	5	5.68
Merchant	1	1.14

Source: Research data, processed

Discussion

A study by Vijaya, *et al.* (2014) found a risk that increased four times in the age group older than 60 years old compared to the age of 40–60 years old.⁵ The results are in line with this study. Most of the patients were in the age group 60–69 years old (37.50%). A study by Riva, *et al.* (2018) stated that older age is associated with more severe visual function loss.⁶ It is due to POAG being a chronic and asymptomatic disease.⁶ In this study, males (55.70%) dominated female patients. The results of this study are similar to a study by Musch, *et al.* (2012), which found cases of POAG dominated by males, namely 303 patients (61.90%) of a total of 505 patients.⁷ In contrast, Sun, *et al.* (2011) found cases of POAG dominated by females (55.04%).⁸ Vajaranant, *et al.* (2010) stated that the possibility of females developing POAG is greater than males because there is a relationship between women with sex hormones which can provide an effect on protecting the optic nerve.⁹

Most POAG patients were unemployed. The results of this study are in line with the study by Kim, *et al.* (2016), where a total of 710 POAG patients were found, 329 of whom were unemployed (40.8%).¹⁰ The underlying reason is the lack of awareness of this disease. Patients tend to ignore the initial signs obtained from POAG and turn up in a condition of high IOP.⁵

Strength and Limitations

This study showed the characteristics of POAG patients. It can be used as a comparison for future POAG studies. The limitation of this study was the lack of data on patients' medical records.

Conclusion

POAG was a higher risk in older age, with males being more dominant than females. Most of POAG patients were unemployed.

Acknowledgments

We would like to express our gratitude to the entire staff for their dedication and support, which enabled this research to be conducted.

Conflict of Interest

The authors declared there is no conflict of interest.

Funding

This study did not receive any funding.

Ethical Clearance

This study had received ethical clearance from the Ethical Committee of Dr. Soetomo General Academic Hospital, Surabaya (No. 0055/KEPK/II/2018).

Authors' Contributions

Designing the study and creating the manuscript: RPR. Collecting data: RPR. Performing data result: RPR. Supervising results and discussion: RPR, IW, DRI. All authors contributed and approved the final version of the manuscript.

References

1. Kwon YH, Fingert JH, Kuehn MH, *et al.* Primary Open-Angle Glaucoma. *N Engl J Med* 2009; 360: 1113–1124. [PubMed]
2. Weinreb RN, Aung T, Medeiros FA. The Pathophysiology and Treatment of Glaucoma: A Review. *JAMA* 2014; 311: 1901–1911. [PubMed]
3. Davis BM, Crawley L, Pahlitzsch M, *et al.* Glaucoma: The Retina and Beyond. *Acta Neuropathol* 2016; 132: 807–826. [PubMed]
4. He J, Zou H, Lee RK, *et al.* Prevalence and Risk Factors of Primary Open-Angle Glaucoma in a City of Eastern China: A Population-Based Study in Pudong New District, Shanghai. *BMC Ophthalmol* 2015; 15: 134. [PubMed]
5. Vijaya L, Rashima A, Panday M, *et al.* Predictors for Incidence of Primary Open-Angle Glaucoma in a South Indian Population: The Chennai Eye Disease Incidence Study. *Ophthalmology* 2014; 121: 1370–1376. [PubMed]
6. Riva I, Legramandi L, Katsanos A, *et al.* Influence of Sociodemographic Factors on Disease Characteristics and Vision-Related Quality of Life in Primary Open-Angle Glaucoma Patients: The Italian Primary Open Angle Glaucoma Study (IPOAGS). *J Glaucoma* 2018; 27: 776–784. [PubMed]
7. Musch DC, Shimizu T, Niziol LM, *et al.* Clinical Characteristics of Newly Diagnosed Primary, Pigmentary and Pseudoexfoliative Open-Angle Glaucoma in the Collaborative Initial Glaucoma Treatment Study. *Br J Ophthalmol* 2012; 96: 1180–1184. [PubMed]
8. Sun J, Zhou X, Kang Y, *et al.* Prevalence and Risk Factors for Primary Open-Angle Glaucoma in a Rural Northeast China Population: A Population-Based Survey in Bin County, Harbin. *Eye (Lond)* 2012; 26: 283–291. [PubMed]
9. Vajaranant TS, Nayak S, Wilensky JT, *et al.* Gender and Glaucoma: What We Know and What We Need to Know. *Curr Opin Ophthalmol* 2010; 21: 91–99. [PubMed]
10. Kim KE, Kim MJ, Park KH, *et al.* Prevalence, Awareness, and Risk Factors of Primary Open-Angle Glaucoma: Korea National Health and Nutrition Examination Survey 2008-2011. *Ophthalmology* 2016; 123: 532–541. [PubMed]