



## RELATIONSHIP BETWEEN BODY MASS INDEX AND ACTIVITIES OF DAILY LIVING ELDERLY IN RURAL COMMUNITIES

Joko Susanto<sup>1\*</sup>, Rukanah<sup>2</sup>, Iswatun<sup>1</sup>

<sup>1</sup>Faculty of Vocational Studies, Universitas Airlangga Surabaya, Indonesia

<sup>2</sup>Independent Midwifery Academy Gresik, Indonesia

Research Report

### ABSTRACT

**Introduction:** A high Body Mass Index (BMI) is known to be associated with mortality and morbidity in the elderly. However, other studies also show the opposite results regarding high BMI as a protector against Activity Daily Living (ADL). Accurate quantification of the role of BMI in the incidence of disability in ADL. Given the increasing prevalence of overweight and obese individuals and the long life spans in the population. This study aims to determine the relationship between Body Mass Index and Daily Life Activities in population-based elderly people. **Methods:** The method used in this research is a cross-sectional observational study conducted on 167 respondents from rural areas in the Lamongan Regency, using a consecutive sampling approach. BMI scores and the Barthel Index were recorded among respondents. Data were analyzed using the Spearman Test with a significance level of  $p < 0.05$ . **Results:** A total of 167 respondents came from rural areas in Lamongan Regency, dominated by 96 (57.5%) women and 71 (42.5%) men. The Spearman Test results found a weak and significant positive correlation between body weight and daily activities ( $r = 0.167$ ;  $p < 0.05$ ). Apart from that, a significant weak positive correlation was also found between BMI and ADL ( $r = 0.157$ ;  $p < 0.05$ ). **Conclusions:** The results of the study show that increasing the BMI value has a protective effect on ADL, so it can be concluded that higher body weight and BMI are associated with increasing Barthel ADL scores.

### ARTICLE INFO

Received March 12, 2024

Accepted May 17, 2024

Online May 30, 2024

\*Correspondence:

Joko Susanto

\*Email:

[jsusanto@vokasi.unair.ac.id](mailto:jsusanto@vokasi.unair.ac.id)

**Keywords:**

Activity Daily Living, Body Mass Index, Elderly

### INTRODUCTION

Body Mass Index (BMI) is used as a useful population-level measure of overweight and obesity (Estrella-Castillo & Gomez-de-Regil, 2019; Nuttall, 2015). Overweight or obesity may cause many chronic illnesses. Furthermore, several studies have shown that high BMI is associated with mortality and morbidity among the elderly (Ozturk, Egici, Bukhari, & Toprak, 2017; Yue-Bin et al., 2018). Individuals with overweight and obesity are a significant public health problem due to an increased risk of hypertension, dyslipidemia, coronary heart disease, stroke, type 2 diabetes, musculoskeletal disorders, and several types of cancer (Kelishadi, 2014). High BMI also shows associations with impaired physical functioning. Mobility is another important topic regarding seniors. Accurate quantification of the role of BMI in the incidence of disability in Activities of Daily Living (ADL) is necessary given the increasing prevalence of overweight and obese individuals and the long life span in the population (Sampaio et al., 2020).

Mobility difficulties are often the first sign of functional decline and may indicate the need for preventive measures (Freiberger, Sieber, & Kob, 2020). Mobility problems have been reported as a predictor of all-cause mortality, and patients with BMI  $> 30$  kg/m<sup>2</sup> have low scores in the "Time Up to Go" test, which assesses mobility (Donini et al., 2020). Mobility problems, as well as diseases that cause cognitive impairment, are causes of addiction. Weight loss through dieting may be associated with cognitive impairment. Therefore, obesity or being overweight can have a negative impact on the performance of daily activities (ADL) in the elderly (Y. Zhang et al., 2021). It still has many controversies since some studies report that obesity is protective against ADL, but the opposite is reported in some others. Many explanations still can be debated about those issues (Donini et al., 2020).

Barthel index is a simple index of independence, which is used to score the patients' ability to perform ADLs (Yi et al., 2020). The Barthel index comprises 10 items,



including the presence or absence of fecal and urinary incontinence and the need for assistance with grooming, toilet use, feeding, transfers (e.g., from the chair to bed), walking, dressing, climbing stairs, and bathing (Y. Zhang et al., 2021). Based on those mentioned above, this study aims to investigate the association between BMI and ADL in the population-based elderly population of Lamongan Regency.

**MATERIALS AND METHODS**

The research was conducted in a rural area in Lamongan Regency with a cross-sectional observational study of 286 populations conducted on 167 respondents, using a consecutive sampling approach. Measurement of Body Mass Index (BMI) on physical examination, body weight, and height were measured by trained medical personnel using standard protocols. Body weight was measured to the nearest 1 kg for individuals wearing light clothing. Height is measured by observing the patient's height from the base of the feet to the highest point of the head when standing in a straight position against the wall. BMI is measured using the formula a person's weight in kilograms divided by their height in meters squared. BMI is divided into 4 categories according to WHO guidelines: underweight (BMI <18.5), normal weight (BMI 18.5 to <24.9), overweight (BMI 25.0 to <29.9), and obesity (BMI >30.0)(Yue-Bin et al., 2018).

Measurement of Daily Living (ADL) activities uses the interview method with the Barthel score. The score consists of 10 variables regarding performance in daily life such as the presence or absence of fecal incontinence, urinary continence, assistance required in care, use of the toilet, eating, transfer (e.g. from chair to bed), mobility, dressing, and climbing stairs, and shower. Each variable has a score of 0-2, except for the bathing and grooming variables which have a score of 0-1, and mobility and moving have a score of 0-3. Higher scores indicate a higher level of independence. The Barthel index score is classified as follows: a) 0-4 points: total dependence; b) 5-8 points: severe dependence; c) 9-11 points: moderate dependence; d) 12-19 points: mild dependence; and e) 20 points: total independence. Data were analyzed in percentage terms, then Spearman tested with a significance level of  $p < 0.05$ .

**RESULTS**

**Table 1.** Characteristic of respondent by Gender, BMI, and ADL (n = 167)

Characteristic of respondent	n	%
Gender		
Male	71	42,5
Female	96	57,5
BMI (kg/m2)		
Underweight	64	38,3
Normal Weight	77	46,1
Overweight	21	12,6
Obese	5	3,0
ADL Barthel (score)		
Moderate dependence (9-11)	2	1,2
Mild dependence (12-19)	39	23,4
<b>Total dependence (20)</b>	<b>126</b>	<b>75,4</b>

BMI=body mass index; ADL=activity daily living

A total of 167 elderly from rural areas of Lamongan regencies were enrolled in this study. About 71 (42.5%) respondents were males and 96 (57,5%) of them were females. Most of the respondents were normal (46,1%), followed by underweight (38,3%), overweight (12,6%), and obese (3,0%) in BMI. Based on the ADL Barthel Index assessment, most of them were total independence (75,4%), followed by mild dependence (23,4%), and moderate dependence (1,2%).

**Table 2.** Spearman correlation test between weight BMI and ADL Barthel (n = 167)

Parameter	r	p
ADL Barthel		
Weight (kg)	0,167	0,029
BMI (kg/m2)	0,157	0,043

BMI: body mass index; ADL: activity daily living; \*p-value: statistically significant if less than 0,05

The results of analysis using the Spearman correlation test found a significant weak positive correlation between body weight and ADL ( $r = 0,167$ ;  $p = 0,029$ ), while between BMI and ADL ( $r = 0,157$ ;  $p = 0,043$ ).

**DISCUSSION**

The elderly population in Indonesia was projected at about 26.66 million (9,03%) in 2017. Predicted 27,08 million in 2020, 33,69 million in 2025, 40,95 million in 2030 and 48,19 million in 2035 (Keating, 2022). According to the World Health Organization (WHO), 71,4 years (males: 69,1 years; females: 73,7 years) is the average life expectancy at birth, and the life expectancy over the age of 60 years was 20,4 years (males: 18,9 years; females: 21,7 years) among the global population in 2015 (Robine, Jagger, Crimmins, Saito, & Van Oyen, 2020). Similarly, 57,5% of our study group was female, which is probably the result of the longer life expectancy of women.

The research results found a positive correlation between BMI and Barthel's ADL. This is supported by previous research that obesity is protective against ADL (Chen, Cui, Zhang, & Peng, 2020). Similarly, we obtained controversial results for whether overweight increases ADL disability or has protective effects (Kivimaki et al., 2022). This is because the ability to perform ADL changes not only based on body weight but also eating habits, and eating habits in individuals who are overweight or obese can reduce the risk of malnutrition and can protect against a decrease in ADL ability, considering that nutritional status is related to ADL performance and ability in elderly (Y. Zhang et al., 2021). High waist circumference (WC) is not associated with poor physical performance and ADL disability (Andreacchi et al., 2021). The combination of a higher level of BMI and lower level of WC has been found to protect from mortality in older adults, they also had a decreased likelihood of having ADL disability (F.-L. Zhang et al., 2021).

But another study found an opposite correlation, e.g. Lv YB et al. concluded that higher BMI was associated with a lower risk of disability in ADL among Chinese adults aged 80 years or older, which suggests that current recommendations for the BMI may need to be revisited (Momin et al., 2020). More attention should be paid to underweight, rather than overweight and obesity for the prevention of disability in ADL after age 80 years. Being overweight or obese is associated with a higher risk of death and disabling pathological problems before the age of 80 years (Kivimaki et al., 2022). Other research has also found that higher odds of disability among obese individuals in high-income countries may mean living longer with disability (Salinas-Rodriguez, Rivera-Almaraz, Scott, & Manrique-Espinoza, 2020). The same thing also happens in several countries with higher incomes. Obesity is an independent indicator and does not include chronic conditions (Piqueras et al., 2021). Overall, the inverse association between BMI and disability in ADL is likely multifactorial. Malnutrition, frailty, survival bias or selection bias, and reverse causation were considered potential explanations (Gill, Han, Gahbauer, Leo-Summers, & Murphy, 2020).

Many studies have shown that sarcopenia which is one of the main problems in the elderly, is associated with a decline in muscle mass and strength and is a predictor of poor outcomes, including mortality, disability, and poor quality of life (Nawawi, Justine, & Mazzuin Razali, 2020). Malnutrition and weight loss are the causes of sarcopenia which is associated with functional dependence in the

elderly. Accordingly, in our study, when the BMI decreased, the ADL-performing ability decreased, possibly because of the slowing down of activity due to malnutrition or sarcopenia (Y. Zhang et al., 2021).

## CONCLUSIONS

The studies show increased BMI values as a protective effect of Barthel ADL, so be concluded higher body weight and BMI are associated with increased Barthel ADL scores. In general, healthcare professionals can advise all patients to receive adequate nutrition and perform exercises to get an ideal BMI, which will protect them from most chronic diseases and help them with healthy aging.

## REFERENCES

- Andreacchi, A. T., Griffith, L. E., Guindon, G. E., Mayhew, A., Bassim, C., Pigeys, M., ... Anderson, L. N. (2021). Body mass index, waist circumference, waist-to-hip ratio, and body fat in relation to health care use in the Canadian Longitudinal Study on Aging. *International Journal of Obesity*, 45(3), 666–676. <https://doi.org/10.1038/s41366-020-00731-z>
- Chen, X., Cui, J., Zhang, Y., & Peng, W. (2020). The association between BMI and health-related physical fitness among Chinese college students: a cross-sectional study. *BMC Public Health*, 20(1), 444. <https://doi.org/10.1186/s12889-020-08517-8>
- Donini, L. M., Rosano, A., Di Lazzaro, L., Lubrano, C., Carbonelli, M., Pinto, A., ... Siervo, M. (2020). Impact of Disability, Psychological Status, and Comorbidity on Health-Related Quality of Life Perceived by Subjects with Obesity. *Obesity Facts*, 13(2), 191–200. <https://doi.org/10.1159/000506079>
- Estrella-Castillo, D. F., & Gómez-de-Regil, L. (2019). Comparison of body mass index range criteria and their association with cognition, functioning, and depression: a cross-sectional study in Mexican older adults. *BMC Geriatrics*, 19(1), 339. <https://doi.org/10.1186/s12877-019-1363-0>
- Freiberger, E., Sieber, C. C., & Kob, R. (2020). Mobility in Older Community-Dwelling Persons: A Narrative Review. *Frontiers in Physiology*, 11. <https://doi.org/10.3389/fphys.2020.00881>
- Gill, T. M., Han, L., Gahbauer, E. A., Leo-Summers, L., & Murphy, T. E. (2020).

- Risk Factors and Precipitants of Severe Disability Among Community-Living Older Persons. *JAMA Network Open*, 3(6), e206021. <https://doi.org/10.1001/jamanetworkopen.2020.6021>
- Keating, N. (2022). A research framework for the United Nations Decade of Healthy Ageing (2021–2030). *European Journal of Ageing*, 19(3), 775–787. <https://doi.org/10.1007/s10433-021-00679-7>
- Kelishadi, R. (2014). Health impacts of Obesity. *Pakistan Journal of Medical Sciences*, 31(1). <https://doi.org/10.12669/pjms.311.7033>
- Kivimaki, M., Strandberg, T., Pentti, J., Nyberg, S. T., Frank, P., Jokela, M., ... Ferrie, J. E. (2022). Body-mass index and risk of obesity-related complex multimorbidity: an observational multicohort study. *The Lancet Diabetes & Endocrinology*, 10(4), 253–263. [https://doi.org/10.1016/S2213-8587\(22\)00033-X](https://doi.org/10.1016/S2213-8587(22)00033-X)
- Momin, M., Fan, F., Li, J., Jia, J., Zhang, L., Zhang, Y., & Huo, Y. (2020). Joint Effects of Body Mass Index and Waist Circumference on the Incidence of Hypertension in a Community-Based Chinese Population. *Obesity Facts*, 13(2), 245–255. <https://doi.org/10.1159/000506689>
- Nawawi, A., Justine, M., & Mazzuin Razali, R. (2020). Quality of life, hospitalization, and sarcopenia among the elderly: a systematic review. *Journal of Gerontology and Geriatrics*, 69(1), 45–52. <https://doi.org/10.36150/2499-6564-340>
- Nuttall, F. Q. (2015). Body Mass Index. *Nutrition Today*, 50(3), 117–128. <https://doi.org/10.1097/NT.00000000000000092>
- Ozturk, G. Z., Egici, M. T., Bukhari, M. H., & Toprak, D. (2017). Association between body mass index and activities of daily living in homecare patients. *Pakistan Journal of Medical Sciences*, 33(6). <https://doi.org/10.12669/pjms.336.13748>
- Piqueras, P., Ballester, A., Durá-Gil, J. V., Martínez-Hervas, S., Redón, J., & Real, J. T. (2021). Anthropometric Indicators as a Tool for Diagnosis of Obesity and Other Health Risk Factors: A Literature Review. *Frontiers in Psychology*, 12. <https://doi.org/10.3389/fpsyg.2021.631179>
- Robine, J.-M., Jagger, C., Crimmins, E. M., Saito, Y., & Van Oyen, H. (2020). Trends in Health Expectancies. [https://doi.org/10.1007/978-3-030-37668-0\\_2](https://doi.org/10.1007/978-3-030-37668-0_2)
- Salinas-Rodríguez, A., Rivera-Almaraz, A., Scott, A., & Manrique-Espinoza, B. (2020). Severity Levels of Disability Among Older Adults in Low- and Middle-Income Countries: Results From the Study on Global Ageing and Adult Health (SAGE). *Frontiers in Medicine*, 7. <https://doi.org/10.3389/fmed.2020.562963>
- Sampaio, A., Marques-Aleixo, I., Seabra, A., Mota, J., Marques, E., & Carvalho, J. (2020). Physical fitness in institutionalized older adults with dementia: association with cognition, functional capacity and quality of life. *Aging Clinical and Experimental Research*, 32(11), 2329–2338. <https://doi.org/10.1007/s40520-019-01445-7>
- Yi, Y., Ding, L., Wen, H., Wu, J., Makimoto, K., & Liao, X. (2020). Is the Barthel Index Suitable for Assessing Activities of Daily Living in Patients With Dementia? *Frontiers in Psychiatry*, 11. <https://doi.org/10.3389/fpsyg.2020.00282>
- Yue-Bin, L., Yuan, J.-Q., Mao, C., Gao, X., Yin, Z.-X., Kraus, V. B., ... Shi, X.-M. (2018). Association of Body Mass Index With Disability in Activities of Daily Living Among Chinese Adults 80 Years of Age or Older. *JAMA Network Open*, 1(5), e181915. <https://doi.org/10.1001/jamanetworkopen.2018.1915>
- Zhang, F.-L., Ren, J.-X., Zhang, P., Jin, H., Qu, Y., Yu, Y., ... Yang, Y. (2021). Strong Association of Waist Circumference (WC), Body Mass Index (BMI), Waist-to-Height Ratio (WHtR), and Waist-to-Hip Ratio (WHR) with Diabetes: A Population-Based Cross-Sectional Study in Jilin Province, China. *Journal of Diabetes Research*, 2021, 1–9. <https://doi.org/10.1155/2021/8812431>
- Zhang, Y., Xiong, Y., Yu, Q., Shen, S., Chen, L., & Lei, X. (2021). The activity of daily living (ADL) subgroups and health impairment among Chinese elderly: a latent profile analysis. *BMC Geriatrics*, 21(1), 30. <https://doi.org/10.1186/s12877-020-01986-x>