

Ergothioneine Modulates Biomarkers of Oxidative Stress, Biochemical Profiles and Rectal Temperatures of Arabian Stallions Following Exercise in A Hot-Humid Environment

Adakole S. Adah^{1*}, Joseph O. Ayo², Peter I. Rekwot³, Tagang Aluwong² and Deborah A. Adah⁴

¹Department of Veterinary Physiology and Biochemistry, ⁴Department of Veterinary Medicine University of Ilorin, P. M. B. 1515, Ilorin, Nigeria

²Department of Veterinary Physiology, Ahmadu Bello University, P. M. B. 1044, Samaru- Zaria, Nigeria

³National Animal Production Research Institute, P. M. B. 1096, Shika-Zaria, Nigeria

*E-mail: adakole.as@unilorin.edu.ng

ABSTRACT

These experiments were performed to determine the effect of ergothioneine (EGT) and environmental parameters on some physiological profiles of Arabian stallions following an 1800 m race in a hot-humid environment. Twelve stallions having a mean weight of 401 ± 7.33 kg and aged 5.28 ± 1.14 years were used as subjects. They were divided into two groups of treated stallions ($n = 6$) and untreated stallions ($n = 6$). Group I which was the experimental group was administered with EGT (0.5 mg/kg orally) every week for two months while group II which served as controls was not treated. The temperature and the relative humidity of the experimental site were determined for six days and on the day of the experiment. The temperature-humidity index (THI) was also calculated. Vital parameters and some biochemical parameters of all stallions were determined before the commencement, immediately after, and one hour after the exercise. Some biomarkers of oxidative stress and serum biochemical parameters (aspartate aminotransferase, creatinine and lactate dehydrogenase) of the stallions were also determined. The biochemical parameters were higher ($P < 0.05$) in the untreated group than in the treated group. The results obtained showed that EGT lowered the rectal temperature and modulated biomarkers of oxidative stress and biochemical profiles.

Keywords: biochemical parameters; ergothioneine; heart rate; rectal temperature; respiratory rate

INTRODUCTION

Horses are unique animals when compared with other animals; their respiratory, circulatory systems and thermoregulatory are among the most important systems involved in exercise (Mcmiken *et al.*, 1983; Cottin *et al.*, 2006; Adah *et al.*, 2020). Physiological and biochemical parameters are used to evaluate the responses of horses to exercise following interventions. These interventions include training, periods of rest and dietary supplementation (Mader *et al.*, 2006; Schimdt *et al.*, 2010). Exercise is necessary for health improvement and fitness (Courouce *et al.*, 2000; Padilla *et al.*, 2006) and training status is used to understand how exercise influences physiological and biochemical responses (Marques *et al.*, 2002; Gerard *et al.*, 2014). Biochemical and physiological responses are closely connected to changes in the autonomic nervous system; therefore, they are useful to determine changes under different supplements interventions (Aubert *et al.*, 2003; Hamlin and Hopkins, 2003). The work output of the muscles of horses during exercise depends on physiological capacities to deliver large volumes of blood to the tissue and the splenic reserve supply (Evans and Rose 1988; Mader *et al.*, 2006; Bashir and Rasedee, 2009; Kerley *et al.*, 2018).

Ambient temperatures and relative humidity that are higher than the thermoneutral zone of horses induce

higher metabolic rates associated with decreased evaporative heat losses, impaired thermoregulation, hyperthermia, electrolyte imbalance, and alteration in biochemical parameters (Hodgson *et al.*, 1994; Rensis and Scaramuzzi, 2003; Hartmann *et al.*, 2015). A high ambient temperature and relative humidity impair the performance of a horse due to cardio-respiratory deficiencies, effects on core body temperatures, and changes in biochemical profiles (Schrama *et al.*, 1996; Nenaber *et al.*, 1999).

Ergothioneine (EGT) is a unique supplement because it does not auto-oxidize at physiological pH and does not promote the generation of hydroxyl radicals from H₂O₂ and Fe²⁺ ions (Fenton reaction) (Cheah *et al.*, 2017; Yoshida *et al.*, 2017). Important physiologic features of EGT include its prompt clearance from peripheral circulation into tissues with little metabolism and high stability and its long half-life (approximately 30 days). It has a reduced tendency to auto-oxidize or produces free radicals from peroxide and iron at normal body pH (Kenneth *et al.*, 2004; Cheah *et al.*, 2016). EGT requires the specific carrier protein called organic cation transporter novel type-1 (OCTN-1) to be transported across the cell membrane (Morris *et al.*, 1991; Grigat *et al.*, 2007). OCTN-1 is found mainly in the kidneys, trachea, erythrocytes, lungs, heart, and the bone marrow (Richard *et al.*, 2009; Marone *et al.*, 2016). Contrary to other

antioxidants such as glutathione or N-acetyl-L-cysteine, EGT metabolises slowly and is resistant to disulfide formation (Tamai *et al.*, 1997; Nakamura *et al.*, 2008). The formation of ergothioneine disulfide occurs only at low body pH in the presence of Cu²⁺ or H₂O₂. Ergothioneine acts as a cation chelator (Cheah *et al.*, 2018, Halliwell *et al.*, 2018), has a bioenergetics effect (Tamai *et al.*, 1997), and acts as an immune regulator (Paul and Snyder, 2009). It is also an antioxidant (Cheah *et al.*, 2016; Tang *et al.*, 2018) and is widely distributed within organs and body tissues prone to oxidative stress (Kerley *et al.*, 2018).

The aim of this study is to evaluate the effect of EGT on selected physiological and biochemical profiles following exercise in Arabian stallions in a hot-humid environment.

MATERIAL AND METHODS

Experimental animals

Twelve healthy stallions of the Arabian breed with a mean weight of 401 ± 7.33 kg served as experimental animals. The stallions which were used for pleasure riding were obtained from a standard stable in Ilorin (8° 30' N, 4° 33' E), situated in the Southern Guinea Savannah vegetation zone of Nigeria. The experiment was performed during the hot-humid season characterized by high ambient temperatures and relative humidity. The stallions were housed in a concrete-walled stable covered with

corrugated iron roofing sheet. They were fed with hay and their feed was supplemented with concentrate (groundnut bran). Also, clean and cold water was also provided *ad libitum*. The stallions were screened for gastrointestinal parasites, haemoparasites, and ectoparasites, and only healthy animals were used as subjects. The ethical review committee of the Ahmadu Bello University, Zaria approved and endorsed the research study (ABUCAUC/2017/VET.MED/APP/019).

Determination of thermal environmental parameters

The ambient temperatures were taken at 06.00 h, 12.00 h and 18.00 h for six consecutive days and on the day of the experiment using dry- and wet-bulb thermometers (Brannan, UK). From the data obtained, relative humidity (RH) and temperature-humidity index (THI) at each hour of measurement were determined. The THI was determined using the equation of Hartmann *et al.* 2015, as follows:

$$\text{THI} = (\text{DBT} \times 0.8) + \{(\text{RH}/100 \times (\text{DBT}-14.4) + 46.4)\}$$

Where, DBT = Dry Bulb Temperature and RH = Relative Humidity

Experimental design

The stallions were divided into two groups of six stallions each: Group I stallions were administered with EGT at a dose of 0.5 mg/kg by oral administration every week for two

months before subjecting them to exercise while stallions in Group II serve as control were not administered with EGT before exercise.

Before the commencement of the experiment, the heart rate (HR), respiratory rate (RR) and rectal temperature (RT) of each stallion were determined in all the stallions. The HR of each horse was measured using the Equine Polar Heart Rate monitor (Polar Electro Oy, Kempele, Finland), while RR was determined by counting the number of respiratory flank movements per minute as described by Williams (2016). The RT was determined using a digital clinical thermometer (The Hartman's Company, England). The thermometer was inserted about 5 cm into the rectum of each animal, and the RT value was read after a beep sound was heard, indicating the end of the reading.

Exercise protocol

The stallions were mounted by riders of an average weight of 70.43 ± 3.78 Kg and subjected to a race of 1800 m on a race track. Rectal temperature, heart, and respiratory rates were determined in the stallions pre-exercise, 5 min post-exercise, and 1-hour post-exercise. Immediately after exercise, stallions rested under a tree shade.

Collection of blood samples

A blood sample of 10 ml was collected by jugular venipuncture from each of the stallions pre-exercise and post-exercise. The samples were

collected into plain containers without anticoagulant so as to harvest serum. The blood samples were transported to a physiology research laboratory in a Coleman box for the analyses of biomarkers of oxidative stress and some biochemical parameters.

Determination of biochemical parameters

The serum activities of the enzymes lactate dehydrogenase, aspartate aminotransferase and concentration of creatinine were determined using a clinical chemistry analyzer Kone Pro (Konelab, Thermo Clinical Lab systems Oy, Finland) while the activities of superoxide dismutase, catalase, glutathione peroxidase and concentrations of malondialdehyde were determined using spectrophotometric methods.

Analyses of data

Data generated from the study were expressed as mean \pm SEM and analysed using the Student *t*-test to compare the two groups. Data generated from thermal environmental parameters were analysed using the one-way analysis of variance. Data analyses were done using GraphPad Prism (Version 5.3).

RESULTS

Table 1 shows the thermal environmental parameters of the study site during the hot-dry season. The dry-bulb temperature (DBT) increased

significantly ($P < 0.05$) from 22.6 ± 1.23 °C at 06.00 h (6 am) to 38.6 ± 6.53 °C at 12.00 h (12 pm). There was a significant ($P < 0.05$) increase in relative humidity from 64.4 ± 2.34 % at 06.00 h (6 am) to 74.3 ± 6.73 % at 12.00 h (12 pm). The

temperature-humidity index (THI) increased significantly ($P < 0.05$) from 76.41 ± 0.56 at 06.00 h (6 am) to 83.36 ± 4.53 at 12.00 h (12 pm) when it attained its peak.

Table 1. Thermal environmental parameters of the experimental site during the hot-humid season

Time of Day	Dry-Bulb Temperature (°C)	Relative Humidity (%)	Temperature-Humidity Index
06.00	22.6 ± 1.23^a (22 - 24)	64.4 ± 2.34^a (63 - 68)	76.41 ± 0.56^a (68.71 - 83.65)
12.00	38.6 ± 6.53^b (37-39)	74.3 ± 6.73^b (72 - 78)	83.36 ± 4.53^b (81.32 - 89.01)
18.00	36.5 ± 0.17 (36-37)	78.8 ± 5.98 (76 - 81)	83.24 ± 3.49 (83.21 - 84.95)
Overall Mean \pm SEM	37.22 ± 4.17 (36.11 - 37.33)	75.19 ± 5.98 (73.65 - 81.11)	81.45 ± 5.18 (78.11-84.76)

^{a,b}Means for the same column having different superscript letters are significantly ($P < 0.05$) different Values in Parentheses are Minimum - Maximum

Table 2 shows the biochemical response of the stallions. All biochemical parameters measured were significantly lower ($P < 0.05$) in the treated group than in the untreated group.

Figure 1 shows heart rate responses of stallions before exercise, immediately

after exercise, and 1 hour after exercise. There was a significant difference ($P < 0.05$) between the treated group and the untreated group immediately after exercise with the treated group having a higher heart rate.

Table 2. Biochemical parameters of stallions subjected to exercise in the hot-humid season following administration of ergothioneine

Parameters		Treated (n = 6)	Not-treated (n = 6)
Aspartate aminotransferase U/L	Pre-Exercise	124.60 ± 10.92	137.43 ± 10.16
	Post-Exercise	183.4 ± 9.33 ^a	230.14 ± 17.02 ^b
Lactate dehydrogenase U/L	Pre-Exercise	316.0 ± 18.25	314.23 ± 21.97
	Post-Exercise	365.9 ± 10.09 ^a	457.13 ± 21.75 ^b
Creatinine µmol/L	Pre-Exercise	40.57 ± 5.37	47.64 ± 6.86
	Post-Exercise	43.27 ± 1.43 ^a	53.07 ± 7.87 ^b
Superoxide dismutase u/ml	Pre-Exercise	0.82 ± 0.21	1.77 ± 0.29 ¹
	Post-Exercise	1.69 ± 0.04 ^a	4.89 ± 2.63 ^{b2}
Catalase u/ml	Pre-Exercise	4.97 ± 0.23	6.93 ± 0.76 ¹
	Post-Exercise	5.24 ± 0.13 ^a	12.54 ± 5.14 ^{b2}
Glutathione peroxidase U/L	Pre-Exercise	259.90 ± 6.32	262.7 ± 7.02 ¹
	Post-Exercise	288.0 ± 8.43 ^a	393.0 ± 17.82 ^{b2}
Malondialdehyde µmol/L	Pre-Exercise	5.61 ± 0.34	5.23 ± 0.14 ¹
	Post-Exercise	7.26 ± 0.56 ^a	11.53 ± 5.83 ^b

^{1,2}Means for the same column having different superscript numbers are significantly (P < 0.05) different.

^{a,b}Means for the same row having different superscript letters are significantly (P < 0.05) different




Figure 1. Heart rate of stallions subjected to exercise during the hot-humid season

The respiratory rates of the stallions as shown in Figure 2 show that the respiratory rate of the treated group was significantly higher ($P < 0.05$) than

the untreated group immediately after exercise. The rectal temperature of the untreated group was higher ($P < 0.05$) than the treated group (Figure 3).




Figure 2. Respiratory rate of stallions subjected to exercise during the hot-humid season




Figure 3. Rectal temperature of stallions subjected to exercise during the hot-humid season

DISCUSSION

The thermal environmental parameters recorded during the study period were characterised by high THI, AT, and RH, typical of the hot-humid season of the Southern Guinea Savannah zone of Nigeria. The mean AT values recorded during the study period was higher than the established thermoneutral zones of 5–25 °C for stallions (Kohn *et al.* 1999; Wallsten *et al.*, 2012) The mean 74.3 ± 0.73 and 78.8 ± 0.77 % relative humidity recorded at 12.00 h and 18.00 h are clearly above the 70 % recommended for stallions (Schrama *et al.*,1996; Wallsten *et al.*, 2012). High ambient temperature and high relative humidity are the most important thermal environmental indices, causing heat stress in stallions (Nienaber *et al.* 1999). The heat generated during exercise

which is a by-product of metabolic production of energy may accumulate when ambient temperatures and relative humidity are high. The hot-humid environment in this study imposed additional heat stress on stallions during exercise resulting in the inability of the horses to dissipate heat effectively (Hogson *et al.*, 1993; Hodgson *et al.*, 1994).

The high THI (76.41 ± 0.24 to 83.24 ± 0.49) recorded in this present study is higher than the accepted value of 68 recommended for horses (CIGR 1984; Noordhuizen. and Noordhuizen2017) indicating that meteorological conditions prevailing in the study area were unfavourable for horse performance. Therefore, measures and supplements aimed at alleviating the effects of high THI are necessary to reduce the risks of heat stress and enhance performance.

Rectal temperature is the most accurate method of evaluating the core body temperature and is considered an important index of the body's heat load (Geor and Mccutcheon, 1998). Lower rectal temperature post-exercise in the ergothioneine treated group indicates a better heat dissipation and less heat load, compared to the untreated group after exercise. Supplementation with ergothioneine before exercise suggests that all organs and tissues involved in heat dissipation were protected from the negative effects of exercise stress, thereby reducing the risk of exercise-induced hyperthermia, rhabdomyolysis, and heat stroke (Hodgson *et al.*, 1994).

The finding that the heart rate of the stallions administered with EGT was higher although not statistically significant immediately after exercise than in the control group may indicate the modulatory effects of EGT. Ergothioneine has been reported to be a potent antioxidant *in vivo* with cytoprotective effects on organs and tissues prone to oxidative damage and inflammation (Tamai *et al.*, 1997; Lawan *et al.*, 2010). Exercise induces increased oxygen consumption which affects the oxidant/antioxidant status. Increased oxidants cause an imbalanced equilibrium between oxidants and antioxidants, resulting in oxidative stress, bradycardia, and reduced heart rate (Williams 2016; Martin *et al.*, 2000). This study has established that EGT may be a potent antioxidant *in vivo*. Halliwell *et al.* (2018) reported an avid

uptake of EGT via a specific carrier solute carrier 22 family member 4 (SLC22A4) by organs challenged by oxidative stress. The higher mean respiratory rates in the treated group indicate a more effective function by the respiratory system of the treated group. Unlike many species, horses have a high metabolic rate but a small surface area for dissipation of heat (Franklin *et al.*, 2010). The respiratory system plays a very important role in exercise and the dissipation of heat and must function effectively. The evaporative heat loss from the respiratory system is considered one of the important mechanisms for maintaining heat balance (Rensisand Scaramuzzi, 2003; Franklin *et al.*, 2010). The increased respiratory response recorded in this study may be due to the stimulation of peripheral receptors that transmit impulses to the thermal centre in the hypothalamus by ergothioneine which has been reported to cross the blood-brain barrier. This may result in activation of the respiratory centre to send impulses to the diaphragm and intercostal muscles resulting in increased respiratory rates (Evans and Rose, 1988). Ergothioneine has also been reported to accumulate in the mitochondria and lungs which are sites that are very prone to oxidative imbalance (Repine and Elkins, 2012). Horses are obligate nasal breathers, so the integrity of their respiratory system is paramount for their well-being during and post-exercise. The higher respiratory rate recorded after exercise

in the treated group suggests a modulatory role by EGT. Our findings agree with the work of Repine and Elkins (2012) who reported that EGT is a potent antioxidant and anti-inflammatory agent in the respiratory system.

The activities of aspartate aminotransferase (AST) and lactate dehydrogenase (LH) were lower in the treated group than in the untreated group after exercise. AST and LH have very high activities in the skeletal muscles and the elevation of their activities in the serum suggests damage to the muscles (Harris *et al.*, 1998; Verdegaaland Franklin, 2015). The lower activities of these enzymes recorded in the treated group suggest lesser damage to the skeletal muscle due to the stress of exercise and the functional protective effects of ergothioneine. This result agrees with the report of Cheah *et al.* (2017) who demonstrated that the administration of ergothioneine in human subjects of good health status reduced oxidative and inflammatory damage to the skeletal muscle.

Creatinine, a byproduct of muscle metabolism, was lower in the treated group suggesting protection of muscles from damage by ergothioneine. Following intense exercise and muscle damage, creatinine was released from the muscles, increasing serum creatinine levels (Harris *et al.*, 1991; Lai *et al.*, 2016). This finding agrees with the work of Joseph *et al.* (2012), who reported an increased creatinine level

following exercise in a hot environment in individuals not administered with any supplement.

The serum activities of catalase, glutathione peroxidase, and superoxide dismutase which are defense enzymes against oxidative stress were higher in the untreated group than in the treated group (Hanzawa *et al.*, 2002; Younus, 2018). The results demonstrated lower oxidative stress in the treated group than in the untreated group suggesting effective protection by ergothioneine against ROS.

The serum concentration of malondialdehyde, a by-product of lipid peroxidation was higher in the untreated group than in the treated group post-exercise. Lipid peroxidation occurs when ROS overwhelms and reacts with lipids containing double carbon bonds, particularly polyunsaturated fatty acids (Ayala *et al.*, 2014). Under normal peroxidation rates, the cells stimulate their maintenance through highly organized antioxidant defense systems that up-regulate antioxidants, resulting in an adaptive stress response (Khoubnasabjafari *et al.*, 2015). However, when lipid peroxidation rates are high, the degree of oxidative insults overwhelms antioxidant ability, and the cells undergo apoptosis or necrosis programmed-cell death (Hodgson *et al.*, 2014). It was, therefore, concluded that EGT exerted modulatory roles on biochemical parameters and lowered rectal temperatures of Arabian stallions

following exercise in a hot-humid environment and may be useful in preventing hyperthermia in stallions following exercises in a hot-humid environment.

REFERENCES

- Adah, A. S., D.A., Adah, C.O., Nwonuma and J.O. Aiyedun. 2020. Modulatory effects of melatonin on erythrocyte indices in Arabian stallions following a 30-km sub-maximal exercise. *Comp ClinPathol.* 29:921-926.
- Aubert A.E., B. Seps and F. Beckers. 2003. Heart rate variability in athletes. *Oct; Sports Med.* 33:889-919.
- Ayala A., M.F., Munoz, and S. Argüelles 2014 Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde.; *Oxi Med Cell Longev*, doi: 10.1155/2014/360438
- Bashir A, and A. Rasedee 2009. Plasma catecholamine, sweat electrolytes and physiological responses of exercised normal, partial, anhidrotic and anhidrotic horses.; *Am. J. Anim. Vet. Sci.* 4(1):26-31.
- Cheah I.K., R. Tang, P. Ye, T.S.Z. Yew, K.H.S. Lim and B. Halliwell. 2016. Liver ergothioneine accumulation in a guinea pig model of non-alcoholic fatty liver disease. A possible mechanism of defense? *Free Rad Res*, 50(1):14-25.
- Cheah I.K., R.M.Y. Tang, T.S.Z. Yew., H.H.C. Lim and B. Halliwell. 2017. Administration of pure ergothioneine to healthy human subjects: uptake, metabolism, and effects on biomarkers of oxidative damage and inflammation. *Antioxid. Redox. Sign.*; 26(5):193-206.
- CIGR (1984). Report of Working Group of Climatization of Animal Houses, Scottish Farm Buildings Investigation Unit, Craibstone, Bucksburn, Aberdeen, AB2 9TR, Scotland for Commission Internationale du Genie Rural (International Commission of Agricultural Engineering).
- Cottin F, Barrey E, Lopes P. and Billat VL. 2006. Effect of fatigue during five successive heats (800 m at high velocity) and recovery runs on heart rate variability in Standardbred. *Proceeding of the 7th International Conference on Equine Exercise Physiology*, Fontainebleau, France, pp: 68.
- Courouce, A, Corde R, Valette JP, Cassiat G, Hodgson DR and Rose RJ. 2000. Comparison of some responses to exercise on the track and the treadmill in French trotters: determination of the optimal treadmill incline; *Vet J.* 159: 57-63.
- Franklin, S.H., E. Van Erck-Westergren, and W.M. Bayly. 2012. Respiratory responses to exercise in the horse. *Equine Vet. J.*; 44:726-732.
- Geor, R.J. and L.J. McCutcheon. 1998. Thermoregulatory adaptations associated with training and heat acclimation. *Vet. Clin. N Am-Equine*; 14:97-120.

- Gerard, M.P., S. de Graaf, E., Roelfsema, D.R., Hodgson and J.H. van der Kolk. 2014. Energetic considerations of exercise. In: D.R. Hodgson, McKeever K.H. and McGowan C.M. (Eds). *The athletic horse, principles and practice of equine sports medicine* 2. ed. USA: Elsevier Saunders; p 19-33
- Grigat S, S. Harlfinger, S. Pal, R. Striebinger, S. Golz A. Geerts, A. Lazar, E. Schomig and D. Gründemann. 2007. Probing the substrate specificity of the EGT transporter with methimazole, mercynine, and organic cations. *Biochem Pharmacol*; 74:309-316.
- Halliwell B, I.K. Cheah and R.M.Y., Tang. 2018. EGT a diet-derived antioxidant with therapeutic Potential. *FEBS Lett*; 592: 3357-3366
- Hanzawa, K, A. Hiraga, Y. Yoshida, H. Hara, M. Kai, K. Kubo and S. Watanabe. 2002. Effect of exercise on plasma haptoglobin composition in control and splenectomized Thoroughbred horses; *J Equine Sci*13(3): 89-92.
- Harris, P. A., D.J. Marlin and J. Gray. 1998. Plasma aspartate aminotransferase and creatine kinase activities in thoroughbred racehorses in relation to age, sex, exercise and training.; *Vet J* 155(3): 295-304.
- Harris, R.C., D.J. Marlin D.H. Snow, and R.A. Harkness. 1991 Muscle ATP loss and lactate accumulation at different work intensities in exercising Thoroughbred horse.; *Euro J ApplPhysiol*62: 235-244.
- Hartmann, E, R.J. Hopkins, C. vonBromssen and K. Dahlborn.2015. 24-h sheltering behaviour of individually kept horses during Swedish summer weather; *Acta Vet Scand* 57: 45-54.
- Hodgson, D.R., K.H. McKeever and C.M. McGowan. 2014. *The Athletic Horse: Principles and Practice of Equine Sports Medicine*. 2nd Ed. Elsevier Saunders, China. p 497.
- Hodgson, D.R., R.E. Davis and E.F. McConaghy. 1994 Thermoregulation in the horse in response to exercise; *Br. Vet. J.* 150: 219-234.
- Hodgson, D.R., LJ McCutcheon, S.K. Byrd, W.S. Brown, W.M. Bayly, G.L. Brengelmann and P.D. Gollnick. 1993. Dissipation of metabolic heat in the horse during exercise; *J. Appl. Physiol.* 74:1161-1170.
- Joseph, M.R., G.M. Joy, N.I.N.A. Anika, F.U. Di, and A. Dolci. 2013. Exercising in a hot environment with muscle damage: effects on acute kidney injury biomarkers and kidney function; *Am J Physiol: Renal Physio* 305: F813-F820.
- Kenneth, W.H, J.K. Andris, J.G. Raymond and B. Warwick. 2004. *Equine Sports Medicine and Surgery: Basic and Clinical Sciences of the Equine Athlete.*; 4: 697-728.
- Kerley, R.N., C. McCarthy, D.B. Kell and L.C. Kenny. 2018. The potential therapeutic effects of ergothioneine in pre-eclampsia. *Free Rad Biol and Med.* 117:45-157
- Khoubnasabjafari, M, K.Ansarin and A. Jouyban. 2015. Reliability of

- malondialdehyde as a biomarker of oxidative stress in psychological disorders. *BioImpacts*. 5(3): 123-127.
- Kohn, C.W., K. Hinchcliff and K.H.McKeevert. 1999. Effect of ambient temperature and humidity on pulmonary artery temperature of exercising horses. *Equine Vet. J. Suppl.*; 30:404-41
- Lai, L, R.B. Vega, D.P. Kelly, S.R. Smith, and Z. Gan. 2016. Exercise inducible lactate dehydrogenase B regulates mitochondrial function in skeletal muscle. *J. Biol. Chem.* 291(49):25306-25318.
- Lawan, A., M.A. Noraniza, A. Rasedee, and A. Bashir. 2010. 2-1-Effects of race distance on physical, hematological and biochemical parameters of endurance horses. *Am. J. Anim. Vet. Sci*; 5(4): 244-248.
- Mader, T.L., M.S. Davis and T. Brown-Brandl . 2006. Environmental factors influencing heat stress in feedlot cattle. *J. Anim. Sci.* 84: 712-719.
- Marone PA, J. Trampota and S. A. Weisman 2016. Safety evaluation of a nature-identical L-ergothioneine in Sprague-Dawley rats. *Int. J. Toxicol.* 2016 Sep; 35(5): 568-583
- Marques, M.S., W.R. Fernandes, C.S. Coelho, and R.M.S. Mirandola. 2002. Influence of physical exercise on plasma lactate and serum cortisol in race horses (in Portuguese). *A Hora Veterinaria*; 22: 29-32.
- Martin, B.B. V.B. Reef, E.J. Parente and A.D. Sage. 2000. Causes of poor performance of horses during training, racing, or showing: 348 cases (1992-1996). *J. Am. Vet. Med. Assoc.*; 216: 554-558.
- McMiken DF. An energetic basis of equine performance. *Equine Vet.* 1983; J. 15: 123-133
- Morgan, K. 1998. Thermoneutral zone and critical temperatures of horses. *J. Therm. Biol.*; 23(1): 59-61.
- Morris, E.A. and H.J. Seeherman. Clinical evaluation of poor performance in the racehorse: the results of 275 evaluations. *Equine Vet. J.* 1991; 23: 169-174.
- Nakamura T., K. Yoshida, H. Yabuuchi, T. Maeda and I. Tamai. 2008. Functional characterization of EGT transport by rat organic cation/carnitine transporter OCTN1 (SLC22A4). *Biol. Pharm. Bull*; 31:1580-1584
- Nardone, A, B. Ronchi, N. Lacetera, M.S. Ranieri, and U. Bernabucci. 2010. Effects of climate changes on animal production and sustainability of livestock systems. *Livest. Sci.*; 130(1-3): 57-69
- Nienaber, J.A., G.L. Hahn and R.A. Eigenberg. 1999. Quantifying livestock responses for heat stress management: a review. *Int. J. Biometeorol.*; 42:183-188.
- Noordhuizen, J. and T. Noordhuizen. 2012. Heat Stress in (Sport) Horses: (I) Occurrence, Signs and Diagnosis, (II) Practical Management and Preventive Measures. *J. Dairy Vet. Sci.*; 2(5): 1-5.
- Padilla L, T. Matsui, Y. Kamiya, M. Kamiya and M. Tanaka. 2006. Heat stress decreases plasma vitamin C

- concentration in lactating cows. *Livest. Sci*; 101(1-3): 300-304.
- Paul, B.D. and S.H. Snyder. 2009. The unusual amino acid L-ergothioneine is a physiologic cytoprotectant. *Cell Death Differ*; 17: 1-7.
- Rensis, F.D. and RJ Scaramuzzi. 2003. Heat stress and seasonal effects on reproduction in the dairy cow-a review. *Theriogenology*; 60(6): 1139-1151. ,mk,m,687.
- Repine, J.D., and N.D. Elkins. 2012. Effect of ergothioneine on acute lung injury and inflammation in cytokine insufflated rats. *Prev. Med*; 54(Suppl): S79-S82.
- Richard, E.A., D.F. Guillaume, P.H. Pitel, M.C. Dupuis, J.P. Valette, T. Art, J.H. Denoix, P.M. Lekeux, and E. Van Erck1999. Sub-clinical diseases affecting performance in Standardbred trotters: diagnostic methods and predictive parameters. *Vet. J*;184: 282-289.
- Schmidt, A, J. Aurich, E. Moestl, J. Muller and C. Aurich. 2010. Changes in cortisol release and heart rate and heart rate variability during the initial training of 3-year-old sport horses. *Horm. Behav*; 58: 628-636.
- Schrama, J.W., W. Vander Hel, J. Gorssen, A.M. Henken and M.W.A. Verstegen. 1996. Required thermal thresholds during transport of animals. *Vet. Quart.*; 18(3):90-95
- Tamai, I, H. Yabuuchi, J. Nezu, Y. Sai, A. Oku, M. Shimane and A. Tsuji. 1997. Cloning and characterization of a novel human pH-dependent organic cation transporter, OCTN1. *FEBS Lett*; 419: 107-111.
- Tang, R.M., I. K. Cheah, T.S. Yew and B. Halliwell 2018. Distribution and accumulation of dietary ergothioneine and its metabolites in mouse tissues. *Sci. Rep*; 8: 1601-1615
- Verdegaal E.J.M.F and S.H. Franklin. Science in brief: Highlights from the applied physiology, muscle and genetics abstracts at the International Conference on Equine Exercise Physiology, *Equine Vet J*. 2015;48(1): 9-12.
- Wallsten, H, K. Olsson and K. Dahlborn, 2012. Temperature regulation in horses during exercise and recovery in a cool environment. *Acta Vet. Scand*. 54:42-47.
- Williams, C.A. 2016 The effect of oxidative stress during exercise in the horse. *J. Anim. Sci*; 94:4067-4075.
- Yoshida, S, Shime, H., Funami, K., Takaki, H., Matsumoto, M. and Kasahara, M. 2017. The Antioxidant ergothioneine augments the immunomodulatory function of TLR agonists by direct action on macrophages. *PLoS ONE*; 12(1): e0169360.
- Younus, H. 2018. Therapeutic potentials of superoxide dismutase. *Int J Health Sci (Qassim)*; 12(3): 88-93.