SYSTEMATIC REVIEW

Differences in factors associated with postpartum hemorrhage in developed and developing countries: A systematic review

Sri Astuti Handayani®*, Farida Kartini®

Midwifery Master Program, Faculty of Health Science, Universitas 'Aisyiyah, Yogyakarta, Indonesia.

Article Info	ABSTRACT
Received Dec 6, 2024	Objective: This study aimed to assess maternal and fetal risk factors for post-
Revised Mar 20, 2025	partum hemorrhage in both developed and developing countries.
Accepted Apr 25, 2025	Materials and Methods: A systematic review methodology was employed,
Published Aug 1, 2025	incorporating a descriptive design. The study followed Joanna Briggs Institute
	(JBI) techniques, the PCC framework, and the PRISMA-ScR checklist. Article
*Corresponding author:	searches were conducted across four databases: Research Rabbit, Science Direct,
Sri Astuti Handayani	Wiley, and PubMed. Articles published in English between 2019 and 2024 that
sriastuti6115@gmail.com	met the inclusion criteria were selected. The search terms included combinations
77 1	of "Postpartum women" OR "Puerperium" AND "Risk factors" OR "Risky
Keywords:	conditions" AND "Postpartum hemorrhag*" "Postpartum bleeding" and
Developed countries	"Developed countr*" AND "Developing countr*".
Developing countries	Results: Out of 906 articles, 20 met the inclusion criteria, representing both
Maternal health	developed and developing countries. The findings were categorized into two
Postpartum hemorrhage	themes: risk factors in developed versus developing countries. In developed
Puerperium	countries, PPH risk factors are often linked to medical interventions and
Risk factors	prolonged labor management, while in developing countries, the focus shifts to
	conditions like anemia, uterine overdistension, and trauma from delivery
	practices. Sociocultural disparities further exacerbate risks in both settings.
	Conclusion : The study reveals significant differences in PPH risk factors between
	developed and developing countries. By reviewing existing research, it identifies
	factors that can influence PPH occurrence. The findings emphasize the need for
	developing policies to screen pregnant, maternity, and postpartum women to
	mitigate PPH risk.

Copyright: © 2025 Majalah Obstetri & Ginekologi. pISSN:0854-0381 eISSN:2598-1013 This is an open-access article distributed under the terms of the Creative Commons Attribution License as stated in https://creativecommons.org/licenses/by-nc-sa/4.0/deed.id



How to cite: Handayani SA, Kartini F. Differences in factors associated with postpartum hemorrhage in developed and developing countries: A systematic review. Majalah Obstetri & Ginekologi (Journal of Obstetrics & Gynecology Science). 2025;33(2):141-151. doi: 10.20473/mog.V33I22025.141-151.

Highlights:

- 1. Analysis of maternal and fetal risk factors associated with postpartum hemorrhage, comparing data from both developed and developing countries.
- 2. There are differences in the risk factors for postpartum hemorrhage between developed and developing countries.



INTRODUCTION

Maternal health has been a global concern. In some countries, especially developed and developing countries, mothers face various risks during childbirth. This situation has prompted the international community to make greater efforts in addressing this issue. According to estimates from the World Health Organization (WHO), 287.000 women will pass away in 2020 as a result of pregnancy and delivery. By 2020, problems from pregnancy and childbirth will kill over 800 women daily. In 2020, rich countries accounted for about 95% of all maternal deaths; the majority of these deaths might have been avoided.²

The Maternal Mortality Ratio (MMR) in low-income countries in 2020 was 430 per 100.000 live births versus 13 per 100 000 live births in high income countries. Indonesia has a comparatively high MMR of 305 per 100,000 live births, whereas the IMR is 16.9 per 1,000 live births, according to data from the Central Bureau of Statistics. It falls short of the 2020–2024 National Medium-Term Development Plan (RPJMN) MMR and Infant Mortality Rate (IMR) objectives, which are 183 per 100,000 live births and 16 per 1,000 live births, respectively. Among Southeast Asian countries, Indonesia's MMR remains in the third-highest ranking.

Complications that arise during or after pregnancy and childbirth can have devastating consequences for women. Pregnancy is the time when many of these problems first appear, but they are frequently avoidable or treatable with the right care. If left untreated, several pre-existing diseases can also get worse during pregnancy. Roughly 75% of maternal deaths are caused by high blood pressure conditions like pre-eclampsia and eclampsia, infections that usually occur after childbirth, severe bleeding, especially postpartum hemorrhage (PPH), delivery complications, and unsafe abortion practices.²

The standard definition of PPH is blood loss of more than 500 milliliters after vaginal delivery or more than 1000 milliliters after cesarean delivery. 6 However, in 2017, PPH was redefined by the American College of Obstetrics and Gynecologists (ACOG) as a blood loss associated with hypovolemia symptoms or indicators, as well as a cumulative blood loss of 1000 milliliters or more, regardless of the delivery method, within the first 24 hours of the birth. Clinical signs and symptoms typical of hypovolemia resulting from postpartum hemorrhage, symptoms like tachycardia hypotension could not show up until blood loss exceeds 25% of the total volume of blood (>1500 ml in late pregnancy).⁸ Maternal mortality rates remain a significant challenge, with PPH being a leading cause.

The complexity of contributing factors complicates the issue. Currently, there are no review articles comparing the risk factors for postpartum hemorrhage in developed and developing countries. The purpose of this systematic review is to distinguish between postpartum hemorrhage risk factors in developed and developing countries.

MATERIALS AND METHODS

A systematic review maps and identifies research gaps by following Arkshey & O'Malley and Daniel Levac's methodology, which includes formulating research questions, selecting articles based on inclusion and exclusion criteria, charting data, and extracting relevant information.

Identifying research questions

Usage of the PCC framework is recommended (Population, Concept, Context) to establish specific objectives and prerequisites for a systematic study (<u>Table 1</u>). This will enable the organized identification of pertinent data items.⁹ This review addresses the research question: "What are the differences in factors contributing to PPH in developed versus developing countries?"

Table 1. PCC framework

P	С	С
(Population)	(Concept)	(Context)
Postpartum	The factor which	Developed countries,
women	influence PPH	developing countries

Identifying relevant articles

In article searches for this systematic review, specific inclusion and exclusion criteria are established to streamline the selection process (<u>Table 2</u>).

Table 2. Eligibility criteria

"Inclusion Criteria"	"Exclusion Criteria"
 Original research article 	 Review/ comment
2. Articles published in the last 5	articles
years (2019-2024)	Book chapters
3. Research articles published in	
English	
4. Full text	
Factors contributing to PPH	

Literature searching

The researcher employed various search strategies, including truncation and Boolean operators, ¹⁰ to identify relevant articles. Databases used were PubMed, Science



Direct, Wiley Online Library, and manual searches through Research Rabbit. Keywords were chosen to align with inclusion criteria and bibliography: "Postpartum women" OR "Puerperium" AND "Risk factors" OR "Risky condition" AND "Postpartum hemorrhag*" OR "Postpartum bleeding" AND "Developed countr*" AND "Developing countr*".

Article selection

Researchers followed the Arksey & O'Malley framework for article selection, which included database identification, duplication screening, title and abstract review, full-text screening, and categorization for inclusion or exclusion. Documentation was provided using PRISMA flow charts (Figure 1).¹¹




Figure 1. PRISMA flowchart



Based on the search for articles using keywords and inclusion and exclusion criteria in 3 databases and 1 manual research. It was found as many as 906 articles and then duplication checks were carried out. After that, 870 items were eliminated based on a screening of the abstract and title of 902 articles. Twelve papers were eliminated after 32 were vetted for publication in full. Twenty articles were reviewed in this systematic review. The articles are subsequently chosen for critical appraisal to evaluate their quality. In order to assess

articles quality, the Joanna Brigs checklist from the Joana Brigs Institute (JBI) is needed.

Data charting

At this point, all of the chosen articles data were entered into a table and included the following: the article title, the name of the author, the year, the nation, the goal of the study, the study design, the data collection method, the number of participants and sample size, and the results, as shown in Table 3.

Table 3. Data charting

No	Title/author/year	Country	Aim	Study design	Data collection/ participant/Sample size	Results
A1	Risk factors for postpartum hemorrhage in The Northern Province of Rwanda ¹²	Republic of South Africa	To research and simulate primary PPH risk factors	Case Control	Between January 1 and June 30, 2020, information was gathered from medical records for 430 women, 108 of whom had PPH.	The most common risk factors for primary PPH were hemoglobin levels less than 11 g/dL before admission, cultural differences, and multiple pregnancies.
A2	Association of oxytocin augmentation and duration of labour with postpartum hemorrhage 13	Norway	To look into the relationship between the length of labor and the increase of oxytocin for PPH	Cohort	From December 2014 to January 2017, a secondary analysis was carried out on 14 birth-care units located in all four health regions of Norway. In the LaPS study, 4,943 of the 7,277 women satisfied the inclusion requirements.	The interpretation of oxytocin and labor time as predictors of PPH is complicated by the intricacy of clinical practice.
A3	Twin growth discordance and risk of postpartum hemorrhage ¹⁴	China	To determine whether prenatal growth abnormalities and PPH are related	Cohort	All twin pregnant women who had cesarean sections between August 2013 and July 2020 had their obstetric records examined. 3,180 of them were examined.	Rather than FGR, twin growth discordance was linked to a higher incidence of PPH.
A4	Relation of meconium- stained amniotic fluid and postpartum hemorrhage ¹⁵	China	To determine whether MSAF poses a risk for PPH following vaginal birth	Cohort	Medical records of all patients with vaginal deliveries from January 1, 2013, to December 31, 2018, were analyzed. Of 55,197 deliveries, 13,686 were classified in the MSAF group, and 41,511 in the clear AF group.	To ascertain whether MSAF increases the risk of PPH after vaginal delivery
A5	Postpartum hemorrhage and risk for postpartum readmission ¹⁶	The United States of America	To evaluate and to determine risk factors for postpartum readmission	Cohort	The analysis utilized the 2010–2014 Nationwide Readmissions Database (NRD) and included women aged 15–54 years.	The general risk of all- cause readmission shortly after delivery suggests that these women may benefit from closer postpartum surveillance.
A6	Analysis of perinatal outcome of forceps delivery and risk factors of postpartum hemorrhage ¹⁷	China	To exploring the perinatal outcomes of forceps delivery and the risk factors of PPH	Cohort	The study included mothers aged 15 to 54 years with PPH extending up to 60 days. Of 1,520 pregnant women, 1,454 were in the normal vaginal delivery group.	The incidence of PPH is higher in the forceps-assisted delivery group, particularly with occipital posterior position, abnormal coagulation function, soft birth canal tears, and lateral perineal incisions.



A7	Incidence and risk factors for postpartum hemorrhage among transvaginal deliveries at a tertiary perinatal medical facility in Japan ¹⁸	Japan	To assess the incidence of, and risk factors for PPH among transvaginal deliveries	Cohort	Health care providers were trained to collect data and measure PPH. Among 1,920 deliveries from June 2013 to July 2016, 93 cases of PPH were identified.	Risk factors for PPH included the use of ART, pregnancy-induced hypertension (PIH), severe vaginal/perineal lacerations, and macrosomia neonates.
A8	Postpartum hemorrhage in Sub- Saharan Africa—a prospective study in metropolitan Mozambique ¹⁹	Mozambique	To analyze MM, PPH, and associated risk factors	Cohort	Data were extracted from the paper charts of all pregnant women delivering at MCH, between February 2019 and January 2021. Of 8,879 women, 5,011 were local and 3,495 were non-local; these groups were analyzed and compared separately.	For women from nonmetropolitan areas, PPH was strongly associated with correctable risk factors such as prepartum anemia and uterine atony.
A9	Risk factors for postpartum hemorrhage following cesarean delivery ²⁰	Israel	To identify risk factors for PPH following cesarean delivery (CD)	Cohort	Data were collected from departmental electronics between January 2014 and December 2015. Out of 15,564 deliveries, 3,208 women met the inclusion criteria.	In women undergoing cesarean section, the duration of surgery and the number of previous cesarean deliveries are associated with the risk of PPH.
A10	Risk factors and recurrence of cause-specific postpartum hemorrhage ²¹	Norway	To explore and recurrence risks in a subsequent delivery	Cohort	Data from The Medical Birth Registry of Norway and Statistics Norway (1967–2017) identified 277,746 cases of PPH out of 3,003,025 births, representing 9.3% of the total.	Retained placenta was most commonly associated with severe PPH, while a history of cesarean delivery increased the risk of PPH due to dystocia.
A11	Incidence and risk factors of postpartum hemorrhage in China ²²	China	To investigate the incidence and the risk factors of PPH in Chinese women	Cohort	This study, conducted across 14 medical centers in 10 provinces from October 1, 2016, to September 30, 2017, enrolled 99,253 women, including 95,967 with singleton pregnancies and 3,286 with twin pregnancies.	Risk factors for PPH varied slightly with the number of fetuses and the mode of delivery, with placenta previa and placenta accreta being the two major contributors.
A12	Prevalence and risk factors of severe postpartum hemorrhage ²³	China	To determine the prevalence and risk factors of SPPH	Cohort	This study utilized a database from medical records, including all women who gave birth after 28 weeks of gestation from January 2015 to August 2019, totaling 34,178 women.	Risk factors for SPPH included previous cesarean section, history of PPH, predelivery anemia, prolonged labor, placenta previa, placental abruption, and macrosomia.
A13	Risk factors for postpartum hemorrhage and its severe forms with blood loss evaluated objectively ²⁴	Brazil	To identify risk factors related to PPH and severe PPH with blood loss quantified objectively	Cohort	Data were collected through interviews and medical records from February 1, 2015, to March 31, 2016. Of 319 eligible women, 8 declined participation and 41 underwent cesarean sections, resulting in the inclusion of 270 women.	A prolonged second stage of labor, use of forceps, and episiotomy are associated with PPH and should prompt delivery assistants to recognize and treat PPH early.
A14	Postpartum hemorrhage in maternal mothers at Anutapura Public Hospital in Palu ²⁵	Indonesia	To know the risk factors of the incidence of PPH in birth mothers	Case control	This study used secondary data from complete patient medical records in 2017. Out of 50 PPH cases, 27 involved mothers aged below 20 or over 35 years, indicating high risk.	Anemia, parity, and maternal age are risk factors for PPH in childbirth.
A15	Risk factors for postpartum hemorrhage in a tertiary hospital in South-Central	The United States of America	To determine prevalence, risk factors, and causes for PPH	Cohort	Data were extracted from electronic medical records for 30,674 women between October 1, 2015, and September 30, 2020. Among them, 1,869 experienced	Black race, increased BMI, cesarean delivery, and anemia are associated with PPH, with anemia at



	Louisiana ²⁶				РРН.	delivery increasing the risk of severe maternal morbidity in PPH cases.
A16	Prevalence, related factors and maternal outcomes of primary postpartum haemorrhage in governmental hospitals in Kabul-Afghanistan ²⁷	Afghanistan	To determine the prevalence, related factors and maternal outcomes of primary PPH in governmental hospitals	Cohort	This study focused on women who gave birth between August and October 2018. Data were collected using a structured checklist for those with primary PPH. Out of 8,652 women observed, 215 had primary PPH, and 2 died during cesarean section.	Among demographic characteristics of women with primary PPH, education level, being a housewife, and the self-employment status of their husbands were the most common related factors.
A17	Incidence and risk factors for postpartum hemorrhage ²⁸	Greece	To estimate the rate of primary PPH, as well as to investigate the potential risk factors	Case control	Singleton pregnancies complicated by PPH were recorded from January 2015 to December 2021. Out of 8,545 deliveries, 219 cases (2.5%) of PPH were identified.	Most cases of PPH were managed with uterotonic medications. Advanced maternal age, prematurity, and multiparity significantly impacted PPH occurrence.
A18	Risk factors for postpartum hemorrhage in a Thai– Myanmar border community hospital ²⁹	Thailand	To assess the incidence and relevant risk factors for PPH	Cohort	Medical and obstetric records were reviewed to identify a cohort of 4,845 women who delivered at the hospital between 2014 and 2018.	Risk factors for PPH included nulliparity, a history of PPH, BMI ≥ 35 kg/m², manual placenta removal, fetal weight > 4000 g, ethnicity, and cultural beliefs.
A19	Risk factors for postpartum hemorrhage after elective cesarean deliveries for twin pregnancies ^{3.0}	China	To identify the high-risk factors associated with PPH after an elective cesarean delivery of twins	Cohort	The study, conducted at the Women's Hospital, School of Medicine, Zhejiang University from September 2014 to April 2019, enrolled 532 pregnant women. PPH occurred in 70 of these women (13.2%).	Twin pregnancies and elective cesarean deliveries are high-risk factors for PPH, leading to increased PPH rates in women undergoing elective cesarean sections for twin pregnancies.
A20	Factors associated with primary postpartum hemorrhage in elderly women undergoing repeated cesarean deliveries ³¹	China	To analyze the risk factors for primary PPH in elderly parturients undergoing repeat cesarean deliveries.	Cohort	Medical records were analyzed for all patients from May 2011 to March 2020, dividing them into a PPH group (116 women) and a non-PPH group (1,559 women).	The number of cesarean sections, placenta previa, placenta implantation, placenta attachment position, and fetal position are independent risk factors for PPH.

RESULT AND DISCUSSION

The data charting process produced 20 articles from the database search. These articles were categorized based on several characteristics. First, the study characteristics by country revealed that 13 articles came from developing nations, including South Africa, China, Mozambique, Brazil, Indonesia, Afghanistan, and Thailand, while seven articles were from developed countries, such as Norway, the United States, Japan, Israel, and Greece. In terms of article type, 16 of the selected papers were cohort studies, four were casecontrol studies, and none were qualitative. Finally, using theme mapping, the researchers identified two main

topics: Both the risk factors for postpartum hemorrhage in industrialized and underdeveloped nations should be considered.

Risk factors for postpartum hemorrhage in developed countries

One dose-dependent risk factor for PPH has been found as the use of oxytocin for labor augmentation, with studies demonstrating that prolonged administration increases PPH risk. 13 Specifically, a significant retrospective study in the United States found that oxytocin administered for longer than four hours when in spontaneous labor and longer than seven hours when



in labor induction significantly raises the likelihood of PPH.³² Additionally, prolonged labor duration, particularly beyond 16 hours, is strongly associated with a higher incidence of PPH.³³ Severe PPH is also more frequent in cases where active labor exceeds 12 hours,³⁴ and the duration of the third stage of labor (TSL) plays a critical role in postpartum bleeding risk. Bleeding correlates positively with TSL duration, increasing sharply up to 10 minutes and more gradually thereafter, with a notable rise in PPH risk when TSL exceeds 18 minutes.³⁵

Women with a prior diagnosis of PPH are more likely to experience a recurrence in subsequent pregnancies. 16,21 This recurrent risk is often linked to persistent underlying conditions such as uterine atony, coagulation disorders, or trauma from previous deliveries. It may also indicate that initial risk factors were not adequately managed or have worsened in later pregnancies.36 Compounding these issues, challenges in recognizing bleeding signs, particularly in patients with darker skin tones, can delay timely intervention, potentially increasing PPH morbidity.37 Black patients, in particular, face higher rates of severe PPH-related complications, including transfusions, disseminated intravascular coagulation, and hysterectomy, partly due to receiving less extensive treatment for PPH, which exacerbates maternal morbidity and mortality.38

Preterm birth is another factor that raises the likelihood of PPH. Women who deliver prematurely are more susceptible to postpartum bleeding due to the immaturity of the vascular and uterine muscle systems, which can exacerbate uterine atony, a leading cause of PPH. Furthermore, preterm births often involve complex medical interventions, such as cesarean sections, which further elevate the risk of bleeding.³⁹ Larger infant birth weight is similarly associated with an increased risk of PPH. Mothers who deliver infants weighing over 4,000 grams are twice as likely to develop PPH as those delivering infants of average weight, primarily due to excessive uterine distension, which impairs normal uterine contractions. 40 Uterine atony, a primary cause of PPH, often results from the uterus's inability to contract efficiently after the delivery of a large baby.

Another important risk factor for postpartum hemorrhage is cesarean sections; studies show that women who have cesarean deliveries are more likely to have PPH than those who deliver vaginally. The heightened risk is attributed to several factors, including greater trauma to the uterine tissue during surgery, which impairs the uterus's ability to contract effectively after delivery. Complications such as uterine atony, where the uterine muscles fail to contract properly, are more

frequently observed in women following cesarean sections. $\frac{42}{}$

Risk factors for postpartum hemorrhage in developing countries

Uterine atony is what causes PPH, as evidenced by multiple studies emphasizing its high prevalence across various populations. 12,27,43 This condition is marked by inadequate contraction of myometrial cells when endogenous oxytocin is released during childbirth. Inadequate compression of the blood vessels within myometrial fibers leads to improper closure at the placental implantation site, resulting in the inability to stop postpartum bleeding. This is more frequently seen in more pregnancies than in ones that were singletons, primarily due to excessive uterine distension, which impairs contractions and increases the risk of uterine atony. Mechanical factors such as uterine overdistension, which compromise myometrial contractility, contribute to approximately 80% of PPH cases.

Additionally, meconium-stained amniotic fluid (MSAF) has been consistently shown as a significant risk factor for PPH after vaginal delivery. Studies reveal that MSAF, compared to clear amniotic fluid, is linked to a higher risk of moderate and severe PPH. Epidemiological and retrospective studies further confirm meconium contamination as a notable contributor to PPH. Another identified risk factor is the use of forceps during labor, which, although it expedites delivery and reduces labor duration, is associated with an increased risk of PPH due to trauma to vaginal and perineal tissues, potential uterine atony, and the need for episiotomy or larger tears.

Hemoglobin levels below 10 g/dL also show a strong negative correlation with the likelihood of PPH. 19,25,49 Severe prenatal anemia, by impairing uterine contractions during labor, increases PPH risk, as weaker contractions are more common in anemic women and lead to inadequate uterine contraction and subsequent bleeding after delivery. 50 Similarly, women with placenta previa are at a significantly higher risk, facing a 6- to 20-fold increased chance of PPH compared to those without the condition. 22,30,31,49 Placenta previa hinders the uterus's ability to contract efficiently, which is critical for controlling postpartum bleeding, and it is often associated with placenta accreta, further increasing the risk of PPH through disruption of uterine smooth muscle contractions. 22

Furthermore, obesity, specifically a BMI over 35 kg/m², serves as an independent risk factor for both PPH and severe PPH.²⁹ Research shows that women who are overweight or classified as class I obese are more likely



to experience PPH and uterine atony compared to those with a normal BMI. Desity impacts uterine contractility, as high-fat and high-cholesterol diets reduce the expression of contractility-related proteins, thus decreasing contractile activity during labor and elevating PPH risk. Another contributing factor is preeclampsia, which is linked to a higher likelihood of transfusion following twin deliveries and is an independent risk factor for severe PPH. Preeclampsia may exacerbate PPH risk through decreased antithrombin III activity or platelet counts, with severe preeclampsia (SPE) also correlating with increased prenatal plasma D-dimer levels, further linking it to PPH.

Multiparity, particularly in women with higher parity, is another significant risk factor for primary PPH. 27,29 Women with more than four births may experience reduced uterine muscle elasticity, resulting in less effective contractions, thus raising the risk of PPH. 54 Additionally, higher parity increases the likelihood of placenta previa, further elevating PPH risk. 44 Racial and ethnic differences also play a crucial role in PPH incidence, with the Southeast Asian ethnic group identified as a higher-risk population. 12,29 Ethnic disparities in PPH outcomes 55 are influenced by biological, cultural, and socioeconomic factors, impacting treatment options and the overall management of peripartum complications, including peripartum hysterectomy. 56

The strength of this article lies in its selection of studies from various countries, both developed and developing, encompassing a wide range of postpartum hemorrhage risk factors, thus providing a more comprehensive perspective. However, the limitation of this article is the restricted database used, which narrows the scope of the findings obtained.

CONCLUSION

Based on the 20 articles reviewed, 7 came from developed countries and 13 reports came from developing countries. The research findings reveal differences in the risk factors contributing to PPH between developed and developing countries. PPH in developed countries is primarily linked to medical interventions and prolonged labor, while in developing countries, anemia, uterine overdistension, and delivery trauma are key factors, with sociocultural disparities exacerbating risks globally. It is hoped that this systematic review will serve as evidence of previous research, thereby providing a reference for identifying factors that can influence the occurrence of PPH. Given the various risk factors identified, it is hoped that a policy will be

established for screening pregnant, maternity, and postpartum women to reduce the risk of PPH.

DISCLOSURES

Acknowledgment

The Master of Midwifery study program at 'Aisyiyah University in Yogyakarta provided funding for this study. Furthermore, the writer expresses gratitude to all those who contributed to the preparation of this systematic review.

Conflict of interest

Every author has no competing interests.

Funding

There has been no external funding for this research.

Author Contribution

All authors participated in all phases of this research, including planning, collecting and analyzing data, drafting, and approving the manuscript for publication.

REFERENCES

- Gustiani R, Kartini F. Factors affecting postpartum haemorrhage on postpartum mother. Placentum. Jurnal Ilmiah Kesehatan dan Aplikasinya. 2023;11(1):13-22. doi: 10.20961/placentum.v11i1. 55528
- 2. WHO. Maternal Mortality [Internet]. 2024 [Cited 2024 Jul 13]. Available From: https://www.who.int/news-room/fact-sheets/detail/maternal-mortality
- 3. Ministry of Health, Republic of Indonesia. Turunkan angka kematian ibu melalui deteksi dini dengan pemenuhan USG di Puskesmas [Reduction of maternal mortality rate by early detection with USG at health centers] [Internet]. 2023 [Cited 2024 Jul 13]. Available from: https://kemkes.go.id/id/turunkan-angka-kematian-ibu-melalui-deteksi-dinidengan-pemenuhan-usg-di-puskesmas
- 4. Ministry of Health, Republic of Indonesia. Dana Alokasi Khusus (DAK) fisik bidang kesehatan dalam mendukung target penurunan angka kematian ibu dan anak [Physical Special Allocation Fund (DAK) for the health sector to support the target of reducing maternal and child mortality rates]. In: Analisis RKP dan Pembicaraan Pendahuluan APBN [RKP Analysis and Preliminary Discussion of the State Budget]. 2021.



- Available from: https://share.google/sUplaWL0Ybi CbKLfr.
- World Bank. Angka Kematian Ibu Indonesia Ketiga Tertinggi Di Asia Tenggara [Indonesia's Maternal Mortality Rate is the Third Highest in Southeast Asia] [Internet]. 2021. [Cited 2024 Jul 13]. Available from: https://databoks.katadata.co.id/demografi/statistik/3df25d0a2d01f36/angka-kematian-ibu-indonesia-ketiga-tertinggi-di-asia-tenggara
- Abecassis A, Wainstock T, Sheiner E, et al. Risk factors for early postpartum hemorrhage: A retrospective, population-based, cohort analysis. Int J Gynaecol Obstet. 2024;166(2):812-8. doi: 10.10/02/ijgo.15414. Epub 2024 Feb 6. PMID: 38321817.
- Committee on Practice Bulletins-Obstetrics. Practice Bulletin No. 183: Postpartum hemorrhage. Obstet Gynecol. 2017;130(4):e168-e186. doi: 10.1097/AOG.00000000000002351. PMID: 289375 71.
- 8. Bienstock JL, Eke AC, Hueppchen NA. Postpartum hemorrhage. N Engl J Med. 2021;384(17):1635-45. doi: 10.1056/NEJMra1513247.
- 9. Pollock D, Peters MDJ, Khalil H, et al. Recommendations for the extraction, analysis, and presentation of results in scoping reviews. JBI Evid Synth. 2023;21(3):520-32. doi: 10.11124/JBIES-22-00123. PMID: 36081365.
- Bramer WM, de Jonge GB, Rethlefsen ML, et al. A systematic approach to searching: an efficient and complete method to develop literature searches. J Med Libr Assoc. 2018;106(4):531-41. doi: 10.5195/jmla.2018.283. Epub 2018 Oct 1. PMID: 30271302; PMCID: PMC6148622.
- 11. Tricco AC, Lillie E, Zarin W, et al. PRISMA extension for scoping reviews (PRISMA-ScR): Checklist and explanation. Ann Intern Med. 201;169(7):467-73. doi: 10.7326/M18-0850. Epub 2018 Sep 4. PMID: 30178033.
- 12. Bazirete O, Nzayirambaho M, Umubyeyi A, et al. Risk factors for postpartum haemorrhage in the Northern Province of Rwanda: A case control study. PLoS One. 2022;17(2):e0263731. doi: 10.1371/journal.pone.0263731. PMID: 35167600; PMCID: PMC8846539.
- Bernitz S, Betran AP, Gunnes N, et al. Association of oxytocin augmentation and duration of labour with postpartum haemorrhage: A cohort study of nulliparous women. Midwifery. 2023;123:103705. doi: 10.1016/j.midw.2023.103705. Epub 2023 May 9. PMID: 37244235.
- 14. Cao X, Luo Y, Zhou S, et al. Twin growth discordance and risk of postpartum hemorrhage: A Retrospective Cohort Study. Front Med (Lausanne). 2022;9:876411. doi: 10.3389/fmed. 2022.876411. PMID: 35692549; PMCID: PMC91 74790.

- Fang ZJ, Liu HF, Zhang YL, et al. Relation of meconium-stained amniotic fluid and postpartum hemorrhage: a retrospective cohort study. Eur Rev Med Pharmacol Sci. 2020;24(20):10352-10358. doi: 10.26355/eurrev_202010_23384. PMID: 3315 5191.
- 16. Fein A, Wen T, Wright JD, et al. Postpartum hemorrhage and risk for postpartum readmission. J Matern Fetal Neonatal Med. 2021;34(2):187-94. doi: 10.1080/14767058.2019.1601697. Epub 2019 Apr 9. PMID: 30919702; PMCID: PMC7135873.
- 17. Feng J, Zhang X. Analysis of perinatal outcome of forceps delivery and risk factors of postpartum hemorrhage. Altern Ther Health Med. 2024;30 (4):102-7. PMID: 38064601.
- Fukami T, Koga H, Goto M, et al. Incidence and risk factors for postpartum hemorrhage among transvaginal deliveries at a tertiary perinatal medical facility in Japan. PLoS One. 2019;14(1):e0208873. doi: 10.1371/journal.pone.0208873. PMID: 30625154; PMCID: PMC6326562.
- Glenzer MM, Correia M, Nhantumbo V, et al. Postpartum hemorrhage in Sub-Saharan Africa-a prospective study in metropolitan Mozambique. J Thromb Haemost. 2023;21(12):3463-76. doi: 10.1016/j.jtha.2023.09.002. Epub 2023 Sep 13. PMID: 37709148.
- 20. Ashwal E, Bergel Bson R, Aviram A, et al. Risk factors for postpartum hemorrhage following cesarean delivery. J Matern Fetal Neonatal Med. 2022;35(18):3626-30. doi: 10.1080/14767058.20 20.1834533. Epub 2021 Jan 28. PMID: 33508987.
- 21. Linde LE, Rasmussen S, Moster D, et al. Risk factors and recurrence of cause-specific postpartum hemorrhage: A population-based study. PLoS One. 2022;17(10):e0275879. doi: 10.1371/journal.pone. 0275879. PMID: 36240210; PMCID: PMC9565
- Li S, Gao J, Liu J, et al. Incidence and Risk Factors of Postpartum Hemorrhage in China: A Multicenter Retrospective Study. Front Med (Lausanne). 2021;8:673500. doi: 10.3389/fmed.2021.673500. PMID: 34497812; PMCID: PMC8419315.
- Liu CN, Yu FB, Xu YZ, et al. Prevalence and risk factors of severe postpartum hemorrhage: a retrospective cohort study. BMC Pregnancy Childbirth. 2021;21(1):332. doi: 10.1186/s12884-021-03818-1. PMID: 33902475; PMCID: PMC80 77797.
- 24. Borovac-Pinheiro A, Ribeiro FM, Pacagnella RC. Risk Factors for Postpartum Hemorrhage and its Severe Forms with Blood Loss Evaluated Objectively A Prospective Cohort Study. Rev Bras Ginecol Obstet. 2021;43(2):113-8. doi: 10.1055/s-0040-1718439. Epub 2021 Jan 28. PMID: 33511619; PMCID: PMC10183850.



- 25. Nur R, Sarina HS, Patui NS, et al. Postpartum hemorrhage in maternal mothers at Anutapura Public Hospital in Palu, 2017. Gac Sanit. 2021;35 Suppl 2:S148-S151. doi: 10.1016/j.gaceta.2021.10.014. PMID: 34929799.
- Taylor K, Noel E, Chapple AG, et al. Risk factors for postpartum hemorrhage in a tertiary hospital in South-Central Louisiana. J Matern Fetal Neonatal Med. 2022;35(25):7353-9. doi: 10.1080/14767058.
 2021.1948528. Epub 2021 Jul 25. PMID: 34304 671.
- 27. Shahbazi Sighaldeh S, Nazari A, Maasoumi R, et al. Prevalence, related factors and maternal outcomes of primary postpartum haemorrhage in governmental hospitals in Kabul-Afghanistan. BMC Pregnancy Childbirth. 2020;20(1):428. doi: 10.1186/s12884-020-03123-3. PMID: 32723320; PMCID: PMC7390104.
- Mitta K, Tsakiridis I, Dagklis T, et al. Incidence and Risk Factors for Postpartum Hemorrhage: A Case-Control Study in a Tertiary Hospital in Greece. Medicina (Kaunas). 2023;59(6):1151. doi: 10.3390/medicina59061151. PMID: 37374355; PMCID: PMC10303199.
- Thepampan W, Eungapithum N, Tanasombatkul K, et al. Risk Factors for Postpartum Hemorrhage in a Thai-Myanmar Border Community Hospital: A Nested Case-Control Study. Int J Environ Res Public Health. 2021;18(9):4633. doi: 10.3390/ijerph 18094633. PMID: 33925427; PMCID: PMC8123 817.
- 30. Wan X, Zhao W, Zhao L, et al. Risk factors for postpartum hemorrhage after elective cesarean deliveries for twin pregnancies. Ginekol Pol. 2024;95(7):531-5. doi: 10.5603/GP.a2023.0071. Epub 2023 Aug 7. PMID: 37548501.
- Yang Y, He J, Deng N. Factors associated with primary postpartum hemorrhage in elderly women undergoing repeated cesarean deliveries. Int J Womens Health. 2021;13:1261-1267. doi: 10.2147/IJWH.S332020. PMID: 35002333; PMCID: PMC8721015.
- Klemettilä E, Rahkonen L, Nuutila M, et al. Role of oxytocin in prevention of postpartum hemorrhage in unplanned out-of-hospital deliveries treated by emergency medical services. Acta Obstet Gynecol Scand. 2020;99(7):901-908. doi: 10.1111/aogs. 13804. Epub 2020 Jan 28. PMID: 31943125.
- Bernitz S, Betran AP, Gunnes N, et al. Association of oxytocin augmentation and duration of labour with postpartum haemorrhage: A cohort study of nulliparous women. Midwifery. 2023;123:103705. doi: 10.1016/j.midw.2023.103705. Epub 2023 May 9. PMID: 37244235.
- 34. Chikkamath SB, Katageri GM, Mallapur AA, et al. Duration of third stage labour and postpartum blood

- loss: a secondary analysis of the WHO Champion trial data. Reprod Health. 2021;18 (1):230. doi: 10.1186/s12978-021-01284-8. PMID: 34775959; PMCID: PMC8591926.
- 35. Nyfløt LT, Stray-Pedersen B, Forsén L, et al. Duration of labor and the risk of severe postpartum hemorrhage: A case-control study. PLoS One. 2017;12(4):e0175306. doi: 10.1371/journal.pone. 0175306. PMID: 28384337; PMCID: PMC5383
- Gill P, Patel A, Van Hook JW. Uterine Atony. 2023 Jul 4. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2025 Jan—. <u>PMID</u>: 29630290.
- Sekar H, Joash K. Why is ethnicity a risk factor for postpartum haemorrhage? J Obstet Gynaecol. 2024;44(1):2380084. doi: 10.1080/01443615.2024. 2380084. Epub 2024 Jul 17. PMID: 39016305.
- Guan CS, Boyer TM, Darwin KC, et al. Racial disparities in care escalation for postpartum hemorrhage requiring transfusion. Am J Obstet Gynecol MFM. 2023;5(6):100938. doi: 10.1016/j.ajogmf.2023.100938. Epub 2023 Mar 21. PMID: 36948294; PMCID: PMC10481353.
- Wang W, Liao C, Zhang H, et al. Postpartum haemorrhage risk prediction model developed by machine learning algorithms: A single-centre retrospective analysis of clinical data. Clinical and Experimental Obstetrics & Gynecology. 2024;51 (3):60. doi: 10.31083/j.ceog5103060.
- 40. Quezada-Robles A, Quispe-Sarmiento F, Bendezu-Quispe G, et al. Fetal macrosomia and postpartum hemorrhage in Latin American and Caribbean Region: Systematic review and meta-analysis. Rev Bras Ginecol Obstet. 2023;45(11):e706-e723. doi: 10.1055/s-0043-1772597. Epub 2023 Nov 29. PMID: 38029773; PMCID: PMC10686756.
- 41. Gong J, Chen Z, Zhang Y, et al. Risk-factor model for postpartum hemorrhage after cesarean delivery: a retrospective study based on 3498 patients. Sci Rep. 2022;12(1):22100. doi: 10.1038/s41598-022-23636-5. PMID: 36543795; PMCID: PMC9772 352.
- 42. Banu D, Jikria N, Naim J, et al. Comparison of risk factors of postpartum hemorrhage among normal versus cesarean delivery cases at a secondary care center in Naogaon. Int J Reprod Contracept Obstet Gynecol. 2024;13(3):535-9. doi: 10.18203/2320-1770.ijrcog20240451.
- 43. Sultana R, Manzoor S, Humayan S. Primary post-partum hemorrhage: Risk factors, causes and maternal outcome. J Soc Obstet Gynaecol Pak. 2020;10(1):40–6. Avalable from: https://jsogp.net/index.php/jsogp/article/view/320.
- 44. Basir F, Gnd HA, Handayani D, et al. Risk factors for uterine atony in postpartum hemorrhage patients



- at Dr. Mohammad Hoesin General Hospital, Palembang, Indonesia. Bioscientia Medicina: Journal of Biomedicine & Translational Research. 2023;6(17):2891–4. doi: 10.37275/bsm. v6i17.719.
- 45. di Marco G, Bevilacqua E, Passananti E, et al. Multiple pregnancy and the risk of postpartum hemorrhage: retrospective analysis in a tertiary level center of care. Diagnostics (Basel). 2023; 13(3):446. doi: 10.3390/diagnostics13030446. PMID: 36766551; PMCID: PMC9914228.
- Bouchè C, Wiesenfeld U, Ronfani L, et al. Meconium-stained amniotic fluid: a risk factor for postpartum hemorrhage. Ther Clin Risk Manag. 2018;14:1671-5. doi: 10.2147/TCRM.S150049. PMID: 30254448; PMCID: PMC6140737.
- 47. Gallo DM, Romero R, Bosco M, et al. Meconium-stained amniotic fluid. Am J Obstet Gynecol. 2023;228(5S):S1158-S1178. doi: 10.1016/j.ajog. 2022.11.1283. Epub 2023 Apr 1. PMID: 37012128; PMCID: PMC10291742. xxx
- 48. Sivasambu G, Kempaiah SB, Thukral U. Risk factors for post-partum hemorrhage in patients who underwent operative vaginal delivery in a tertiary care center. International Journal of Reproduction Contraception Obstetrics and Gynecology. 2021;10(11):4085-9. doi: 10.18203/2320-1770. ijrcog20214066.
- 49. Liu CN, Yu FB, Xu YZ, et al. Prevalence and risk factors of severe postpartum hemorrhage: A retrospective cohort study. BMC Pregnancy Childbirth. 2021;21(1):332. doi: 10.1186/s12884-021-03818-1. PMID: 33902475; PMCID: PMC8077797.
- 50. Omotayo MO, Abioye AI, Kuyebi M, et al. Prenatal anemia and postpartum hemorrhage risk: A

- systematic review and meta-analysis. J Obstet Gynaecol Res. 2021;47(8):2565-76. doi: 10.1111/jog.14834. Epub 2021 May 17. PMID: 34002432; PMCID: PMC9258034.
- 51. Butwick AJ, Abreo A, Bateman BT, et al. Effect of maternal body mass index on postpartum hemorrhage. Anesthesiology. 2018;128(4):774-83. doi: 10.1097/ALN.0000000000002082. PMID: 29346134; PMCID: PMC5849500.
- 52. Polic A, Curry TL, Louis JM. The impact of obesity on the management and outcomes of postpartum hemorrhage. Am J Perinatol. 2022;39(6):652-7. doi: 10.1055/s-0040-1718574. Epub 2020 Oct 14. PMID: 33053594.
- Shao H, Gao S, Dai D, et al. The association of antenatal D-dimer and fibrinogen with postpartum hemorrhage and intrauterine growth restriction in preeclampsia. BMC Pregnancy Childbirth. 2021; 21(1):605. doi: 10.1186/s12884-021-04082-z. PMID: 34482843; PMCID: PMC8420001.
- 54. Hidayati SN, Budihastuti UR, Widyaningsih V. Path analysis on determinants of postpartum bleeding at Dr. Moewardi Hospital, Central Java. Journal of Maternal and Child Health. 2019;4(6): 35-47. doi: 10.26911/thejmch.2019.04.06.04.
- 55. Dai J, Shi Y, Wu Y, et al. The interaction between age and parity on adverse pregnancy and neonatal outcomes. Front Med (Lausanne). 2023;10:1056 064. doi: 10.3389/fmed.2023.1056064. PMID: 36910494; PMCID: PMC9995429.
- 56. Wang L, Pan JY. Predictive model for postpartum hemorrhage requiring hysterectomy in a minority ethnic region. World J Clin Cases. 2024;12(22): 4865-72. doi: 10.12998/wjcc.v12.i22. 4865. PMID: 39109042; PMCID: PMC11238818.

